



भारत सरकार
GOVERNMENT OF INDIA



शुभो चटर्जी
Subho Chatterjee
जन्म वर्ष / Year of Birth : 1988
पुरुष / Male



6190 3560 5968

आधार — आम आदमी का अधिकार

Subho Chatterjee

SUBURBAN DIAGNOSTIC (I) PVT LTD.
FLAT NO.101 ANAND SAGAR CHS
ABOVE RAJKAMAL SHOP
SECTOR - 17, VASHI,
NAVI MUMBAI - 400703

Date: 08/03/2025

To,
Suburban Diagnostics (India) Private Limited

VASHI CENTER
FLAT NO 101 ANAND SAGAR CHS
ABOVE RAJKAMAL SHOP
SECTOR 17 VASHI
NAVI MUMBAI:- 400703
Phone No:- 022 6170 0000

SUBJECT- TO WHOMSOEVER IT MAY CONCERN

Dear Sir/ Madam,

This is to inform you that I, Myself Mr/ Mrs/ Ms. Susho Chatterjee

don't want to performed the following tests:

- 1) Stool Test
- 2) Phy & Eye check. up
- 3) USG
- 4) _____
- 5) _____

CID No. & Date : 393765797 8/3/2025

Corporate/ TPA/ Insurance Client Name : M/S Medi wheel

Thanking you.

Yours sincerely,

(Mr/Mrs/Ms. Susho Chatterjee)

Age 36 years 4 months 8 days

Gender Male


Heart Rate 88bpm

Patient Vitals

BP: NA
Weight: 92 kg
Height: 178 cm
Pulse: NA
Spo2: NA
Resp: NA
Others:

Measurements

QRSD: 72ms
QT: 330ms
QTcB: 399ms
PR: 136ms
P-R-T: 60° 42° 38°



* ECG Within Normal Limits: Sinus Rhythm. Please correlate clinically.

REPORTED BY

Dasgupta

Dr Anirban Dasgupta
MBBS DNB
Reg. 2005020920

Disclaimer: 1) Analysis in this report is based on ECG alone and should be used as an adjunct to clinical history, symptoms, and results of other invasive and non-invasive tests and must be interpreted by a qualified physician. 2) Patient vitals are not captured by the clinician and not derived from the ECG.

CID : 393765797
Name : Mr. CHATTERJEE SUBHO
Age / Sex : 36 Years/Male
Ref. Dr : self
Reg. Location : Vashi Main Centre
Reg. Date : 08-Mar-2025
Reported : 11-Mar-2025 / 18:53

USG WHOLE ABDOMEN

LIVER:

The liver is normal in size (14.5 cm), shape and smooth margins. It shows **raised** parenchymal echo pattern. The intra hepatic biliary and portal radical appear normal. The main portal vein and CBD appears normal. A **thin walled anechoic cyst measuring 0.9 x 0.6 cm is noted in segment VIII of right lobe of liver suggestive of simple hepatic cyst.**

GALL BLADDER:

The gall bladder is physiologically distended and appears normal. No evidence of gall stones or mass lesions seen.

PANCREAS:

The pancreas is well visualised and appears normal. No evidence of solid or cystic mass lesion.

KIDNEYS:

Both the kidneys are normal in size, shape and echotexture.
No evidence of any calculus, hydronephrosis or mass lesion seen.
Right kidney measures 10.8 x 4.5 cm. Left kidney measures 10.7 x 5.1 cm.

SPLEEN:

The spleen is normal in size (9.7 cm) and echotexture. No evidence of focal lesion is noted.
Gaseous distention of bowel loops is noted.

URINARY BLADDER:

The urinary bladder is well distended and reveal no intraluminal abnormality.


PROSTATE:

The prostate is normal in size, measures 3.2 x 3.3 x 3.2 cm and volume is 18.6 cc.

IMPRESSION:

Grade I - II fatty liver.
Simple hepatic cyst.

-----End of Report-----



Dr. Swapnil Nisal
MBBS, DMRE
MMC Reg. No. 2015/06/3297

Click here to view images <<ImageLink>>

Page no 1 of 1

CID : 393765797
Name : Mr. CHATTERJEE SUBHO
Age / Sex : 36 Years/Male
Ref. Dr : self
Reg. Location : Vashi Main Centre
Reg. Date : 08-Mar-2025
Reported : 10-Mar-2025 / 9:29

X-RAY CHEST PA VIEW

Both lung fields are clear.

Both costo-phrenic angles are clear.

The cardiac size and shape are within normal limits.


The domes of diaphragm are normal in position and outlines.

The skeleton under review appears normal.

IMPRESSION:

NO SIGNIFICANT ABNORMALITY IS DETECTED.

-----End of Report-----



Dr Shilpa Beri
MBBS DMRE
Reg No 2002/05/2302
Consultant Radiologist

[Click here to view images <<ImageLink>>](#)

Page no 1 of 1

NAME :- MR.CHATTERJEE SUBHO	AGE :- 36 YRS	P
SEX :- MALE	DATE :-08/03/2025	O
CID NO :-393765797		R
		T

2D Echo and Colour doppler report

All cardiac chambers are normal in dimension
 No obvious resting regional wall motion abnormalities (RWMA)
 Interatrial and Interventricular septum – Appears Normal
 Valves – Structurally normal
 Good biventricular function.
 IVC appears normal.
 Pericardium is normal.
 Great vessels - Origin and visualized proximal part are normal.
 No coarctation of aorta.

Doppler study

Normal flow across all the valves.
 No pulmonary hypertension.
 No diastolic dysfunction.

Measurements

Aorta annulus	18 mm
Left Atrium	31 mm
LVID(Systole)	18 mm
LVID(Diastole)	36 mm
IVS(Diastole)	10 mm
PW(Diastole)	9 mm
LV ejection fraction.	55-60%

Conclusion


Good biventricular function

No RWMA

Valves – Structurally normal

No diastolic dysfunction

No PAH



Dr. Anirban Dasgupta

MBBS DNB

Reg. No.2005/02/0920

Performed by: Dr. Anirban Dasgupta
D.N.B. Internal Medicine, Diploma Cardiology (PGDCC-IGNOU).

SUBURBAN DIAGNOSTIC (I) PVT LTD.
FLAT NO.101 ANAND SAGAR CHS
ABOVE RAJKAMAL SHOP
SECTOR - 17, VASHI,
NORTH MUMBAI - 400703

Ref By: SELF
 Collected: 8/3/2025 9:37:00AM
 A/c Status: P
 Collected at: WALKIN - VASHI (MAIN CENTRE)
 Shop No. 22, Raikar Bhavan, Near Navaratna
 Hotel, Sector 17, Vashi, Navi Mumbai,
 Maharashtra - 400703

Age: 36 Years
 Gender: Male
 Reported: 8/3/2025 7:40:37PM
 Report Status: Final
 Processed at: SDRL, VIDYAVIHAR

MediWheel Full Body Health Checkup Male >40/2D ECH
PROSTATE SPECIFIC ANTIGEN (PSA)

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
TOTAL PSA, Serum	0.46	<4.00 ng/mL	CLIA

Clinical Significance:

- PSA is detected in the serum of males with normal, benign hyper-plastic, and malignant prostate tissue.
- Monitoring patients with a history of prostate cancer as an early indicator of recurrence and response to treatment.
- Prostate cancer screening 4. The percentage of Free PSA (FPSA) in serum is described as being significantly higher in patients with BPH than in patients with prostate cancer. 5. Calculation of % free PSA (ie. FPSA/TPSA x 100), has been suggested as way of improving the differentiation of BPH and Prostate cancer.

Interpretation:

Increased In- Prostate diseases, Cancer, Prostatitis, Benign prostatic hyperplasia, Prostatic ischemia, Acute urinary retention, Manipulations like Prostatic massage, Cystoscopy, Needle biopsy, Transurethral resection, Digital rectal examination, Radiation therapy, Indwelling catheter, Vigorous bicycle exercise, Drugs (e.g., testosterone), Physiologic fluctuations. Also found in small amounts in other cancers (sweat and salivary glands, breast, colon, lung, ovary) and in Skene glands of female urethra and in term placenta, Acute renal failure, Acute myocardial infarction,
Decreased In- Ejaculation within 24-28 hours, Castration, Antandrogen drugs (e.g., finasteride), Radiation therapy, Prostatectomy, PSA falls 17% in 3 days after lying in hospital, Artifactual (e.g., improper specimen collection; very high PSA levels), Finasteride (5- α -reductase inhibitor) reduces PSA by 50% after 6 months in men without cancer.

Reflex Tests: % FREE PSA, USG Prostate

Limitations:

- PSA values determined on patient samples by different testing procedures cannot be directly compared with one another and could be the cause of erroneous medical interpretations. If there is a change in the tPSA assay procedure used while monitoring therapy, then the tPSA values obtained upon changing over to the new procedure must be confirmed by parallel measurements with both methods. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels.
- Patients who have been regularly exposed to animals or have received immunotherapy or diagnostic procedures utilizing immunoglobulins or immunoglobulin fragments may produce antibodies, e.g. HAMA, that interferes with immunoassays. PSA results should be interpreted in light of the total clinical presentation of the patient, including: symptoms, clinical history, data from additional tests, and other appropriate information.
- Serum PSA concentrations should not be interpreted as absolute evidence for the presence or



Lab No. : 393765797
 Ref By : SELF
 Collected : 8/3/2025 9:37:00AM
 A/c Status : P
 Collected at : WALKIN - VASHI (MAIN CENTRE)
 Shop No. 22, Raikar Bhavan, Near Navaratna
 Hotel, Sector 17, Vashi, Navi Mumbai,
 Maharashtra - 400703

MediWheel Full Body Health Checkup Male >40/2D ECH
PROSTATE SPECIFIC ANTIGEN (PSA)
 RESULTS
 BIOLOGICAL REF RANGE
 METHOD

absence of prostate cancer.
Note : The concentration of PSA in a given specimen, determined with assay from different manufacturers, may not be comparable due to differences in assay methods and reagent specificity.
Reference:
 • Wallach's Interpretation of diagnostic tests
 • Total PSA Pack insert



R
F
P
O
R
T



Ref By: **SELF**
 Collected: **8/3/2025 9:37:00AM**
 A/c Status: **P**
 Collected at: **WALKIN - VASHI (MAIN CENTRE)**
 Shop No. 22, Raikar Bhavan, Near Navarata
 Hotel, Sector 17, Vashi, Navi Mumbai,
 Maharashtra - 400703

Age: **36 Years**
 Gender: **Male**
 Reported: **8/3/2025 7:40:37PM**
 Report Status: **Final**
 Processed at: **SDRL, VIDYAVIHAR**

MediWheel Full Body Health Checkup Male >40/2D ECH
BLOOD GROUPING & Rh TYPING

RESULTS

PARAMETER
 ABO GROUP
 Rh Typing

B
 Positive

NOTE: Test performed by automated Erythrocytes magnetized technology (EMT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

Clinical significance:

ABO system is most important of all blood group in transfusion medicine

Limitations:

- ABO blood group of newborn is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

References:

1. Denise M Harming, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia



Age : 36 Years
 Gender : Male
 Reported : 8/3/2025 7:40:37PM
 Report Status : Final
 Processed at : SDRL, VIDYAVIHAR

MediWheel Full Body Health Checkup Male >40/2D ECH

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
TOTAL PSA, Serum	0.46	<4.00 ng/mL	CLIA

Clinical Significance:

- PSA is detected in the serum of males with normal, benign hyper-plastic, and malignant prostate tissue.
- Monitoring patients with a history of prostate cancer as an early indicator of recurrence and response to treatment.
- Prostate cancer screening 4. The percentage of Free PSA (FPSA) in serum is described as being significantly higher in patients with BPH than in patients with prostate cancer. 5. Calculation of % free PSA (ie. FPSA/TPSA x 100), has been suggested as way of improving the differentiation of BPH and Prostate cancer.

Interpretation:

Increased In-Prostate diseases, Cancer, Prostatitis, Benign prostatic hyperplasia, Prostatic ischemia, Acute urinary retention, Manipulations like Prostatic massage, Cystoscopy, Needle biopsy, Transurethral resection, Digital rectal examination, Radiation therapy, Indwelling catheter, Vigorous bicycle exercise, Drugs (e.g., testosterone), Physiologic fluctuations. Also found in small amounts in other cancers (sweat and salivary glands, breast, colon, lung, ovary) and in Skene glands of female urethra and in term placenta, Acute renal failure, Acute myocardial infarction, Decreased In-Ejaculation within 24-28 hours, Castration, Antiandrogen drugs (e.g., finasteride), Radiation therapy, Prostatectomy, PSA falls 17% in 3 days after lying in hospital, Artifactual (e.g., improper specimen collection; very high PSA levels). Finasteride (5- α -reductase inhibitor) reduces PSA by 50% after 6 months in men without cancer.

Reflex Tests: % FREE PSA, USG Prostate

Limitations:

- tPSA values determined on patient samples by different testing procedures cannot be directly compared with one another and could be the cause of erroneous medical interpretations. If there is a change in the tPSA assay procedure used while monitoring therapy, then the tPSA values obtained upon changing over to the new procedure must be confirmed by parallel measurements with both methods. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels.
- Patients who have been regularly exposed to animals or have received immunotherapy or diagnostic procedures utilizing immunoglobulins or immunoglobulin fragments may produce antibodies, e.g. HAMA, that interfere with immunoassays. PSA results should be interpreted in light of the total clinical presentation of the patient, including: symptoms, clinical history, data from additional tests, and other appropriate information.
- Serum PSA concentrations should not be interpreted as absolute evidence for the presence or



R
E
O
R
T





- Wallach's Interpretation of diagnostic tests
- Total PSA Pack insert

Reference:


Note : The concentration of PSA in a given specimen, determined with assay from different manufacturers, may not be comparable due to differences in assay methods and reagent specificity.
 absence of prostate cancer.

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
PROSTATE SPECIFIC ANTIGEN (PSA)			
MediWHEEL Full Body Health Checkup Male >40/2D ECH			

Name: CHATTERJEE SUBHO
 Lab No.: 393765797
 Ref By: SELF
 Collected: 8/3/2025 9:37:00AM
 A/c Status: P
 Collected at: WALKIN - VASHI (MAIN CENTRE)
 Shop No. 22, Raikar Bhavan, Near Navarata
 Hotel, Sector 17, Vashi, Navi Mumbai,
 Maharashtra - 400703

Age: 36 Years
 Gender: Male
 Reported: 8/3/2025 7:40:37PM
 Report Status: Final
 Processed at: SDRL, VIDYAVIHAR

R
E
P
O
R
T



Ref By: SELF
Collected: 8/3/2025 9:37:00AM
A/c Status: P
Collected at: WALKIN - VASHI (MAIN CENTRE)
Shop No. 22, Raikar Bhavan, Near Navarata
Hotel, Sector 17, Vashi, Navi Mumbai,
Maharashtra - 400703

Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SURL, VIDYAVIHAR

MediWheel Full Body Health Checkup Male >40/2D ECH
BLOOD GROUPING & Rh TYPING

RESULTS

ABO GROUP: B
Rh Typing: Positive

NOTE: Test performed by automated Erythrocytes magnetized technology (EMT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

Clinical significance:

ABO system is most important of all blood group in transfusion medicine

Limitations:

- ABO blood group of new born is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

References:

1. Denise M Harmering, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia




Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SDRL, VIDYAVIHAR

T
R
O
M
R

MediWheel Full Body Health Checkup Male >40/2D ECH
BLOOD GROUPING & Rh TYPING

PARAMETER

Dr Trupti Shetty
MD Pathology
Deputy HOD



Dr Leena Salunkhe
DPB
HOD



Dr Namrata Rauli
MD, Biochemistry
Consultant Biochemist



Dr Priyanka Sunil Pagare
MD Pathology
Sr. Pathologist



Dr Urushali Shroff
MD Pathology
Sr. Pathologist




Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SHREEYASH HOSPITAL LAB, PANVEL

MediWhee Full Body Health Checkup Male >40/2D ECH

GBC (Complete Blood Count), Blood


PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
RBC PARAMETERS			
Haemoglobin	15.3	13.0 - 17.0 g/dL	Spectrophotometric
RBC	5.5	4.5 - 5.5 mil/cmm	Elect. Impedance
PCV	45.5	40.0 - 50.0 %	Calculated
MCV	83.3	81.0 - 101.0 fL	Measured
MCH	28.0	27.0 - 32.0 pg	Calculated
MCHC	33.6	31.5 - 34.5 g/dL	Calculated
RDW	14.1	11.6 - 14.0 %	Calculated
WBC PARAMETERS			
WBC Total Count	7360	4000 - 10000 /cmm	Elect. Impedance
WBC DIFFERENTIAL AND ABSOLUTE COUNTS			
Lymphocytes	31.0	20.0 - 40.0 %	Calculated
Absolute Lymphocytes	2281.6	1000.0 - 3000.0 /cmm	Calculated
Monocytes	7.2	2.0 - 10.0 %	Calculated
Absolute Monocytes	529.9	200.0 - 1000.0 /cmm	Calculated
Neutrophils	58.8	40.0 - 80.0 %	Calculated
Absolute Neutrophils	4327.7	2000.0 - 7000.0 /cmm	Calculated
Eosinophils	2.2	1.0 - 6.0 %	Calculated
Absolute Eosinophils	161.9	20.0 - 500.0 /cmm	Calculated
Basophils	0.8	0.1 - 2.0 %	Calculated
Absolute Basophils	58.9	20.0 - 100.0 /cmm	Calculated
PLATELET PARAMETERS			



MediWheel Full Body Health Checkup Male >40/2D ECH
CBC (Complete Blood Count), Blood

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
Platelet Count	325000	150000 - 410000 /cmm	Elect. Impedance
MPV	10.8	6.0 - 11.0 fL	Measured
PDW	14.3	11.0 - 18.0 %	Calculated
RBC MORPHOLOGY			
Others	Normocytic		
	Normochromic		

Specimen: EDTA whole blood



Name : CHATTERJEE SUBHO
Lab No. : 393765/97
Ref By : SELF

Collected : 8/3/2025 9:37:00AM

A/c Status : P

Collected at : WALKIN - VASHI (MAIN CENTRE)

Shop No. 22, Raikar Bhavan, Near Navaratna
Hotel, Sector 17, Vashi, Navi Mumbai,

Maharashtra - 400703

MediWheel Full Body Health Checkup Male >40/2D ECH
ERYTHROCYTE SEDIMENTATION RATE (ESR)

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
ESR, EDTA WB	8.00	2.00 - 15.00 mm/hr	Sedimentation

Clinical Significance: The erythrocyte sedimentation rate (ESR), also called a sedimentation rate is the rate red blood cells sediment in a period of time.

Interpretation:

Factors that increase ESR: Old age, Pregnancy, Anemia
Factors that decrease ESR: Extreme leukocytosis, Polycythemia, Red cell abnormalities- Sickle cell disease

Limitations:

- It is a non-specific measure of inflammation.
- The use of the ESR as a screening test in asymptomatic persons is limited by its low sensitivity and specificity.

Reflex Test: C-Reactive Protein (CRP) is the recommended test in acute inflammatory conditions.

Reference:


- Pack Insert
- Bridgen ML. Clinical utility of the erythrocyte sedimentation rate. American family physician. 1999 Oct 1;60(5):1443-50.



Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SHREEYASH HOSPITAL LAB, PANVEL

MediWheel Full Body Health Checkup Male >40/2D ECH

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
GLUCOSE (SUGAR) FASTING, Fluoride Plasma Fasting	84.31	Non-Diabetic: < 100 mg/dl Impaired Fasting Glucose: 100-125 mg/dl Diabetic: >= 126 mg/dl	Hexokinase
Note : ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition			
GLUCOSE (SUGAR) PP, Fluoride Plasma PP	101.12	Non-Diabetic: < 140 mg/dl Impaired Glucose Tolerance: 140-199 mg/dl Diabetic: >= 200 mg/dl	Hexokinase
Note : ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition			



Age : 36 Years
 Gender : Male
 Reported : 8/3/2025 7:40:37PM
 Report Status : Final
 Processed at : SHREEYASH HOSPITAL LAB, PANVEL
 Collected at : WALKIN - VASHI (MAIN CENTRE)
 Shop No. 22, Raikar Bhavan, Near Navaratna Hotel,
 Sector 17, Vashi, Navi Mumbai, Maharashtra -
 400703

Medi/Wheel Full Body Health Checkup Male >40/2D ECH

KIDNEY FUNCTION TESTS

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
BLOOD UREA, Serum	16.72	12.80 - 42.80 mg/dL	Urease GLDH
BUN, Serum	7.81	6.00 - 20.00 mg/dL	Calculated
CREATININE, Serum	0.98	0.67 - 1.17 mg/dL	Enzymatic
eGFR, Serum	102.49	(ml/min/1.73sqm) Normal or High: Above 90 Mild decrease: 60-89 Mild to moderate decrease: 45-59 Moderate to severe decrease: 30-44 Severe decrease: 15-29 Kidney failure: <15	Calculated
TOTAL PROTEINS, Serum	6.78	6.40 - 8.30 g/dL	Buret
Albumin Serum	4.52	3.50 - 5.20 g/dL	BCG
GLOBULIN Serum	2.27	2.30 - 3.50 g/dL	Calculated
AVG RATIO Serum	1.99	1.00 - 2.00	Calculated
URIC ACID, Serum	6.10	3.50 - 7.20 mg/dL	Enzymatic
PHOSPHORUS, Serum	3.42	2.70 - 4.50 mg/dL	Molybdate UV
CALCIUM, Serum	9.06	8.60 - 10.00 mg/dL	N-BAPTA
SODIUM, Serum	139.00	135.00 - 148.00 mmol/L	Indirect ISE
POTASSIUM, Serum	4.6	3.50 - 5.30 mmol/L	Indirect ISE
CHLORIDE Serum	107.00	98.00 - 107.00 mmol/L	Indirect ISE

Note: eGFR estimation is calculated using 2021 CKD-EPI GFR equation



T R O P E R

Age : 36 Years
Gender : Male
Reported : 8/3/2025 7:40:37PM
Report Status : Final
Processed at : SHREYASH HOSPITAL LAB, PANVEL
Collected at : WALKIN - VASHI (MAIN CENTRE)
Shop No. 22, Raikar Bhavan, Near Navaratna Hotel,
Sector 17, Vashi, Navi Mumbai, Maharashtra -
400703

MediWHEEL Full Body Health Checkup Male >40/2D ECH
GLYCOSYLATED HEMOGLOBIN (HbA1c)

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
Glycosylated Hemoglobin (HbA1c), EDTA WB	5.3	Non-Diabetic Level: < 5.7 % Prediabetic Level: 5.7-6.4 % Diabetic Level: >= 6.5 %	HPLC
Estimated Average Glucose (eAG), EDTA WB	105.4		Calculated

Intended use:

- In patients who are meeting treatment goals, HbA1c test should be performed at least 2 times a year
- In patients whose therapy has changed or who are not meeting glycemic goals, it should be performed quarterly
- For microvascular disease prevention, the HbA1c goal for non pregnant adults in general is Less than 7%.

Clinical Significance:


- HbA1c, Glycosylated hemoglobin or glycated hemoglobin, is hemoglobin with glucose molecule attached to it.
- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of glycosylated hemoglobin in the blood.

Test Interpretation:

- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of Glycosylated hemoglobin in the blood.
- HbA1c test may be used to screen for and diagnose diabetes or risk of developing diabetes.
- To monitor compliance and long term blood glucose level control in patients with diabetes.
- Index of diabetic control, predicting development and progression of diabetic micro vascular complications.

Factors affecting HbA1c results:

Increased in: High fetal hemoglobin, Chronic renal failure, Iron deficiency anemia, plentectomy, Increased serum triglycerides, Alcohol ingestion, Lead/opiate poisoning and Salicylate treatment.
Decreased in: Shortened RBC lifespan (Hemolytic anemia, blood loss), following transfusions, pregnancy, ingestion of large amount of Vitamin E or Vitamin C and Hemoglobinopathies
Reflex tests: Blood glucose levels, CGM (Continuous Glucose monitoring)
References: ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition.



**R
E
P
O
R
T**

Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SHREEYASH HOSPITAL LAB, PANVEL

R
E
P
O
R
T

MediWHEEL Full Body Health Checkup Male >40/2D ECH

FUS and KETONES

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
Urine Sugar (Fasting)	Absent	Absent	
Urine Ketones (Fasting)	Absent	Absent	



Ref By : SELF

Collected : 08/03/2025 09:37:00AM

A/c Status : P

Collected at : WALKIN - VASHI (MAIN CENTRE)

Shop No. 22, Raikar Bhavan, Near Navaratna Hotel,
Sector 17, Vashi, Navi Mumbai, Maharashtra -
400703

Age : 36 Years
Gender : Male
Reported : 8/3/2025 7:40:37PM
Report Status : Final
Processed at : SHREEYASH HOSPITAL LAB, PANVEL

MediWheel Full Body Health Checkup Male >40/2D ECH
Glucose & Ketones, Urine

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
Urine Ketones (PP)	Absent	Absent	
Urine Sugar (PP)	Absent	Absent	



Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SHREEYASH HOSPITAL LAB, PANVEL

MediWHEEL Full Body Health Checkup Male >40/2D ECH

LIPID PROFILE

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
CHOLESTEROL, Serum	118	Desirable: <200 mg/dl Borderline High: 200-239mg/dl High: >=240 mg/dl	CHOD-POD
TRIGLYCERIDES, Serum	111	Normal: <150 mg/dl Borderline-high: 150 - 199 mg/dl High: 200 - 499 mg/dl Very high: >=500 mg/dl	GPO-POD
HDL CHOLESTEROL Serum	28	Desirable: >60 mg/dl Borderline: 40 - 60 mg/dl Low (High risk): <40 mg/dl	Homogeneous enzymatic colorimetric assay
NON HDL CHOLESTEROL, Serum	90	Desirable: <130 mg/dl Borderline-high: 130 - 159 mg/dl High: 160 - 189 mg/dl Very high: >=190 mg/dl	Calculated
LDL CHOLESTEROL Serum	68	Optimal: <100 mg/dl Near Optimal: 100 - 129 mg/dl Borderline High: 130 - 159 mg/dl High: 160 - 189 mg/dl Very High: >= 190 mg/dl	Calculated
VLDL CHOLESTEROL Serum	22	>= 30 mg/dl	Calculated
CHOL / HDL CHOL RATIO,	4	0-4.5 Ratio	Calculated
LDL CHOL / HDL CHOL RATIO,	2	0-3.5 Ratio	Calculated

Reference:

1) Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III).

2) Pack Insert.



Age : 36 Years
Gender : Male
Reported : 8/3/2025 7:40:37PM
Report Status : Final
Processed at : SHREYASH HOSPITAL LAB, PANVEL
Shop No. 22, Raikar Bhavan, Near Navaratna Hotel,
Sector 17, Vashi, Navi Mumbai, Maharashtra -
400703

MediWHEEL Full Body Health Checkup Male >40/2D ECH

THYROID FUNCTION TESTS

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
Free T3, Serum	5.90	3.50 - 6.50 pmol/L	ECLIA
Free T4 Serum	16.36	11.50 - 22.70 pmol/L	ECLIA
sensitive TSH Serum	5.00	0.35 - 5.50 microIU/ml	ECLIA

Interpretation:


A thyroid panel is used to evaluate thyroid function and/or help diagnose various thyroid disorders.

Clinical Significance:

1. TSH Values between high abnormal upto 15 microIU/ml should be correlated clinically or repeat the test with new sample as physiological factors can give falsely high TSH.
2. TSH values may be transiently altered because of non thyroidal illness like severe infections, liver disease, renal and heart severe burns, trauma and surgery etc.

TSH	FT4 / T4	FT3 / T3	Interpretation
High	Normal	Normal	Subclinical hypothyroidism, poor compliance with thyroxine, drugs like amiodarone recovery phase of nonthyroidal illness, TSH Resistance
High	Low	Low	Hypothyroidism, Autoimmune thyroiditis, post radioiodine rx, post thyroidectomy, anti thyroid drugs, tyrosine kinase inhibitors & amiodarone amyloid deposits in thyroid, thyroid tumors & congenital hypothyroidism.
Low	High	High	Hypothyroidism, graves disease, toxic multinodular goiter, toxic adenoma, excess iodine or thyroxine intake, pregnancy related (hyperemesis gravidarum hydatiform mole)
Low	Normal	Normal	Subclinical Hypothyroidism, recent rx for hyperthyroidism, drugs like steroids & dopamine, Non thyroidal illness.
Low	Low	Low	Central Hypothyroidism, Non Thyroidal illness, Recent rx for Hyperthyroidism.
High	High	High	Interfering anti TPO antibodies, drug interference: Amiodarone, Heparin, Beta blockers, steroids & anti epileptics.

Diurnal Variation: TSH follows a diurnal rhythm and is at maximum between 2 am and 4 am, and is at a minimum between 6 pm and 10 pm. The variation is on the order of 50 to 206%. Biological variation: 19.7%



A/c Status : P
 Collected : 08/03/2025 09:37:00AM
 Ref By : SELF
 Collected at : WALKIN - VASHI (MAIN CENTRE)
 Shop No. 22, Raikar Bhavan, Near Navaratna Hotel,
 Sector 17, Vashi, Navi Mumbai, Maharashtra -
 400703
 Processed at : SHREEYASH HOSPITAL LAB, PANVEL
 Report Status : Final
 Reported : 8/3/2025 7:40:37PM
 Gender : Male
 Age : 36 Years

MediWHEEL Full Body Health Checkup Male >40/2D ECH

THYROID FUNCTION TESTS

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
-----------	---------	-----------------------	--------

(with in subject variation)

Reflex Tests: Anti thyroid Antibodies, USG Thyroid, TSH receptor Antibody, Thyroglobulin, Calcitonin

Limitations:

1. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. >5 mg/day) until atleast 8 hours following the last biotin administration.
2. Patient samples may contain heterophilic antibodies that could react in immunoassays to give falsely elevated or depressed results. this assay is designed to minimize interference from heterophilic antibodies.

Reference:

1. O.koulouri et al. / Best Practice and Research clinical Endocrinology and Metabolism 27(2013)
2. Interpretation of the thyroid function tests, Dayan et al. THE LANCET . Vol 357
3. Tietz , Text Book of Clinical Chemistry and Molecular Biology -5th Edition
4. Biological Variation: From principles to Practice-Callum G Fraser (AACCC Press)



Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SHREEYASH HOSPITAL LAB, PANVEL

MediWHEEL Full Body Health Checkup Male >40/2D ECH
LIVER FUNCTION TESTS

PARAMETER	RESULTS	BIOLOGICAL REF RANGES	METHOD
BILIRUBIN (TOTAL), Serum	1.45	0.10 - 1.20 mg/dL	Colorimetric
BILIRUBIN (DIRECT), Serum	0.51	0.00 - 0.30 mg/dL	Diazo
BILIRUBIN (INDIRECT), Serum	0.94	0.10 - 1.00 mg/dL	Calculated
TOTAL PROTEINS, Serum	6.78	6.40 - 8.30 g/dL	Buret
Albumin Serum	4.52	3.50 - 5.20 g/dL	BCG
GLOBULIN Serum	2.27	2.30 - 3.50 g/dL	Calculated
A/G RATIO Serum	1.99	1.00 - 2.00	Calculated
SGOT (AST), Serum	23.76	5.00 - 40.00 U/L	NADH (w/o P-5-P)
SGPT (ALT), Serum	36.47	5.00 - 45.00 U/L	NADH (w/o P-5-P)
GAMMA GT, Serum	19.56	3.00 - 60.00 U/L	Enzymatic
ALKALINE PHOSPHATASE, Serum	121.71	40.00 - 130.00 U/L	Colorimetric



R E P O R T

Name: CHATTERJEE SUBHO
Lab No.: 393765797
Ref By: SELF
Collected: 8/3/2025 9:37:00AM
A/c Status: P
Collected at: WALKIN - VASHI (MAIN CENTRE)
 Shop No. 22, Raikar Bhavan, Near Navaratna Hotel, Sector 17, Vashi, Navi Mumbai, Maharashtra - 400703

Age: 36 Years
Gender: Male
Reported: 8/3/2025 7:40:37PM
Report Status: Final
Processed at: SHREEYASH HOSPITAL LAB, PANVEL

URINE EXAMINATION REPORT

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	METHOD
PHYSICAL EXAMINATION			
Color	Pale Yellow	Pale Yellow	-
Transparency	CLEAR	Clear	-
CHEMICAL EXAMINATION			
Specific Gravity	Absent	1.002-1.035	Chemical Indicator
Reaction (pH)	6.5	5-8	pH Indicator
Proteins	Absent	Absent	Protein error principle
Glucose	Absent	Absent	GOD-POD
Ketones	Absent	Absent	Legals Test
Blood	Absent	Absent	Peroxidase
Bilirubin	Absent	Absent	Diazonium Salt
Urobilinogen	Normal	Normal	Diazonium Salt
Nitrite	Absent	Negative	Griess Test
MICROSCOPIC EXAMINATION			
(WBC) Pus cells / hpf	1-2	0-5/hpf	
Red Blood Cells / hpf	Absent	0-2/hpf	
Epithelial Cells / hpf	2-3	0-5/hpf	
Hyaline Casts	Absent	Absent	
Pathological cast	Absent	Absent	
Calcium oxalate monohydrate crystals	Absent	Absent	
Calcium oxalate dihydrate crystals	Absent	Absent	
Triple Phosphate crystals	Absent	Absent	
Uric acid crystals	Absent	Absent	
Amorphous debris	Absent	Absent	
Bacteria / hpf	4-5	0-20/hpf	
Yeast	Absent	Absent	
OTHERS	--		





Chief of Lab
 MD Pathology
 Dr Imran Mujawar

Imran Mujawar

URINE EXAMINATION REPORT

PARAMETER RESULTS BIOLOGICAL REF RANGE METHOD

SUBURBAN
 DIAGNOSTICS
 PRECISE TESTING - HAVING
 Name: CHATTERJEE SUBHO
 Lab No.: 393765797

Ref By : SELF
 Collected : 8/3/2025 9:37:00AM
 A/c Status : P
 Collected at : WALKIN - VASHI (MAIN CENTRE)
 Shop No. 22, Raikar Bhavan, Near Navaratna
 Hotel, Sector 17, Vashi, Navi Mumbai,
 Maharashtra - 400703

Age : 36 Years
 Gender : Male
 Reported : 8/3/2025 7:40:37PM
 Report Status : Final
 Processed at : SHREEYASH HOSPITAL LAB, PANVEL

T
R
O
P
E
R

MediWheel Full Body Health Checkup Male >40/2D ECH

PARAMETER RESULTS BIOLOGICAL REF RANGE METHOD

PHYSICAL EXAMINATION

EXAMINATION OF FAECES
Sample Not Received

CHEMICAL EXAMINATION

MICROSCOPIC EXAMINATION



-----End of report-----

IMPORTANT INSTRUCTIONS
The published test results relate to the submitted specimen. All test results are dependent on the quality of the sample received by the laboratory. Laboratory tests should be clinically correlated by a physician and are merely a tool to help arrive at a diagnosis. Unforeseen circumstances may cause a delay in the delivery of the report. Inconvenience is regretted. Certain tests may require further testing at an additional cost for derivation of exact value. Kindly submit the request within 72 hours post-reporting. The Court/Forum at Mumbai shall have exclusive jurisdiction in all disputes/claims concerning the test(s) & or results of the test(s). Test results are not valid for medico-legal purposes. This computer-generated medical diagnostic report has been verified by a doctor or an authorized medical professional. A physical signature is not required for this report. (#) sample drawn from an external source. If test results are alarming or unexpected, the client is advised to contact customer care immediately for possible remedial action. Tel: 022-61700000, Email: customer.service@suburbanandiagnostics.com <mailto:customer.service@suburbanandiagnostics.com>
West Reference Lab, Mumbai, is a CAP (0036028) Accredited laboratory.

