



INV. No. Patient Name Age/Gen QLSR-INV-K-11184/(2024-2025)(11138) **Mr. PRAVEEN BARDAN TOPNO**

Age/Gen 40 Years | Male Referred By **Dr. Self**

Source BERLIN DIAG INS CORP - (7)

Patient ID 11184

Sample Collected 09/11/2024 01:00 PM Sample Received 09/11/2024 04:58 PM Report Generated 09/11/2024 06:10 PM



Report Of Biochemistry Examination

Investigation	Result	Unit(s)	Reference Range	
GLUCOSE FASTING (FBS Plasma Glucose(F) Method (GOD-POD Method)	109.7	mg/dL	65 - 110	

Comments:

Fasting Blood Sugar/Glucose test a blood sample will be taken after an overnight fast. A fasting blood sugar level of less than 100mg/dL is normal. A fasting blood sugar level from 100 to 125 mg/dL is considered prediabetes. If it's 126 mg/dL or higher on two separate tests, you have diabetes.

GLUCOSE, POST PRANDIAL 2 HOURS

Plasma Glucose(PP) 128.2 mg/dL 75 - 140

Method (GOD-POD Method)

Note:

- 1. The diagnosis of Diabetes requires a fasting plasma glucose of > or = 126 mg/dL and/or a random / 2 hr post glucose value of > or = 200 mg/dL on at least 2 occasions
- 2. Very low glucose levels cause severe CNS dysfunction
- 3. Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis & is considered critical

GLYCOSYLATED HAEMOGLOBIN

Whole blood HbA1c	_	5.4	%	Non diabetic	level(< 6.0)
Method (HPLC)				Goal(< 7.0)	
Whole blood eAG (Estimated		108	mg/dl	-	
AverageGlucose Level)					
Method (CALCULATION)					

Note:

The Parameter indicates control over the last 90 Days

In the Blood, glucose adheres to haemoglobin (Hb) and make Glycosylated haemoglobin/HbA₁C, which provides a clue about the average blood glucose level over the last 8-12 weeks and it is an indicator for chronic glycaemic control along with effects of drug, diet and exercise.

In normal individuals, 90% is the adult haemoglobin fraction and the rest 8% is formed by HbA. Reduction of HbA_1C value reduces diabetic and cardiological related morbidity and mortality.

The short life span of RBC in haemoglobinopathy and chemically modified derivatives of haemoglobin (carbamylated Hb in renal failure and acetylated Hb, who are taking aspirin) can affect the results. Iron deficiency anaemia, liver disease, opiate addiction may interfere the test value.

HPLC, ion exchange chromatography is the ideal method for HbA₁C estimation. The target goal is <7%.

Report ID:- 44592 | Page 1/4





INV. No. QLSR-INV-K-11184/(2024-2025)(11138)

Patient Name Mr. PRAVEEN BARDAN TOPNO

Age/Gen

40 Years | Male **Dr. Self**

Referred By Source

BERLIN DIAG INS CORP - (7)

Patient ID 11184

Sample Collected 09/11/2024 01:00 PM Sample Received 09/11/2024 04:58 PM

Report Generated 09/11/2024 06:10 PM

Report Of Biochemistry Examination

Investigation Result Unit(s) Reference Range

Besides HbA₁C serum fructosamine can be measured.

American diabetes association guideline

Reference range

Non diabetic adult > 18 years : < 5.7%
Pediabetes : 5.7% - 6.4%
Diagnosing diabetes : > 6.5%

Lipid Profile

Lipid Profile				
Serum Triglyceride Method (Enzymatic,end point)		93.1	mg/dL	< 150
Serum Cholesterol Method (Oxidase, Esterase, Peroxid	ase)	196.6	mg/dL	125 - 200
Serum HDL-Chol Method (PTA/MgC12, Reflectance ph	notometry)	49.1	mg/dL	30 - 65
Serum LDL-Chol Method (Direct Homogeneous, Spec	ctrophotometry)	128.5	mg/dL	85 - 150
Serum VLDL-Chol		19	mg/dL	5 - 40
Serum LDL/HDL Choleste Method (Calculated)	erol Ratio	2.62		1.5 - 3.5
Serum Cholesterol/ HDL I	Ratio	4 00		Low Risk(0

Serum Cholesterol/ HDL Ratio 4.00 Low Risk(0 - 3) High Risk(5 - 10)

Method (Calculated)

Interpretation:

NATIONAL LIPID	TOTAL CHOLESTEROL in	TRIGLYCERIDE	LDL	NON HDL
ASSOCIATION	mg/dL	in mg/dL	CHOLESTEROL	CHOLESTEROL
RECOMMENDATIONS			in mg/dL	in mg/dL
(NLA-2014)			.91	
Optimal	<200	<150	<100	<130
Above Optimal	-	-	100- 129	130 - 159
Borderline High	200-239	150-199	130-159	160 - 189
High	>=240	200-499	160-189	190 - 219
Very High	-	>=500	>=190	>=220

Report ID:- 44592 | Page 2/4





E Certificate No.

INV. No.
Patient Name
Age/Gen
Referred By
Source

QLSR-INV-K-11184/(2024-2025)(11138) **Mr. PRAVEEN BARDAN TOPNO**

40 Years | Male

Dr. Self

BERLIN DIAG INS CORP - (7)

Patient ID 11184

Sample Collected 09/11/2024 01:00 PM Sample Received 09/11/2024 04:58 PM

Report Generated 09/11/2024 06:10 PM

Report Of Biochemistry Examination

Investigation Result Unit(s) Reference Range

Note:

- 1. Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.
- 2. Lipid Association of India (LAI) recommends screening of all adults above the age of 20 years for Atherosclerotic Cardiovascular Disease (ASCVD) risk factors especially lipid profile. This should be done earlier if there is family history of premature heart disease, dyslipidemia, obesity or other risk factors.
- 3. Indians tend to have hi<mark>gher triglyceride levels & Lower HDL choleste</mark>rol combined with small dense LDL particles, a pattern known as atherogenic dyslipidemia.
- 4. Non HDL Cholesterol comprises the cholesterol carried by all atherogenic particles, including LDL, IDL, VLDL & VLDL remnants, Chylomicron remnants & Lp(a).
- 5. LAI recommends LDL cholesterol as primary target and Non HDL cholesterol as co-primary treatment target.
- 6. Apolipoprotein B is an optional, secondary lipid target for treatment once LDL & Non HDL goals have been achieved.
- 7. Additional testing for Apolipoprotein B, hsCRP, Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement

Liver Function Test (I	FT)			
Serum Bilirubin (Total) Method (By Diphylline, Diazonium Sa		0.93	mg/dL	0.2 - 1.3
Serum Bilirubin (Direct) Method (Diphylline, Diazonium Salt)		0.22	mg/dL	0.1 - 0.4
Serum Bilirubin (Indirect) Method (Calculated)		0.71	mg/dL	0.2 - 1.1
Serum SGOT Method (IFCC)		32.8	U/L	17 - 59
Serum SGPT Method (IFCC)		31.9	U/L	21 - 72
Alkaline phosphatase (AL Method (IFCC)	P)	110.5	U/L	Adult (38 - 126)
Serum Total Protein Method (Biuret Method)		9.2	g/dL	Adult(6.2 - 8.2) Children(5.6 - 8.4)
Serum Albumin Method (BCG)		5.0	gm/dL	Newborn Children(2.4 - 4.8) Adult(3.5 - 5.0)
Serum Globulin Method (Calculated)		4.20	g/dL	Adult(2.3 - 3.6)

Report ID:- 44592 | Page 3/4





INV. No. Patient Name Age/Gen Referred By Source

QLSR-INV-K-11184/(2024-2025)(11138) Mr. PRAVEEN BARDAN TOPNO 40 Years | Male

Dr. Self

BERLIN DIAG INS CORP - (7)

11184 Patient ID

Sample Collected 09/11/2024 01:00 PM Sample Received 09/11/2024 04:58 PM Report Generated

09/11/2024 06:10 PM

Report Of Biochemistry Examination

Investigation	Result	Unit(s)	Reference Range
Serum A/G Ratio	1.19		1.0 - 2.3

Note

- 1. In an asymptomatic patient, Non alcoholic fatty liver disease (NAFLD) is the most common cause of increased AST, ALT levels. NAFLD is considered as hepatic manifestation of metabolic syndrome.
- 2. In most type of liver disease, ALT activity is higher than that of AST; exception may be seen in Alcoholic Hepatitis, Hepatic Cirrhosis, and Liver neoplasia. In a patient with Chronic liver disease, AST:ALT ratio>1 is highly suggestive of advanced liver fibrosis.
- 3. In known cases of Chr<mark>onic Liver disease due to Viral Hepatitis B & C</mark>, Alcoholic liver disease or NAFLD, Enhanced liver fibrosis (ELF) test may be used to evaluate liver fibrosis.
- In a patient with Chronic Liver disease, AFP and Des-gamma carboxyprothrombin (DCP)/PIVKA II can be used to assess risk for development of Hepatocellular Carcinoma.

Kidney Function Test	t (KFT)		
Serum Urea Method (GLDH,Kinetic Assay)	31.9	mg/dL	Adult (17 - 43) New Born (8.4 - 25.8)
Serum Creatinine Method (Modified Jaffe, Kinetic)	1.1	mg/dL	Infant (10.8 - 38.4) Male:(0.72-1.16) Female: (0.72-1.18) Neonate: (0.26 - 1.01) Infant (2months - less than 3yrs): (0.15-0.37) Children (3 yrs - less than 15 yrs): (0.24-0.73)
Serum Uric Acid Method (Uricase PAP)	5.2	mg/dL	3.5 - 7.2
Serum Sodium Method (By Indirect ISE)	137.8	mmol/L	136 - 145
Serum Potassium Method (By Indirect ISE)	4.1	mmol/L	3.5 - 5.1
Serum Chloride Method (By Ion-selective Electrode)	105.0	mmol/L	98 - 107

~~~~~ End of report ~~~~~

Report ID:- 44592 | Page 4/4







INV. No. Patient Name Age/Gen QLSR-INV-K-11184/(2024-2025)(11138) **Mr. PRAVEEN BARDAN TOPNO** 

40 Years | Male

Referred By Dr. Self

Source BERLIN DIAG INS CORP - (7)

Patient ID 11184

Sample Collected 09/11/2024 12:59 PM Sample Received 09/11/2024 01:11 PM Report Generated 09/11/2024 06:31 PM



#### **Report Of Haematology Examination**

| Investigation                | Result       | Unit(s) | Reference Range |
|------------------------------|--------------|---------|-----------------|
| ERYTHROCYTE SEDIMEN          | ITATION RATE |         |                 |
| ESR                          | 12           | mm      | < 20            |
| Method (Westergren & Manual) |              |         |                 |

#### Note

- 1. C-Reactive Protein (CRP) is the recommended test in acute inflammatory conditions.
- 2. Test conducted on EDTA whole blood at 37°C.
- 3. ESR readings are auto- corrected with respect to Hematocrit (PCV) values

| COMPLETE BLOOD C           | OUNT       |            |              |                                 |
|----------------------------|------------|------------|--------------|---------------------------------|
| Haemoglobin (Hb)%          | /          | 13.3       | gm%          | Adult Men (13 - 18)             |
| Method (By Sahlis Method ) |            |            |              | Adult Women (11.5 - 16.5)       |
|                            |            |            |              | Children (11 - 13)              |
|                            |            |            |              | Children (1-6) : (12 - 14)      |
|                            |            |            |              | Children (6-12) : (12 - 14)     |
| PCV                        |            | 42.6       | %            | 35 - 45                         |
| Total Platelets Count (P   |            | 1.3        | Lacs Per cmm | 1.5 - 4                         |
| Total RBC (Red Cell Cou    | nt)        | 5.2        | mill./uL     | Women (4.2 - 5.4)               |
|                            |            |            |              | Male (4.7 - 6.1)                |
|                            |            |            |              | Children (4.6 - 4.8)            |
| Total Leucocyte Count (    | TLC)       | 5,000      | Per cmm      | Adult :- (4,000 - 11,000)       |
| Method (Flow Cytometry)    |            |            |              | New Born (10,000 - 26,000)      |
|                            |            |            |              | (1-4) Years : (6,000 - 18,000)  |
|                            |            |            |              | (5-7) Years : (5,000 - 15,000)  |
| MCV                        |            | 70         | £I           | (8-12) Years : (4,500 - 12,500) |
| MCV                        |            | 79<br>24.7 | fL           | 76 - 96                         |
| MCH                        |            | 24.7       | pg<br>~ /d!  | 22 - 32                         |
| MCHC                       | Lauranutaa | 31.2       | g/dL         | 30 - 35                         |
| Differential count of      | Leucocytes | FF         | 0/           | 40. 70                          |
| Neutrophils                |            | 55<br>37   | %<br>%       | 40 - 70                         |
| Lymphocytes                |            | 37         | %<br>%       | 15 - 40                         |
| Monocytes                  |            | 02         |              | 00 - 6                          |
| Eosinophils                |            | 06         | %<br>%       | 0.5 - 7                         |
| Basophils                  |            | 00         | 70           | 00 - 01                         |

#### **Comment:**

CBC is a powerful diagnostic tool in various hematological and non-hematological conditions. It can be

Report ID:- 44623 | Page 1/2





INV. No. Patient Name Age/Gen

Referred By

QLSR-INV-K-11184/(2024-2025)(11138) Mr. PRAVEEN BARDAN TOPNO

40 Years | Male

Dr. Self

Source BERLIN DIAG INS CORP - (7) Patient ID 11184

Sample Collected 09/11/2024 12:59 PM Sample Received 09/11/2024 01:11 PM Report Generated

09/11/2024 06:31 PM

#### **Report Of Haematology Examination**

**Investigation** Result Unit(s) Reference Range

used to diagnose various conditions like anemia, hemoglobinopathies, infections. leukemia, nutritional deficiencies, parasitemias, etc. For microcytic indices, a Mentzer index of less than 13 suggests that the patient may have thalassemia trait, and an index of more than 13 suggests that the patient may have iron deficiency.

#### Blood Grouping (A B O) and Rh Type

Whole blood Blood Group Whole blood Rh Type

O" Positive

#### Note:

- 1. Both forward and reverse grouping performed.
- 2. Test conducted on EDTA whole blood.

~~~~ End of report ~~~~

Report ID:- 44623 | Page 2/2







INV. No.
Patient Name
Age/Gen
Referred By
Source

QLSR-INV-K-11184/(2024-2025)(11138) **Mr. PRAVEEN BARDAN TOPNO**

40 Years | Male

Dr. Self

BERLIN DIAG INS CORP - (7)

Patient ID 11184

Sample Collected 09/11/2024 04:53 PM Sample Received 09/11/2024 04:54 PM

Report Generated 09/11/2024 04:34 PM



Report Of Clini Patho Examination

Investigation Result Unit(s) Reference Range **FASTING URINE SUGAR** Nil Urine Glucose (Sugar) gm% ~~~~~ End of report ~~~~

Report ID:- 44634 | Page 1/1





E Certificate No.

INV. No. Patient Name Age/Gen Referred By

Source

QLSR-INV-K-11184/(2024-2025)(11138) **Mr. PRAVEEN BARDAN TOPNO**

40 Years | Male

Dr. Self

BERLIN DIAG INS CORP - (7)

Patient ID 11184

Sample Collected 09/11/2024 04:53 PM Sample Received 09/11/2024 04:54 PM

Report Generated 09/11/2024 06:44 PM



Report Of Clini Patho Examination

| Investigation | | Result | Unit(s) | Reference Range |
|---------------------------------------|---------|-----------------------|-----------|-----------------|
| | | | | |
| | crosco | pic Examination (R/M) | | |
| Physical Examination | | Chana | | Dala Vallani |
| Colour | | Straw | | Pale Yellow |
| Urine Appearance | | Transparent | | |
| Urine Deposit | | Absent | | |
| Urine Specific Gravity | | 1.010 | | 1.010 - 1.030 |
| Urine Reaction | | Acidic | | |
| Chemical Examination | | | | |
| Urine Glucose (Sugar) | | Nil | gm% | |
| Urine Protein (Albumin) | | Absent | | |
| Urine pH | | 6.0 | | 6.0 |
| Urine Ketone Body | | Absent | | |
| Urine Blood | | Negative | | |
| Urine Phosphate (Amor <mark>ph</mark> | ious de | eposits) Absent | | |
| <u> Urine Microscopic Exar</u> | ninati | <u>on</u> | | |
| Urine Red blood cells | | Absent | /HPF | 0-2 |
| Urine Pus Cells | | 1-2 | /HPF | 0-5 |
| Urine Epithelial cells | | 0-2 | /HPF | 0-4 |
| Urine Bacteria | | Absent | | |
| Urine Cast | | Absent | /HPF | |
| Urine Crystals | | Absent | /HPF | |
| Urine Yeast cells | | Absent | , | |
| Urine Spermatozoa | | Absent | /HPF | |
| 5c 5pcaco20a | | 7.650110 | , | |
| | | ~~~~~ End of repo | ort ~~~~~ | |
| | | Ena or repo | ,, , | |
| | | | | |

Report ID:- 44635 | Page 1/1





INV. No. Patient Name Age/Gen Referred By Source

QLSR-INV-K-11184/(2024-2025)(11138) Mr. PRAVEEN BARDAN TOPNO

40 Years | Male Dr. Self

BERLIN DIAG INS CORP - (7)

Patient ID 11184 Sample Collected

09/11/2024 01:00 PM Sample Received 09/11/2024 01:12 PM Report Generated 11/11/2024 10:31 AM

Report Of Immunology Examination

| Investigation | Result | Unit(s) | Reference Range |
|---|--------|---------|---|
| (Thursday Durafile 1) | | | |
| (Thyroid Profile-I) Serum T3 Method (ECLIA) | 0.96 | ng/mL | (0.8 - 2.0)
11-15 Years (0.83 - 2.13)
1-10 Years (0.94 - 2.69)
1-12 Months (1.05 - 2.45)
1-7 Days (0.36 - 3.16)
1-4 Weeks (1.05 - 3.45) |
| Serum T4 Method (ECLIA) | 14.64 | μg/dL | 1-4 Weeks (1.03 - 3.43)
(5.1 - 14.1)
1-12 Months (5.9 - 16)
1-7 Days (11 - 22) |
| Serum TSH
Method (ECLIA) | 0.87 | μlU/mL | 1-4 Weeks (8.2 - 17)
1-10 Years (6.4 - 15)
11-15 Years (5.5 - 12)
Up to 1 Week (0.7-11.0)
1 week-4 week (0.7- 11.0) |
| | | | 1-12 Months (0.7- 8.4)
1-19 Years (0.6-4.9)
19 Years Above (0.5-5.5)
1st Trimester (0.6 - 3.4)
2nd Trimester (0.37 - 3.6) 3rd
Trimester(0.38 - 4.04) |

Mild to moderate degree of elevation normal T3&T4 levels indicates impaired thyroid hormone reserves and indicates subclinical hypothyroidism.

Mild to moderate decrease with normal T3 & T4 indicates subclinical hyperthyroidism.

TSH measurement is used for screening & diagnosis of Euthyroidism, hypothyroidism & hyperthyroidism. Suppressed TSH (< 0.01 μ IU/ml) suggests diagnosis of hyperthyroidism.

Elevated concentration of TSH (>7 μ IU/ml) suggest diagnosis of hypothyroidism.

Please correlate clinically.

~~~~~ End of report ~~~~~

Report ID:- 44797 | Page 1/1





INV. No. Patient Name Age/Gen Referred By QLSR-INV-K-11184/(2024-2025)(11138) **Mr. PRAVEEN BARDAN TOPNO** 

40 Years | Male

Dr. Self

Source BERLIN DIAG INS CORP - (7)

Patient ID 11184

Sample Collected 09/11/2024 01:00 PM Sample Received 09/11/2024 01:12 PM

Report Generated 11/11/2024 11:43 AM

#### **Report Of Immunology Examination**

Investigation Result Unit(s) Reference Range

#### Prostate Specific Antigen (PSA) Total

Serum PROSTATE SPECIFIC ANTIGEN (PSA) 0.64

Method (ECLIA)

ng/ml

< 4.0 For Healthy Man

#### P.S.A.

PSA is elevated in benign prostrate hypertrophy. Clinically an elevated PSA value is not of diagnostic value as a specific test for cancer and should only be used in conjunction with other clinical symptom and diagnostic procedure.

~~~~~ End of report ~~~~~

Report ID:- 44801 | Page 1/1







| Patient Name | Mr. PRAVEEN BARDAN | Requested By | MEDIWHEEL |
|--------------|--------------------|----------------|------------------------------------|
| | BER/2024/OPD27885 | Procedure Date | 09.11.2024 |
| Age/Sex | 40Y/MALE | Hospital | BERLIN DIAGNOSTICS AND DAY
CARE |

USG WHOLE ABDOMEN

Liver: The liver is normal in size (15.1 cm) and outline. It shows normal parenchymal echotexture. No obvious focal or diffuse lesion is seen. The intra and extra hepatic biliary passage are not dilated. The portal vein is normal in caliber at the portal hepatis.

Gall bladder: The gall bladder is normal in size, has normal wall thickness with no evidence of calculi.

CBD: The CBD is of normal caliber.

Pancreas: The pancreas is normal in size and echogenicity with distinct outline. No obvious focal lesion is seen.

Kidneys: Both kidneys were normal in position:

Right kidney measures 10.2 cm Left kidney measures 10.2 cm

The renal cortical thickness and corticomedullary differentiation were adequate on both sides. No evidence of renal calculus or hydronephrosis seen on either side.

Spleen: The spleen is normal in size and echogenicity.

Urinary Bladder: The urinary bladder is normal in size. Its walls show a smooth outline. There is no evidence of any intraluminal or perivesical abnormality.

Prostate: The prostate is normal in size, measures (20.7 gm) and shows normal parenchymal echogenicity.

No significant probe tenderness in RIF.

No evidence of pleural effusion on either side.

No evidence of ascites or lymphadenopathy seen.

IMPRESSION: NORMAL STUDY.

Please correlate clinically.

Dr. Ambuj Srivastav M.D. Consultant Radiologist.





TOPNO, PRAVEEN BARDAN Patient ID 202427885 09.11.2024 2:18:49pm


EXERCISE TEST / ECG Strips

96 bpm 110/80 mmHg


PRETEST SUPINE 0:08

BRUCE 0.0 km/h 0.0 %

BERLIN DIAGNOSTICS AND DAY CARE





Attending MD:




99 bpm 110/80 mmHg


PRETEST HYPERV. 0:42







GE CardioSoft V6.73 (2) 25 man/s 10 mm/mV 60IIz 0.01 - 20Hz FRF- HR(V6,V4) TOPNO, PRAVEEN BARDAN Patient ID 202427885 09.11.2024 2:24:07pm 153 bpm 124/94 mmHg EXERCISE STAGE 2 04:20 12-Lead Report (PEAK EXERCISE) V3 V2 Start of Test: 2:18;40pm BRUCE 4.0 km/h 12.0 % Measured at 60ms Post J (10mm/mV)
Auto Points Lead 0.05 0.05 BERLIN DIAGNOSTICS AND DAY CARE ٧6 V5 35 -0.60 0.00 Page 4



09.11.2024 TOPNO, PRAVEEN BARDAN Patient ID 202427885 2:25:52pm aVF BASEIJINE 0.65 -0.65 -0.15 -0.50 -1.73aVR 0.50 0.20 0.92 0.75 0.25 aVF -1.10 aVR CURRENT 0.05 0.20 0.20 -0.30 0.10 1.180 aVF aVR~ 104 bpm 114/84 mmHg #1 01:50 Comparative Medians Report RECOVERY V2 0.60 45 V6 0.75 1.05 √3 0.04 0.70 1.98 1.45 BRUCE 0.0 km/b 0.0% 0.83 V6 -0.30 0.99 15 -0.10 4 1.37 -835 √3 0.35 12 0.01 ≤ 8 454 BERLIN DIAGNOSTICS AND DAY CARE Lead ST Level (mm) ST Slope (mV/s)

GR CardioSoft V6.73 (2) 25 mm/s 10 mm/mV 60Hz 0.01 - 20Hz FRF- HR(V6,V4)

Start of Test: 2:18:40pm

09,112024 Patient ID 202427885 TOPNO, PRAVEEN BARDAN 2:26:42pm GE CardioSoft V6.73 (2) 25 nom/s 10 nom/mV 60Hz 0.01 - 20Hz FRF - HR(V6,V4) 60 ms post J BASELINE 0.75 aVR aVL -1.73 0.50 Ξ 0.65 aVF 0.65 60 ms post J CURRENT 0.97 aVL aVR 0.50 aVF -0.96 0.20 0.05 0.15 105 bpm 110/80 mmHg aVL THE 02:40 Comparative Medians Report RECOVERY ≤ 0.70 ₹ 1.64 1.45 V5 1.65 0.75 1.05 0.60 V6 BRUCE 0.0 km/h 0.0% Start of Test: 2:18:40pm 1.01 0.10 V3-4 0.30 V6 0.95 V5 0.75 3 3 46 BERLIN DIAGNOSTICS AND DAY CARE J.ead ST Level (mm) ST Slope (mV/s)

Page

| T0mm/mV 60Hz 0.01-20Hz PRF- HEART V5.4 | aVF ₩\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\ | aVE → ↑ ↑ .0.15
-0.65 | aVR | 0.50
0.34 | 0.75 | 0.20 mm
0.33 mV/s | EXERCISE
0:00
103 bpm
110/80 mmHg |
|--|---|--------------------------|---------------------------|-----------------------|--------------------|----------------------|---|
| 2)
1-20Hz FRP- HEAD | -1.05
-0.96 | aVL→
0.45
0.94 | 9.50
0.50 | -1.00 | -0.66 | -0.05
0.65 | MAX. ST
EXERCISE
329
153 bpm
124/94 mmHg |
| | aVF
-0.85
0.96 | aVL
0.25
-0.14 | aVR VIII
0.55
-0.41 | -0.70
0.06 | 0.99 | -0.20 | PEAK EXERCISE
EXERCISE
4:20
153 bpm
124/94 mmHg |
| Unconfirmed | aVF
-0.20
0.75 | avI_
0.00
-0.16 | aVR | 0.15 | 1.00
1.00 | 0.34 | ISE TEST END RECOVERY 2:39 105 bpm 110/80 mmHg |
| | 1.12
0.60 | V5
0.75
1.42 | V4 \\ 1.05 | V3 - 1.65 | 1.45
1.64 | 0.70
0.04 | BASELINE
EXERCISE
0:00
103 bpm
110/80 mmHg |
| | V65
-0.65 | -0.80
-0.80 | V4 V4 V | 0.10
2.65 | V2
0.00
1.90 | 0.65
0.65 | MAX. ST
EXERCISE
3:29
153 bpm
124/94 mmHg |
| | -0.80
1.15 | 0.90
1.41 | V4 V4 V
0.70
1.87 | V3 V3
0.05
2.73 | -0.10
1.79 | 1.05
1.05 | PEAK EXERCISE
EXERCISE
4:20
153 bpm
124/94 mmHg |
| | 0.78
0.78 | V5
-0.25
0.92 | 0.10 | V3
0.30
1.52 | 0.25
0.97 | 0.45
0.04 | E TEST END RECOVERY 2:39 105 bpm 110/80 mmHg |

09.11.2024 Patient ID 202427885

2:18:40pm Male 175 cm 78.4 kg Meds: 40yrs Asian

Medical History: Test Reason:

Comment: Technician: Test Type: Ref. MD: Ordering MD:

BRUCE: Total Exercise Time 04:20

Max HR: 155 hpm 86% of max predicted 180 hpm HR at rest: 97

Max BP: 124/94 mmHg BP at rest: 110/80 Max RPP: 18972 mmHg*bpm Maximum Workload: 7,00 METS

Max. ST: -1.05 mm, 0.00 mV/s in II; EXERCISE STAGE 2 03:29

Arrhythmia: A:68

ST/HR index: 1.57 µV/bpm

Reasons for Termination: Target heart rate achieved

Response to Exercise: appropriate response. Chest Pain: none. Arrhythmias: none. Summary: Functional Capacity: normal. HR Response to Exercise: appropriate. BP

Conclusion:

TMT TEST IS NEGATIVE FOR INDUCIBLE INCHEMIA.

SO Committant Cardiologist

MALENDREG No. GO125

Dr. Amar Kumar

HEER (Bedand)

| 7 | | | | | | | | | | |
|-------------|--|---|--|--|---|---|---|--|---|---|
| Stage Name | Time
in Stage | Speed
(km/h) | Grade
(%) | Workload
(METS) | HR
(bpun) | BP
(mmHg) | RPP
(mmHg*bpm | (huin) | STLevel | Comment |
| | | 1000 | | The state of the s | 7.00 | /Greenway | Comment of the | Comme | (rum m) | |
| CHIDIN'S | 0000 | | | | | | | | | |
| SCIPINE | 00:09 | 0.00 | 0.00 | 0.1 | S | 110/00 | 02201 | = | | |
| STAUDING | 00.72 | 100 | 0 0 0 0 | | OK | 110/00 | 00001 | 0 | 0.60 | |
| Transfer to | 00.60 | 0.00 | 0.00 | 1.0 | 96 | 10000 | 03.60 | ٥ | 250 | |
| DIFLEY. | 14:00 | 0.00 | 0.00 | 1.0 | 99 | 31028 | ウルカラ | 5 | 5 60 | |
| WARM-UP | 00:18 | 1 60 | 00.0 | _ | 103 | 2000 | 1000 | 0 | 0.00 | |
| STACE | 017.00 | 3 70 | 0000 | () | 193 | 110190 | 0.06514 | 0 | 0,75 | |
| CTACCO | 01.00 | 2.70 | 0.00 | 4.6 | 136 | 118/88 | 16048 | \$ | -010 | |
| 31000.2 | 07.10 | 4.00 | 12,00 | 7.0 | Lsa | 124/94 | 18972 | > | 000 | |
| | 02:4[| 0.00 | 0.00 | 0 | 201 | 08/011 | 11550 | ه د | 0.00 | |
| | | | | | | | 2000 | 0 | -0.20 | |
| | Stage Name SUPINE STANDING HYPERV. WARM-UP STAGE I STAGE 2 | Stage Name Tunc in Stage SUPINE 00:09 STANDING 00:23 HYPERV. 00:11 WARM-UP 00:18 STAGE 1 03:00 STAGE 2 01:20 02:41 | Time
in Stage (
00:09
00:11
00:11
00:18
03:00
01:20
01:20
02:41 | Time Speed in Stage (km/h) 00:09 0.00 00:23 0.00 00:11 0.00 00:18 1.60 03:00 2.70 01:20 4.00 02:41 0.00 | Time Speed Grade in Stage (km/h) (%) 00:09 | Time Speed Grade in Stage (km/h) (%) 00:09 | Time Speed Grade Workload in Stage (km/h) (%) (METS) 00:09 | Time Speed Grade Workload HR Bp (%) (METS) (bpm) (mmHg) (00:09 0.00 0.00 1.0 96 110/80 00:11 0.00 0.00 1.0 96 110/80 00:11 0.00 0.00 1.0 99 110/80 00:18 1.60 0.00 1.1 103 110/80 03:00 2.70 [0.00 4.6 136 118/88 01:20 4.00 12.00 7.0 153 124/94 02:41 0.00 0.00 1.0 1.0 105 110/80 | Time in Speed in Stage Speed (km/h) Grade (%) Workload HR (mmHg) (mmHg) (mmHg) (mmHg) (mmHg) (mmHg) (mmHg) BP (mmHg) (mmH | Time in Stage Speed (km/h) Grade (%) Workload HR (mmHg) (mmHg) (mmHg) (mmHg) (mmHg) (mmHg) Bp (mmHg) |



| Patient Name | Mr. PRAVEEN BARDAN | B | |
|--------------|----------------------------|----------------|-------------------------------|
| MRN | TOPNO
PED/2024/CPD2799F | Requested By | MEDIWHEEL |
| PO TO | BER/2024/OPD27885 | Procedure Date | 09.11.2024 |
| Age/Sex | 40Y/MALE | Centre | BERLIN DIAGNOSTICS & DAY CARE |

X-RAY CHEST PA VIEW

OBSERVATIONS:

Both lung fields are clear.

Both hila are normal.

Both CP angles are normal,

Cardiac contour and size are within normal limits.


The bony rib cage is normal.

Soft tissue are normal.

IMPRESSION: NORMAL STUDY.

Dr. Ambuj Srivastav M.D. Consultant Radiologist.

We regret typographical errors if any. Please contact us for correction.



EN BARDAN TOPNO AGE 40Y/M MEDIWHEEL BER/202427885 CHEST PA VIEW 09/11/2024 BERLIN DIAGNOSTICS & DAY CARE, BARIATU ROAD, RANCHI.