



Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169032
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 17:00 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL PATHOLOGY

Complete Urine Examination (CUE)

Investigation	Observed Value	Biological Reference Intervals
Physical Examination		
Colour Method:Physical	Pale Yellow	Straw to Yellow
Appearance Method:Physical	Clear	Clear
Chemical Examination		
Reaction and pH Method:pH- Methyl red & Bromothymol blue	7.5	4.6-8.0
Specific gravity Method:Bromothymol Blue	1.005	1.003-1.035
Protein Method:Tetrabromophenol blue	Negative	Negative
Glucose Method:Glucose oxidase/Peroxidase	Positive(+++)	Negative
Blood Method:Peroxidase	Negative	Negative
Ketones Method:Sodium Nitroprusside Method	Negative	Negative
Bilirubin Method:Dichloroanilinediazonium	Negative	Negative
Leucocytes Method:3 hydroxy5 phenylpyrrole + diazonium	Negative	Negative
Nitrites Method:Diazonium + 1,2,3,4 tetrahydrobenzo (h) quinolin 3-ol	Negative	Negative
Urobilinogen Method:Dimethyl aminobenzaldehyde	0.2	0.2-1.0 mg/dl
Microscopic Examination		
Pus cells (leukocytes) Method:Microscopy	0-1	2 - 3 /hpf
Epithelial cells Method:Microscopy	0-1	2 - 5 /hpf
RBC (erythrocytes) Method:Microscopy	Absent	Absent
Casts Method:Microscopy	Absent	Occasional hyaline casts may be seen



Name : **MR.RAJ SAURABH** TID/SID : UMR2599458/ 29169032
Age / Gender : 55 Years / Male Registered on : 08-Mar-2025 / 08:56 AM
Ref.By : MEDI WHEELS Collected on : 08-Mar-2025 / 09:12 AM
Req.No : BIL5397038 Reported on : 08-Mar-2025 / 17:00 PM
Reference : Arcofemi Health Care Ltd -

TEST REPORT

Crystals	Absent	Phosphate, oxalate, or urate crystals may be seen
Method:Microscopy		
Others	Nil	Nil
Method:Microscopy		

Method: Semi Quantitative test ,For CUE

Reference: Godkar Clinical Diagnosis and Management by Laboratory Methods, First South Asia edition. Product kit literature.

Interpretation:

The complete urinalysis provides a number of measurements which look for abnormalities in the urine. Abnormal results from this test can be indicative of a number of conditions including kidney disease, urinary tract infection or elevated levels of substances which the body is trying to remove through the urine . A urinalysis test can help identify potential health problems even when a person is asymptomatic. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakur

Dr Debleena Thakur
Consultant Pathologist
KMC NO : 89765





Name : **MR.RAJ SAURABH** TID/SID : UMR2599458/ 29169031
Age / Gender : 55 Years / Male Registered on : 08-Mar-2025 / 08:56 AM
Ref.By : MEDI WHEELS Collected on : 08-Mar-2025 / 09:12 AM
Req.No : BIL5397038 Reported on : 08-Mar-2025 / 15:57 PM
Reference : Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF HEMATOPATHOLOGY

Blood Grouping ABO And Rh Typing

Parameter	Results
Blood Grouping (ABO)	B
Rh Typing (D)	POSITIVE

Method: Hemagglutination Tube Method by Forward & Reverse Grouping

Reference: Kit literature

Interpretation: The ABO grouping and Rh typing test determines blood type grouping (A,B, AB, O) and the Rh factor (positive or negative). A person's blood type is based on the presence or absence of certain antigens on the surface of their red blood cells and certain antibodies in the plasma. A,B,H antigens are not fully developed at birth, increase gradually in strength and become fully expressed around 1 year of age. It is mandatory to repeat blood grouping at/after one year of age for new born babies &/or done on cord blood

Note: All individuals carry other blood group system antigens in addition to ABO and Rh. Antibody screening is recommended to all individuals before blood transfusion to detect any unexpected antibodies.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakur

Dr Debleena Thakur
Consultant Pathologist
KMC NO : 89765





Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169031
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 15:14 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF HEMATOPATHOLOGY

Erythrocyte Sedimentation Rate (ESR)

Investigation	Observed Value	Biological Reference Intervals
ESR 1st Hour Method:Modified Westergren	2	<=20 mm/hour

Complete Blood Count (CBC)

Investigation	Observed Value	Biological Reference Interval
Hemoglobin Method:Spectrophotometry	14.0	13.0-18.0 g/dL
Packed Cell Volume Method:Derived from Impedance	41.0	40-54 %
Red Blood Cell Count. Method:Impedance Variation	4.57	4.3-6.0 Mill/Cumm
Mean Corpuscular Volume Method:Derived from Impedance	89.8	78-100 fL
Mean Corpuscular Hemoglobin Method:Derived from Impedance	30.6	27-32 pg
Mean Corpuscular Hemoglobin Concentration Method:Derived from Impedance	34.1	31.5-36 g/dL
Red Cell Distribution Width - CV Method:Derived from Impedance	13.7	11.5-16.0 %
Red Cell Distribution Width - SD Method:Derived from Impedance	38.8	39-46 fL
Total WBC Count. Method:Impedance Variation	5390	4000-11000 cells/cumm
Neutrophils Method:Impedance Variation, Flowcytometry	51.3	40-75 %
Lymphocytes Method:Microscopy	37.9	20-45 %
Eosinophils Method:Impedance Variation,Method_Desc= Flow Cytometry	2.9	01-06 %
Monocytes Method:Impedance Variation, Flowcytometry	7.3	01-10 %
Basophils. Method:Impedance Variation,Method_Desc= Flow Cytometry	0.6	00-02 %



Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169031
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 15:14 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

Absolute Neutrophils Count. Method:Calculated	2765	1500-6600 cells/cumm
Absolute Lymphocyte Count Method:Calculated	2043	1500-3500 cells/cumm
Absolute Eosinophils count. Method:Calculated	156	40-440 cells/cumm
Absolute Monocytes Count. Method:Calculated	393	<1000 cells/cumm
Absolute Basophils count. Method:Calculated	32	<200 cells/cumm
Platelet Count. Method:Impedance Variation	1.40	1.4-4.4 lakhs/cumm
Mean Platelet Volume. Method:Derived from Impedance	13.1	7.9-13.7 fL
Plateletcrit. Method:Derived from Impedance	0.17	0.18-0.28 %

Method: Automated Hematology Analyzer, Microscopy

Reference: Dacie and Lewis Practical Hematology, 12th Edition

Interpretation: A Complete Blood Picture (CBP) is a screening test which can aid in the diagnosis of a variety of conditions and diseases such as anemia, leukemia, bleeding disorders and infections. This test is also useful in monitoring a person's reaction to treatment when a condition which affects blood cells has been diagnosed. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakur

Dr Debleena Thakur
Consultant Pathologist
KMC NO : 89765





Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169034F
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 13:55 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Blood Urea Nitrogen (BUN)

Investigation	Observed Value	Biological Reference Interval
Blood Urea Nitrogen.	11.1	6-20 mg/dL
Method:Kinetic, Urease - GLDH, Calculated		

Interpretation: Urea is a waste product formed in the liver when protein is metabolized. Urea is released by the liver into the blood and is carried to the kidneys, where it is filtered out of the blood and released into the urine. Since this is a continuous process, there is usually a small but stable amount of urea nitrogen in the blood. However, when the kidneys cannot filter wastes out of the blood due to disease or damage, then the level of urea in the blood will rise. The blood urea nitrogen (BUN) evaluates kidney function in a wide range of circumstances, to diagnose kidney disease, and to monitor people with acute or chronic kidney dysfunction or failure. It also may be used to evaluate a person's general health status as well.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics

Creatinine, Serum

Investigation	Observed Value	Biological Reference Interval
Creatinine.	0.64	0.7-1.3 mg/dL
Method:Spectrophotometry, Jaffe - IDMS Traceable		

Interpretation:

Creatinine is a nitrogenous waste product produced by muscles from creatine. Creatinine is majorly filtered from the blood by the kidneys and released into the urine, so serum creatinine levels are usually a good indicator of kidney function. Serum creatinine is more specific and more sensitive indicator of renal function as compared to BUN because it is produced from muscle at a constant rate and its level in blood is not affected by protein catabolism or other exogenous products. It is also not reabsorbed and very little is secreted by tubules making it a reliable marker. Serum creatinine levels are increased in pre renal, renal and post renal azotemia, active acromegaly and gigantism. Decreased serum creatinine levels are seen in pregnancy and increasing age.

Biological reference interval changed; Reference: Tietz Textbook of Clinical Chemistry & Molecular Diagnostics, Fifth Edition.

Glucose Fasting (FBS)

Investigation	Observed Value	Biological Reference Interval
Glucose Fasting	130	Normal: <100 mg/dL Impaired FG: 100-125 mg/dL Diabetes mellitus: >=126 mg/dL
Method:Hexokinase		

Interpretation: It measures the Glucose levels in the blood with a prior fasting of 9-12 hours. The test helps screen a symptomatic/ asymptomatic person who is at risk for Diabetes. It is also used for regular monitoring of glucose levels in people with Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2022



	Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169033
	Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
	Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
	Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 14:23 PM
			Reference	: Arcofemi Health Care Ltd -

TEST REPORT

Glucose Post Prandial (PPBS)

Investigation	Observed Value	Biological Reference Interval
Glucose Post Prandial Method:Hexokinase	187	Normal : <140 mg/dL Impaired PG: 140-199 mg/dL Diabetes mellitus: >=200 mg/dL

Interpretation: This test measures the blood sugar levels 2 hours after a normal meal. Abnormally high blood sugars 2 hours after a meal reflect that the body is not producing sufficient insulin which is indicative of Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2020.

Glycosylated Hemoglobin (HbA1C)

Investigation	Observed Value	Biological Reference Interval
Glycosylated Hemoglobin (HbA1c) Method:High-Performance Liquid Chromatography	8.0	Non-diabetic: <= 5.6 % Pre-diabetic: 5.7 - 6.4 % Diabetic: >= 6.5 %
Estimated Average Glucose (eAG) Method:Calculated	183	mg/dL %

Interpretation: It is an index of long-term blood glucose concentrations and a measure of the risk for developing microvascular complications in patients with diabetes. Absolute risks of retinopathy and nephropathy are directly proportional to the mean HbA1c concentration. In persons without diabetes, HbA1c is directly related to risk of cardiovascular disease.

In known diabetic patients, HbA1c can be considered as a tool for monitoring the glycemic control.
Excellent Control - 6 to 7 %,
Fair to Good Control - 7 to 8 %,
Unsatisfactory Control - 8 to 10 %
and Poor Control - More than 10 %.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2018.

Bun/Creatinine Ratio

Investigation	Observed Value	Biological Reference Interval
BUN/Creatinine Ratio Method:Calculated	17.3	12-16

Reference:

A Manual of Laboratory Diagnostic Tests. Edition 7, Lippincott Williams and Wilkins, By Frances Talaska Fischbach, RN, BSN, MSN, and Marshall Barnett Dunning 111, BS, MS, Ph.D.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kanya-SN



PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE



Name : **MR.RAJ SAURABH**
Age / Gender : 55 Years / Male
Ref.By : MEDI WHEELS
Req.No : BIL5397038

TID/SID : UMR2599458/
Registered on : 08-Mar-2025 / 08:56 AM
Collected on :
Reported on :
Reference : Arcofemi Health Care Ltd -

TEST REPORT

Dr.Kavya S N
Consultant Pathologist
KMC NO : 84851





Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169033
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 14:23 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Lipid Profile

Investigation	Observed Value	Biological Reference Interval
Total Cholesterol Method:Spectrophotometry , CHOD - POD	168	Desirable: < 200 mg/dL Borderline: 200-239 mg/dL High: >= 240 mg/dL
HDL Cholesterol Method:Spectrophotometry , Direct Measurement	42	Optimal : >=60 mg/dL Borderline : 40-59 mg/dL High Risk <40 mg/dL
Non HDL Cholesterol Method:Calculated	126	Optimal : <130 mg/dL Above Optimal : 130-159 mg/dL Borderline : 160-189 mg/dL High Risk : 190-219 mg/dL Very high Risk : >=220 mg/dL
LDL Cholesterol Method:Calculated	93.4	Optimum: <100 mg/dL Near/above optimum: 100-129 mg/dL Borderline: 130-159 mg/dL High: 160-189 mg/dL Very high: >=190 mg/dL
VLDL Cholesterol Method:Calculated	32.60	<30 mg/dL
Total Cholesterol/HDL Ratio Method:Calculated	4	Optimal : <3.3 Low Risk : 3.4-4.4 Average Risk : 4.5-7.1 Moderate Risk : 7.2-11.0 High Risk : >11.0
LDL/HDL Ratio Method:Calculated	2.22	Optimal : 0.5-3.0 Borderline : 3.1-6.0 High Risk : >6.0
Triglycerides Method:Spectrophotometry, Enzymatic - GPO/POD	163	Normal:<150 mg/dL Borderline: 150-199 mg/dL High: 200-499 mg/dL Very high: >=500 mg/dL mg/dl #

Interpretation: Lipids are fats and fat-like substances which are important constituents of cells and are rich sources of energy. A lipid profile typically includes total cholesterol, high density lipoproteins (HDL), low density lipoprotein (LDL), chylomicrons, triglycerides, very low density lipoproteins (VLDL), Cholesterol/HDL ratio. The lipid profile is used to assess the risk of developing a heart disease and to monitor its treatment. The results of the lipid profile are evaluated along with other known risk factors associated with heart disease to plan and monitor treatment. Treatment options require clinical correlation. **Reference:** Third Report of the National Cholesterol Education program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III), JAMA 2001.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---



PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE



Name : **MR.RAJ SAURABH**
Age / Gender : 55 Years / Male
Ref.By : MEDI WHEELS
Req.No : BIL5397038

TID/SID : UMR2599458/
Registered on : 08-Mar-2025 / 08:56 AM
Collected on :
Reported on :
Reference : Arcofemi Health Care Ltd -

TEST REPORT

Kavya SN

Dr.Kavya S N
Consultant Pathologist
KMC NO : 84851





Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169033
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 14:23 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Liver Function Test (LFT)

Investigation	Result	Biological Reference Interval
Total Bilirubin. Method:Spectrophotometry, Diazo method	0.74	Neonates: <=15.0 mg/dL Adults: <=1.2 mg/dL
Direct Bilirubin. Method:Spectrophotometry, Diazo method	0.37	<=0.30 mg/dL
Indirect Bilirubin. Method:Calculated	0.37	Neonates: <= 14.7 mg/dL Adults: <= 1.0 mg/dL
Alanine Aminotransferase ,(ALT/SGPT) Method: IFCC without pyridoxal phosphate activation	27	<=41 U/L
Aspartate Aminotransferase,(AST/SGOT) Method: IFCC without pyridoxal phosphate activation	16	<=40 U/L
ALP (Alkaline Phosphatase). Method:Spectrophotometry , IFCC	66	40-129 U/L
Gamma GT. Method:Spectrophotometry , IFCC	24	<60 U/L
Total Protein. Method:Spectrophotometry, Biuret	7.1	6.4-8.3 g/dL
Albumin. Method:Spectrophotometry, Bromcresol Green	4.5	3.5-5.2 g/dL
Globulin. Method:Spectrophotometry, Bromcresol Green	2.60	2.0-3.5 g/dL
A/GRatio. Method:Calculated	1.73	1.1-2.5

Interpretation: Liver functions tests help to identify liver disease, its severity, and its type. Generally these tests are performed in combination, are abnormal in liver disease, and the pattern of abnormality is indicative of the nature of liver disease. An isolated abnormality of a single liver function test usually means a non-hepatic cause. If several liver function tests are simultaneously abnormal, then hepatic etiology is likely.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist
KMC NO : 84851



Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169033
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 15:31 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Prostate Specific Antigen (PSA) Total

Investigation	Observed Value	Biological Reference Interval
Prostate Specific Antigen (PSA) Total	0.690	0.0-4.0 ng/mL
Method:ECLIA		

Interpretation: PSA is a protein produced by cells in the prostate and is used to screen men for prostate cancer. PSA levels are elevated in Prostate cancer, and other conditions such as benign prostatic hyperplasia (BPH) and inflammation of the prostate. An elevated PSA may be followed by a biopsy and other tests like urinalysis and ultrasound to rule out urinary tract infections and for an accurate diagnosis. PSA levels are vital to determine the effectiveness of treatment and to detect recurrence in diagnosed cases of prostate cancer.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist
KMC NO : 84851





Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169033
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 15:31 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Thyroid Profile (T3,T4,TSH)

Investigation	Observed Value	Biological Reference Interval
Triiodothyronine Total (T3) Method:ECLIA	1.02	0.80-2.00 ng/mL Note: Biological Reference Ranges are changed due to change in method of testing.
Thyroxine Total (T4) Method:ECLIA	7.37	4.6-12.0 µg/dL
Thyroid Stimulating Hormone (TSH) Method:ECLIA	3.69	0.27-4.20 µIU/mL

Interpretation: A thyroid profile is used to evaluate thyroid function and/or help diagnose hypothyroidism and hyperthyroidism due to various thyroid disorders. T4 and T3 are hormones produced by the thyroid gland. They help control the rate at which the body uses energy, and are regulated by a feedback system. TSH from the pituitary gland stimulates the production and release of T4 (primarily) and T3 by the thyroid. Most of the T4 and T3 circulate in the blood bound to protein. A small percentage is free (not bound) and is the biologically active form of the hormones.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics, Carl A. Burtis, David E. Bruns.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist
KMC NO : 84851





Name	: MR.RAJ SAURABH	TID/SID	: UMR2599458/ 29169033
Age / Gender	: 55 Years / Male	Registered on	: 08-Mar-2025 / 08:56 AM
Ref.By	: MEDI WHEELS	Collected on	: 08-Mar-2025 / 09:12 AM
Req.No	: BIL5397038	Reported on	: 08-Mar-2025 / 14:23 PM
		Reference	: Arcofemi Health Care Ltd -

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Uric Acid, Serum

Investigation	Observed Value	Biological Reference Interval
Uric Acid. Method:Enzymatic	5.4	3.4-7.0 mg/dL

Interpretation: It is the major product of purine catabolism. Hyperuricemia can result due to increased formation or decreased excretion of uric acid which can be due to several causes like metabolic disorders, psoriasis, tissue hypoxia, pre-eclampsia, alcohol, lead poisoning, acute or chronic kidney disease, etc. Hypouricemia may be seen in severe hepato cellular disease and defective renal tubular reabsorption of uric acid.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Kavya SN

Dr.Kavya S N
Consultant Pathologist
KMC NO : 84851





PLEASE SCAN QR CODE

Name : Mr . RAJ SAURABH
Age/Gender : 55 Years/Male
Ref By : MEDI WHEELS
Reg.No : BIL5397038

TID : UMR2599458
Registered On : 08-Mar-2025 08:56 AM
Reported On : 08-Mar-2025 11:46 AM
Reference : Arcofemi Health Care Ltd
- Medi Whe

ECHOCARDIOGRAM REPORT

MESUREMENTS

IVS (D): 0.9CM LVID (D): 4.3CM LVPW (D):1.1 CM

IVS(S): 1.8CM LVID (S): 3.2CM LVPW(S):1.9 CM

AO: 2.2 CM LA:2.6 CM

EF: 62%

VALVES:

MITRAL VALVE : NORMAL

AORTIC VALVE : NORMAL

TRICUSPID VALVE : NORMAL

PULMONARY VALVE : NORMAL

CHAMBERS:

LEFT ARTIUM : NORMAL

RIGHT ARTIUM : NORMAL

LEFT VENTRICLE : NORMAL

RIGHT VENTRICLE : NORMAL

SEPTAE:

IVS : INTACT

IAS : INTACT

GREAT ARTERIES:

AORTA : NORMAL

PULMONARY ARTERY : NORMAL



PLEASE SCAN QR CODE

Name : Mr . RAJ SAURABH
Age/Gender : 55 Years/Male
Ref By : MEDI WHEELS
Reg.No : BIL5397038

TID : UMR2599458
Registered On : 08-Mar-2025 08:56 AM
Reported On : 08-Mar-2025 11:46 AM
Reference : Arcofemi Health Care Ltd
- Medi Whe

DOPPLER STUDY:

MITRAL VALVE : E -0.7M/S A -0.5M/S TRIVIAL MR
AORTIC VALVE : 1.2M/S
TRICUSPID VALVE : NORMAL
PULMONARY VALVE : 0.8M/S

WALL MOTION ABNORMALITIES: NO RWMA

PERICARDIUM : NORMAL
VEGETATION / THROMBUS : NO

FINAL DIAGNOSIS:

- **NORMAL CARDIAC CHAMBERS.**
- **NORMAL LV SYSTOLIC FUNCTION.**
- **NO REGIONAL WALL MOTION ABNORMALITIES**
- **NO PE / CLOT / VEGETATION SEEN.**

*** End Of Report ***

Dr Prabhu K
Consultant Cardiologist



PLEASE SCAN QR CODE

Name : Mr . RAJ SAURABH
Age/Gender : 55 Years/Male
Ref By : MEDI WHEELS
Reg.No : BIL5397038

TID : UMR2599458
Registered On : 08-Mar-2025 08:56 AM
Reported On : 08-Mar-2025 12:29 PM
Reference : Arcofemi Health Care Ltd
- Medi Whe

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is distended. No evidence of calculus or wall thickening. No pericholecystic fluid collection. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation preserved. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation preserved. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cm)	Parenchymal thickness (cm)
Right Kidney	10.6	2.0
Left Kidney	10.0	1.9

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern. It measures 4.4 x 2.8 x 3.1 cm volume: 19.6cc.

No evidence of ascites.

IMPRESSION:

- **No significant abnormality detected.**


*** End Of Report ***

Dr Lohith H P
Consultant Radiologist

55 Years
Male

QRS 94 ms
QT / QTcBaz 370 / 405 ms
PR 144 ms
P 92 ms
RR / PP 838 / 833 ms
P / QRS / T 133 / 21 / 48 degrees

Normal sinus rhythm
Normal ECG



GE MAC2000 1 1 12SL™ V241 25 mm/s 10 mm/mV ADS 0.56-20 Hz 2x5x6_25_R1

Unconfirmed