

Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. : -

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

R

E

Use a QR Code Scanner Application To Scan the Code

Collected : 09-Nov-2024 / 09:21 Reported : 09-Nov-2024 / 15:58

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

CBC (Com	plete	Blood	Count)	, Blood

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
RBC PARAMETERS			
Haemoglobin	12.1	12.0-15.0 g/dL	Spectrophotometric
RBC	4.67	3.8-4.8 mil/cmm	Elect. Impedance
PCV	36.8	36-46 %	Calculated
MCV	78.8	81-101 fl	Measured
MCH	25.9	27-32 pg	Calculated
MCHC	32.8	31.5-34.5 g/dL	Calculated
RDW	15.8	11.6-14.0 %	Calculated
WBC PARAMETERS			
WBC Total Count	5340	4000-10000 /cmm	Elect. Impedance
WBC DIFFERENTIAL AND ABSO	LUTE COUNTS		
Lymphocytes	48.6	20-40 %	
Absolute Lymphocytes	2595.2	1000-3000 /cmm	Calculated
Monocytes	6.5	2-10 %	
Absolute Monocytes	347.1	200-1000 /cmm	Calculated
Neutrophils	43.7	40-80 %	
Absolute Neutrophils	2333.6	2000-7000 /cmm	Calculated
Eosinophils	1.1	1-6 %	
Absolute Eosinophils	58.7	20-500 /cmm	Calculated
Basophils	0.1	0.1-2 %	
Absolute Basophils	5.3	20-100 /cmm	Calculated
Immature Leukocytes	-		

WBC Differential Count by Absorbance & Impedance method/Microscopy.

PLATELET PARAMETERS

Platelet Count	299000	150000-410000 /cmm	Elect. Impedance
MPV	8.0	6-11 fl	Measured
PDW	13.8	11-18 %	Calculated

RBC MORPHOLOGY

Hypochromia Mild

Microcytosis Occasional



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. : - Collected : 09-Nov-2024 / 09:21

Reg. Location : Borivali West (Main Centre) Reported : 09-Nov-2024 / 19:00

Macrocytosis -

Anisocytosis -

Poikilocytosis -

Polychromasia -

Target Cells -

Basophilic Stippling -

Normoblasts -

Others -

WBC MORPHOLOGY -

PLATELET MORPHOLOGY -

COMMENT -

Specimen: EDTA Whole Blood

ESR, EDTA WB-ESR 15 2-20 mm at 1 hr. Sedimentation

Clinical Significance: The erythrocyte sedimentation rate (ESR), also called a sedimentation rate is the rate red blood cells sediment in a period of time.

Interpretation:

Factors that increase ESR: Old age, Pregnancy, Anemia

Factors that decrease ESR: Extreme leukocytosis, Polycythemia, Red cell abnormalities- Sickle cell disease

Limitations:

- It is a non-specific measure of inflammation.
- · The use of the ESR as a screening test in asymptomatic persons is limited by its low sensitivity and specificity.

Reflex Test: C-Reactive Protein (CRP) is the recommended test in acute inflammatory conditions.

Reference:

- Pack Insert
- Brigden ML. Clinical utility of the erythrocyte sedimentation rate. American family physician. 1999 Oct 1;60(5):1443-50.

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
*** End Of Report ***





Dr.ANUPA DIXIT M.D.(PATH) Consultant Pathologist

Authenticity Check

Use a OR Code Scanner

Application To Scan the Code

Page 2 of 12



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. :

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

R

E

Use a QR Code Scanner Application To Scan the Code

Collected : 09-Nov-2024 / 09:21 Reported : 09-Nov-2024 / 13:42

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
GLUCOSE (SUGAR) FASTING, Fluoride Plasma Fasting	84.0	Non-Diabetic: < 100 mg/dl Impaired Fasting Glucose: 100-125 mg/dl Diabetic: >/= 126 mg/dl	Hexokinase
GLUCOSE (SUGAR) PP, Fluoride Plasma PP	74.2	Non-Diabetic: < 140 mg/dl Impaired Glucose Tolerance: 140-199 mg/dl Diabetic: >/= 200 mg/dl	Hexokinase
BILIRUBIN (TOTAL), Serum	0.47	0.1-1.2 mg/dl	Colorimetric
BILIRUBIN (DIRECT), Serum	0.22	0-0.3 mg/dl	Diazo
BILIRUBIN (INDIRECT), Serum	0.25	0.1-1.0 mg/dl	Calculated
TOTAL PROTEINS, Serum	7.9	6.4-8.3 g/dL	Biuret
ALBUMIN, Serum	4.3	3.5-5.2 g/dL	BCG
GLOBULIN, Serum	3.6	2.3-3.5 g/dL	Calculated
A/G RATIO, Serum	1.2	1 - 2	Calculated
SGOT (AST), Serum	21.7	5-32 U/L	NADH (w/o P-5-P)
SGPT (ALT), Serum	14.3	5-33 U/L	NADH (w/o P-5-P)
GAMMA GT, Serum	10.3	3-40 U/L	Enzymatic
ALKALINE PHOSPHATASE, Serum	82.4	35-105 U/L	Colorimetric
BLOOD UREA, Serum	21.7	12.8-42.8 mg/dl	Kinetic
BUN, Serum	10.1	6-20 mg/dl	Calculated
CREATININE, Serum	0.66	0.51-0.95 mg/dl	Enzymatic



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. :

eGFR, Serum

Reg. Location

: Borivali West (Main Centre)

125

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Collected: 09-No

:09-Nov-2024 / 09:21

Calculated

Enzymatic

Reported :09-Nov-2024 / 13:42

(ml/min/1.73sqm)

Normal or High: Above 90 Mild decrease: 60-89

Mild to moderate decrease: 45-

59

Moderate to severe decrease:30

-44

Severe decrease: 15-29 Kidney failure:<15

Note: eGFR estimation is calculated using 2021 CKD-EPI GFR equation

URIC ACID, Serum 2.2 2.4-5.7 mg/dl

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West
*** End Of Report ***









Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. : -

Reg. Location: Borivali West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

:09-Nov-2024 / 09:21

Reported :09-Nov-2024 / 17:34

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE GLYCOSYLATED HEMOGLOBIN (HbA1c)

PARAMETER RESULTS BIOLOGICAL REF RANGE METHOD

Glycosylated Hemoglobin 5.4 Non-Diabetic Level: < 5.7 % HPLC (HbA1c), EDTA WB - CC Prediabetic Level: 5.7-6.4 %

Prediabetic Level: 5.7-6.4 % Diabetic Level: >/= 6.5 %

Collected

Estimated Average Glucose 108.3 mg/dl Calculated

(eAG), EDTA WB - CC

Intended use:

- In patients who are meeting treatment goals, HbA1c test should be performed at least 2 times a year
- · In patients whose therapy has changed or who are not meeting glycemic goals, it should be performed quarterly
- For microvascular disease prevention, the HbA1C goal for non pregnant adults in general is Less than 7%.

Clinical Significance:

- HbA1c, Glycosylated hemoglobin or glycated hemoglobin, is hemoglobin with glucose molecule attached to it.
- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of glycosylated hemoglobin in the blood.

Test Interpretation:

- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of Glycosylated hemoglobin in the blood.
- HbA1c test may be used to screen for and diagnose diabetes or risk of developing diabetes.
- To monitor compliance and long term blood glucose level control in patients with diabetes.
- Index of diabetic control, predicting development and progression of diabetic micro vascular complications.

Factors affecting HbA1c results:

Increased in: High fetal hemoglobin, Chronic renal failure, Iron deficiency anemia, Splenectomy, Increased serum triglycerides, Alcohol ingestion, Lead/opiate poisoning and Salicylate treatment.

Decreased in: Shortened RBC lifespan (Hemolytic anemia, blood loss), following transfusions, pregnancy, ingestion of large amount of Vitamin E or Vitamin C and Hemoglobinopathies

 $\textbf{Reflex tests:} \ Blood \ glucose \ levels, \ CGM \ (Continuous \ Glucose \ monitoring)$

References: ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition.

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
*** End Of Report ***





Dr.ANUPA DIXIT M.D.(PATH) Consultant Pathologist

Page 5 of 12



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. : -

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

R

E

Use a QR Code Scanner Application To Scan the Code

Collected : 09-Nov-2024 / 09:21

Reported :09-Nov-2024 / 19:40

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE URINE EXAMINATION REPORT

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
PHYSICAL EXAMINATION			
Color	Pale yellow	Pale Yellow	Light scattering
Transparency	Clear	Clear	Light scattering
CHEMICAL EXAMINATION			
Specific Gravity	1.002	1.002-1.035	Refractive index
Reaction (pH)	6.5	5-8	pH Indicator
Proteins	Absent	Absent	Protein error principle
Glucose	Absent	Absent	GOD-POD
Ketones	Absent	Absent	Legals Test
Blood	Absent	Absent	Peroxidase
Bilirubin	Absent	Absent	Diazonium Salt
Urobilinogen	Normal	Normal	Diazonium Salt
Nitrite	Negative	Negative	Griess Test
MICROSCOPIC EXAMINATION			
(WBC)Pus cells / hpf	0.0	0-5/hpf	
Red Blood Cells / hpf	0.0	0-2 /hpf	
Epithelial Cells / hpf	0.1	0-5/hpf	
Hyaline Casts	0.0	0-1/hpf	
Pathological cast	0.0	0-0.3/hpf	
Calcium oxalate monohydrate crystals	0.0	0-1.4/hpf	
Calcium oxalate dihydrate crystals	0.0	0-1.4/hpf	
Triple phosphate crystals	0.0	0-1.4/hpf	
Uric acid crystals	0.0	0-1.4/hpf	
Amorphous debris	Absent	Absent	
Bacteria / hpf	1.5	0-29.5/hpf	
Yeast	Absent	Absent	



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. : -

Reg. Location : Borivali West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected : 09-Nov-2024 / 09:21

Reported :09-Nov-2024 / 19:40

Note: Microscopic examination performed by Automated Cuvette based technology. All the Abnormal results are confirmed by reagent strips and Manual method. The Microscopic examination findings are mentioned in decimal numbers as the arithmetic mean of the multiple fields scanned using microscopy. Reference: Pack Insert.

Others -

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
*** End Of Report ***

Dr.ANUPA DIXIT M.D.(PATH) Consultant Pathologist



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. : -

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Collected

Reported

: 09-Nov-2024 / 09:21

:09-Nov-2024 / 17:23

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE BLOOD GROUPING & Rh TYPING

PARAMETER RESULTS

ABO GROUP 0

Rh TYPING Negative

NOTE: Test performed by automated Erythrocytes magnetized technology (EMT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

Clinical significance:

ABO system is most important of all blood group in transfusion medicine

Limitations:

- · ABO blood group of new born is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

Refernces:

- 1. Denise M Harmening, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia
- 2. AABB technical manual

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
*** End Of Report ***





Dr.VRUSHALI SHROFF M.D.(PATH) Pathologist

Page 8 of 12



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. :

Reg. Location : Borivali West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

:09-Nov-2024 / 09:21

Reported :09-Nov-2024 / 13:42

Collected

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE LIPID PROFILE

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
CHOLESTEROL, Serum	183.0	Desirable: <200 mg/dl Borderline High: 200-239mg/dl High: >/=240 mg/dl	CHOD-POD
TRIGLYCERIDES, Serum	67.3	Normal: <150 mg/dl Borderline-high: 150 - 199 mg/dl High: 200 - 499 mg/dl Very high:>/=500 mg/dl	GPO-POD
HDL CHOLESTEROL, Serum	54.3	Desirable: >60 mg/dl Borderline: 40 - 60 mg/dl Low (High risk): <40 mg/dl	Homogeneous enzymatic colorimetric assay
NON HDL CHOLESTEROL, Serum	128.7	Desirable: <130 mg/dl Borderline-high:130 - 159 mg/dl High:160 - 189 mg/dl Very high: >/=190 mg/dl	Calculated
LDL CHOLESTEROL, Serum	116.0	Optimal: <100 mg/dl Near Optimal: 100 - 129 mg/dl Borderline High: 130 - 159 mg/dl High: 160 - 189 mg/dl Very High: >/= 190 mg/dl	Calculated
VLDL CHOLESTEROL, Serum	12.7	< /= 30 mg/dl	Calculated
CHOL / HDL CHOL RATIO, Serum	3.4	0-4.5 Ratio	Calculated
LDL CHOL / HDL CHOL RATIO, Serum	2.1	0-3.5 Ratio	Calculated

^{*}Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West
*** End Of Report ***





Dr.JYOT THAKKER..
M.D. (PATH), DPB
Pathologist & AVP(Medical Services)

Page 9 of 12



Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. :

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

R

E

Use a QR Code Scanner Application To Scan the Code

Collected : 09-Nov-2024 / 09:21

Reported :09-Nov-2024 / 13:30

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE THYROID FUNCTION TESTS

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
Free T3, Serum	4.4	3.5-6.5 pmol/L	ECLIA
Free T4, Serum	13.4	11.5-22.7 pmol/L First Trimester:9.0-24.7 Second Trimester:6.4-20.59 Third Trimester:6.4-20.59	ECLIA
sensitiveTSH, Serum	3.45	0.35-5.5 microIU/ml First Trimester:0.1-2.5 Second Trimester:0.2-3.0 Third Trimester:0.3-3.0 microU/ml	ECLIA



CID : 2431420359

Name : MS.POOJA POOJA

Age / Gender : 25 Years / Female

Consulting Dr. Collected Reported Reg. Location : Borivali West (Main Centre) :09-Nov-2024 / 13:30



Authenticity Check

Use a OR Code Scanner Application To Scan the Code

:09-Nov-2024 / 09:21

Interpretation:

A thyroid panel is used to evaluate thyroid function and/or help diagnose various thyroid disorders.

Clinical Significance:

- 1)TSH Values between high abnormal upto15 microIU/ml should be correlated clinically or repeat the test with new sample as physiological
- can give falsely high TSH.
- 2)TSH values may be trasiently altered becuase of non thyroidal illness like severe infections, liver disease, renal and heart severe burns, trauma and surgery etc.

TSH	FT4 / T4	FT3 / T3	Interpretation
High	Normal	Normal	Subclinical hypothyroidism, poor compliance with thyroxine, drugs like amiodarone, Recovery phase of non-thyroidal illness, TSH Resistance.
High	Low	Low	Hypothyroidism, Autoimmune thyroiditis, post radio iodine Rx, post thyroidectomy, Anti thyroid drugs, tyrosine kinase inhibitors & amiodarone, amyloid deposits in thyroid, thyroid tumors & congenital hypothyroidism.
Low	High	High	Hyperthyroidism, Graves disease, toxic multinodular goiter, toxic adenoma, excess iodine or thyroxine intake, pregnancy related (hyperemesis gravidarum, hydatiform mole)
Low	Normal	Normal	Subclinical Hyperthyroidism, recent Rx for Hyperthyroidism, drugs like steroids & dopamine), Non thyroidal illness.
Low	Low	Low	Central Hypothyroidism, Non Thyroidal Illness, Recent Rx for Hyperthyroidism.
High	High	High	Interfering anti TPO antibodies, Drug interference: Amiodarone, Heparin, Beta Blockers, steroids & anti epileptics.

Diurnal Variation: TSH follows a diurnal rhythm and is at maximum between 2 am and 4 am, and is at a minimum between 6 pm and 10 pm. The variation is on the order of 50 to 206%. Biological variation:19.7%(with in subject variation)

Reflex Tests: Anti thyroid Antibodies, USG Thyroid , TSH receptor Antibody. Thyroglobulin, Calcitonin

Limitations:

- 1. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. >5 mg/day) until atleast 8 hours following the last biotin administration.
- 2. Patient samples may contain heterophilic antibodies that could react in immunoassays to give falsely elevated or depressed results. this assay is designed to minimize interference from heterophilic antibodies.

- 1.O.koulouri et al. / Best Practice and Research clinical Endocrinology and Metabolism 27(2013)
- 2.Interpretation of the thyroid function tests, Dayan et al. THE LANCET. Vol 357
- 3. Tietz , Text Book of Clinical Chemistry and Molecular Biology -5th Edition
- 4. Biological Variation: From principles to Practice-Callum G Fraser (AACC Press)

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West *** End Of Report ***





BMhaskar Dr.KETAKI MHASKAR M.D. (PATH) **Pathologist**



CID : 2431420359

Name : MS.POOJA POOJA

: 25 Years / Female Age / Gender

Consulting Dr.

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

:09-Nov-2024 / 14:11

Collected Reported :09-Nov-2024 / 20:07

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

PARAMETER RESULTS BIOLOGICAL REF RANGE METHOD

Urine Sugar (Fasting) Absent **Absent** Urine Ketones (Fasting) Absent Absent

Urine Sugar (PP) Absent Urine Ketones (PP) **Absent Absent**

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab *** End Of Report ***





Dr. VRUSHALI SHROFF M.D.(PATH) **Pathologist**

Page 12 of 12

R E P Age / Gender : 25 Years/Female : 09-Nov-2024 / 09:01 Collected R : 09-Nov-2024 / 17:13 Consulting Dr. : Reported T Reg.Location : Borivali West (Main Centre)

PHYSICAL EXAMINATION REPORT

History and Complaints:

No Complaint

EXAMINATION FINDINGS:

Height (cms):

Afebrile

Temp (0c): Blood Pressure (mm/hg):

Pulse:

76/min

60 Weight (kg):

Skin:

NAD NAD

Nails: Lymph Node:

Not palpable

Systems

Cardiovascular: S1S2-Normal Chest-Clear Respiratory:

Genitourinary: GI System:

NAD NAD

CNS:

NAD

IMPRESSION:

Nowheal

ADVICE:

physician Feb.

CHIEF COMPLAINTS:

Hypertension:

1HD

Nο 3) Arrhythmia No 4) Diabetes Mellitus No

5) Tuberculosis No Asthama

Pulmonary Disease

No

No

No

E P 0 vame of the time of MS.POOJA POOJA : 09-Nov-2024 / 09:01 R Age / Gender : 25 Years/Female Collected : 09-Nov-2024 / 17:13 T Reported Consulting Dr. : : Borivali West (Main Centre) Reg.Location

dore	No
8) Thyroid/ Endocrine disorders	No
9) Nervous disorders	No
co cystem	No
11) Genital urinary disorder 11) Genital urinary diseases or sympto	ms No
miatic lollit discus	No
ninod disease of	No
(4) Cancer/lump growths	No
15) Congenital disease	No
. C. rappries	No
17) Musculoskeletal System	

PERSONAL HISTORY:

No Alcohol No 1) 2) Smoking Mixed Diet No 3) Medication

DR. NITIN SONAVANE M.B.B.S.AFLH, D.DIAR, D.CARD. CONSULTANT CARDIOLOGIST REGD - 0: 87714

Suburban Diagnost'es (I) Put. Ltd. *** End Of Report ***

3018 302, 3rd Floor Med Eleganance Above Taring - L. T. Road, Borivali (V-22), 1 - 7 5 - 420 002

> Dr.NITIN SONAVANE PHYSICIAN



R T CID NO: 2431420359 AGE/SEX: 25Y/F PATIENT'S NAME: MS.POOJA POOJA DATE: 09/11/2024 REF BY: ----

E p

0

2-D ECHOCARDIOGRAPHY

- RA, LA RV is Normal Size.
- No LV Hypertrophy.
- 3. Normal LV systolic function. LVEF 60 % by bi-plane
- No RWMA at rest.
- Aortic, Pulmonary, Mitral valves normal. Trivial TR.
- Great arteries: Aorta: Normal
 - No mitral valve prolaps.
- Inter-ventricular septum is intact and normal.
- Intra Atrial Septum intact.
- Pulmonary vein, IVC, hepatic are normal.
- 10.No LV clot.
- 11. No Pericardial Effusion
- No Diastolic disfunction. No Doppler evidence of raised LVEDP.



AGE/SEX: 25Y/F
DATE: 09/11/2024

1. AO root diameter	2.9 cm
2. IVSd	0.9 cm
3. LVIDd	4.5 cm
4. LVIDs	2.1 cm
5. LVPWd	0.9 cm
6. LA dimension	3.6 cm
7. RA dimension	3.6 cm
8. RV dimension	3.0 cm
9. Pulmonary flow vel:	0.9 m/s
10. Pulmonary Gradient	3.4 m/s
11. Tricuspid flow vel	1.6 m/s
12. Tricuspid Gradient	11 m/s
13. PASP by TR Jet	21 mm Hg
14. TAPSE	2.2 cm
15. Aortic flow vel	1.2 m/s
16. Aortic Gradient	6 m/s
17. MV:E	0.8 m/s
18. A vel	$0.6 \mathrm{m/s}$
19. IVC	15 mm
20. E/E'	8

Impression:
Normal 2d echo study.

Disclaimer

Echo may have interfinite observer variations in measurements as the study is observer dependent and changes with PCs homodynamics. Please co-relate findings with patients clinical status.

End of Report

DR. S. NITIN Consultant Cardiologist Reg. No. 87714

E

0

R



CID

: 2431420359

Name

: Ms Pooja Pooja

Age / Sex

: 25 Years/Female

Ref. Dr

.

Reg. Location

: Borivali West

Reg. Date

: 09-Nov-2024

Reported

: 09-Nov-2024 / 15:46

R

Ε

0

R

X-RAY CHEST PA VIEW

Both lung fields are clear.

Both costo-phrenic angles are clear.

The cardiac size and shape are within normal limits.

The domes of diaphragm are normal in position and outlines.

The skeleton under review appears normal.

IMPRESSION: NO SIGNIFICANT ABNORMALITY IS DETECTED.

Kindly correlate clinically.

Note: Investigations have their limitations. Solitary tridiological investigations never confirm the final diagnosis. They only help in diagnosing the disease in correlation to clinical symptoms and other related tests. X-ray is known to have inter-occurve variations. Further relation up assaging may be needed in some cases for confirmation / exclusion of diagnosis. Please interpret accordingly. In case of any typingraphical error / spelling error in the report, patient is requested to immediately crethed the centre within T days post which the center will not be responsible for any recultivation.

-End of Report---

Dr. Pranali Mahale MD,Radiodiagnosis Consultant Radiologist Reg no. 2019/07/5682

Franch

Click here to view images <<tmageLink>>



CID

: 2431420359

Name

: Ms Pooja Pooja

Age / Sex

: 25 Years/Female

Ref. Dr

Reg. Location

: Borivali West

Authenticity Check

Der a Oft Cade Seatter Application To Scan the Code R

Е

p

0

R

т

: 09-Nov-2024

Reg. Date : 09-Nov-2024 / 11:46 Reported

USG WHOLE ABDOMEN

LIVER: Liver is normal in size, shape and echotexture. There is no intra-hepatic biliary radical dilatation. No evidence of any obvious focal lesion.

GALL BLADDER: Gall bladder is partially distended.

(Tiny polyps/calculi may be missed due to technical limitations, sub-optimal distension of GB, adjacent gases and inter-machine variability in resolution settings)

PORTAL VEIN: Portal vein is normal. CBD: CBD is normal.

PANCREAS: Pancreas obscured due to bowel gases.

KIDNEYS: Right kidney measures 8.9 x 4.5 cm. Left kidney measures 9.3 x 4.5 cm. Both kidneys are normal in shape and echotexture. Corticomedullary differentiation is maintained. There is no evidence of any hydronephrosis, hydroureter or calculus.

SPLEEN: Spleen is normal in size, shape and echotexture. No focal lesion is seen.

URINARY BLADDER: Urinary bladder is distended and normal. Wall thickness is within normal limits.

UTERUS: Uterus is anteverted, normal and measures 7.2 x 2.5 x 4.4 cm. Uterine myometrium shows homogenous echotexture. Endometrium is normal in thickness and measures 7.3 mm. Cervix appears normal.

OVARIES: Both ovaries appear normal in size and echotexture.

The right ovary measures 3.2 x 1.9 cm.

The left ovary measures 2.8×1.6 cm.

Bilateral adnexa is clear.

No free fluid or obvious significant lymphadenopathy is seen.

Click here to view images http://3.111.232.119/iRISViewer/NeoradViewer?

sionNo-2024110909034541



CID

: 2431420359

Name

: Ms Pooja Pooja

Age / Sex

Reg. Location

: 25 Years/Female

Ref. Dr

r

: Borivali West

Authenticity Check



R T

E

0

Cac a QR Code Scanner Application To Sean the Code

Reg. Date

: 09-Nov-2024

Reported

: 09-Nov-2024 / 11:46

Opinion:

· No significant abnormality is detected.

For clinical correlation and follow up.

Note: Investigations have their limitations. Solitary radiological investigations never confirm the final diagnosis. They only help in diagnosing the disease in correlation to clinical symptoms and other related tests. USG is known to have inter-observer variations. Further / Follow-up intaging may be needed in some cases for confirmation / exclusion of diagnosis. Patient was explained in detail verbally about the USG findings. USG measurements and its limitations in case of any typegraphical error in the report, patient is requested to immediately contact the center for restification within 7 days post which the center will not be responsible for any rectification. Please interpret accordingly.

-----End of Report-----

Dr. Pranali Mahale MD,Radiodiagnosis Consultant Radiologist Reg no. 2019/07/5682

REGD. OFFICE: Dr. Lail Path Labs Ltd., Block E. Sector 18, Robini, New Dolbi - 110060. CIN No.: L74899DL 1506/00165389 ...

REPORT

Date:-

Name: Poota

CID: 2431420356

Sex / Age: /

EYE CHECK UP

Chief complaints:

Systemic Diseases:

Past history:

Unaided Vision:

Aided Vision:

Refraction:

Mil

Re Le

6/6 6/6

M/6 H/6

(Right Eye)

(Left Eye)

	Sph	Cyl	Axis	Vn	Sph	Cyl	Axis	Vn
Distance				\\				
Near						, in 2008,		

Colour Vision: Normal / Abnormal

Remark:

Suburing Diagnost to (I) Put. Ltd.
Above Tar Survival Supanence
Corival (Violation Control of Road)

PRECISE FEATING HEALTHER LIVING SUBURBAN

Patient Name: POOJA POOJA 2431420359 Patient ID:

Date and Time: 9th Nov 24 12:39 PM

THE PROPERTY OF THE PRINCE OF

N MM MA

Heart Rate 720 pm

74

Z

aVR

Gender Bornste

120-30 months 60 kg Patient Vitals Weight

168 cm NA Height: Pulse:

N.A. Spo2:

NA Respt

5

aVL

Others:

Measurementss

9/

23

aVF

Ħ

SOms ORSD:

413ms 378ms: QTeB: 6

96ms E

30.00 1999

P.R. T.

Tricog

ECG Within Normal Limits: Sinus Rhythm Sinus Arrhythmia Seen. Please correlate clinically.

25.8 m/s 10.0 mm/mV

H

REPORTED BY

P. Nith Soperate Carotha: Crestigne 8734