

# Arcofemi Healthcare Pvt Ltd

(Formerly known as Arcofemi Healthcare Ltd) F-701A, Lado Sarai, Mehrauli, New Delhi - 110030 Email: wellness@mediwheel.in, Website: www.mediwheel.in

Tel: +91-11-41195959, Fax: +91-11-29523020

CIN: U24240DL2011PTC216307

# **MEDICAL FITNESS CERTIFICATE**

(To be signed by a registered medical practitioner holding a Medical degree)

This is to certify that <u>Miss Swati Jhunjhunwala</u> aged, <u>22yr</u>. Based on the examination, I certify that he is in good mental and physical health and it is free from any physical defects such as deafness, colour blindness, and any chronic or contagious diseases.

Place: Mumbai

Date: 14/05/2024

Dr. Nitesh Kumar Mi-MBBB BCMR 47093

Name & Signature of

Medical officer

Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

**Episode** : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

## **Blood Bank**

Test Name Result

Sample No: 00331737A Collection Date: 14/05/24 08:55 Ack Date: 14/05/2024 10:38 Report Date: 14/05/24 12:49

BLOOD GROUPING/ CROSS-MATCHING BY SEMI AUTOMATION				
BLOOD GROUP (ABO)	'B'			
Rh Type  Method - Column Agglutination	POSITIVE			

REMARK: THE REPORTED RESULTS PERTAIN TO THE SAMPLE RECEIVED AT THE BLOOD CENTRE.

## Interpretation:

Blood typing is used to determine an individual's blood group, to establish whether a person is blood group A, B, AB, or O and whether he or she is Rh positive or Rh negative. Blood typing has the following significance,

- Ensure compatibility between the blood type of a person who requires a transfusion of blood or blood components and the ABO and Rh type of the unit of blood that will be transfused.
- Determine compatibility between a pregnant woman and her developing baby (fetus). Rh typing is especially important during pregnancy because a mother and her fetus could be incompatible.
- Determine the blood group of potential blood donors at a collection facility.
- Determine the blood group of potential donors and recipients of organs, tissues, or bone marrow, as part of a workup for a transplant procedure.

End of Report

Dr.Pooja Vinod Mishra MD Pathology

Jr Consultant Pathologist, MMC Reg No. 2017052191

RegNo: 2017/05/2191



Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

UHID : SHHM.94476 : 14/05/2024 08:31 **Order Date** 

: OP Episode

**Mobile No** Ref. Doctor : Self : 9674611213 DOB : 13/12/2001

: SEVENHILLS HOSPITAL, MUMBAI

**Facility** 

Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

Episode : OP

**Ref. Doctor**: Self Mobile No: 9674611213

2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018

**DOB** : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

# **Biochemistry**

Test Name			Resu	lt		Unit	Bio	logical Reference Interval
Sample No :	O0331737C	Collection Date :	14/05/24 08	:55	Ack Date :	14/05/2024 09:25	Report Date :	14/05/24 12:21
BUN-SERU	<u>JM</u>							
BUN - SERU Method - Urea				8.36			mg/dl	4 - 18
References: 1)Pack Inser	t of Bio system							

End of Report -

Dr.Ritesh Kharche MD, PGD-HM

Consultant Pathologist and Director of Laboratory Services

RegNo: 2006/03/1680



Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

**Episode** : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

Facility: SEVENHILLS HOSPITAL, MUMBAI

## **HAEMATOLOGY**

Test Name			Resul	t	Unit	Bio	logical Reference Interval
Sample No :	O0331737A	Collection Date :	14/05/24 08:	Ack Date :	14/05/2024 09:25	Report Date :	14/05/24 09:46
COMPLETI	BLOOD COUNT	Γ (CBC) - EDTA	WHOLE BLO	OOD			
Total WBC	Count			8.18		x10^3/ul	4 - 10
Neutrophils				65.7		%	40 - 80
Lymphocyte	25			29.6		%	20 - 40
Eosinophils				0.8 ▼ (L)		%	1 - 6
Monocytes				3.7		%	2 - 10
Basophils				0.2 ▼ (L)		%	1 - 2
Absolute Ne	eutrophil Count			5.37		x10^3/ul	2 - 7
Absolute Ly	mphocyte Count			2.42		x10^3/ul	0.8 - 4
Absolute Ed	sinophil Count			0.06		x10^3/ul	0.02 - 0.5
Absolute Mo	onocyte Count			0.31		x10^3/ul	0.12 - 1.2
Absolute Ba	sophil Count			0.02		x10^3/ul	0 - 0.1
RBCs				<b>4.22</b> ▼ (L)		x10^6/ul	4.5 - 5.5
Hemoglobin	l			12.4		gm/dl	12 - 15
Hematocrit				36.9 ▼ (L)		%	40 - 50
MCV				87.4		fl	83 - 101
MCH				29.5		pg	27 - 32



Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

Episode : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

MCHC	33.7	gm/dl	31.5 - 34.5
RED CELL DISTRIBUTION WIDTH-CV (RDW-CV)	13.4	%	11 - 16
RED CELL DISTRIBUTION WIDTH-SD (RDW-SD)	43.8	fl	35 - 56
Platelet	312	x10^3/ul	150 - 410
Mean Platelet Volume (MPV)	8.3	fl	6.78 - 13.46
PLATELET DISTRIBUTION WIDTH (PDW)	15.6	%	9 - 17
PLATELETCRIT (PCT)	0.259	%	0.11 - 0.28
Comment	PS Findings RBCs = Normocytic Normochromic WBCs = Normal Morphology Platelets =Adequate		

## Method:-

HB Colorimetric Method.

RBC/PLT Electrical Impedance Method.

WBC data Flow Cytometry by Laser Method.

MCV,MCH,MCHC,RDW and rest parameters - Calculated.

All Abnormal Haemograms are reviewed confirmed microscopically.

NOTE: Wallach's Interpretation of Diagnostic Tests. 11th Ed, Editors: Rao LV. 2021

## NOTE :-

The International Council for Standardization in Haematology (ICSH) recommends reporting of absolute counts of various WBC subsets for clinical decision making. This test has been performed on a fully automated 5 part differential cell counter which counts over 10,000 WBCs to derive differential counts. A complete blood count is a blood panel that gives information about the cells in a patient's blood, such as the cell count for each cell type and the concentrations of Hemoglobin and platelets. The cells that circulate in the bloodstream are generally divided into three types: white blood cells (leukocytes), red blood cells (erythrocytes), and platelets (thrombocytes). Abnormally high or low counts may be physiological or may indicate disease conditions, and hence need to be interpreted clinically.



Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

Episode : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

- End of Report

Dr.Pooja Vinod Mishra

**MD Pathology**Jr Consultant Pathologist, MMC Reg No.

2017052191 RegNo: 2017/05/2191



ćT; 14/05/2024 08:50:00 SEVENHILLS HEALTHCARE 100B F 50~ 0.50-100 Hz W OPD .....normal P axis, V-rate 50-99 .....SV complex w/ short R-R interval 75 Chest: 10.0 mm/mV - OTHERWISE NORMAL ECG -Limb: 10 mm/mV swati jhunjhunwala Atrial premature complex..... Sinus rhythm.... Female aVR aVL Speed: 25 mm/sec Standard Placement 137 79 376 429 64

Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

Episode : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

Facility: SEVENHILLS HOSPITAL, MUMBAI

#### **HAEMATOLOGY**

Test Name		Result		Unit	Biol	ogical Reference Inter	val		
	Sample No :	O0331737A	Collection Date :	14/05/24 08:55	Ack Date :	14/05/2024 09:25	Report Date :	14/05/24 10:45	

ERYTHROCYTE SEDIMENTATION RATE (ESR)			
ESR	<b>50</b> ▲ (H)	mm/hr	0 - 20

Method: Westergren Method

#### INTERPRETATION :-

ESR is a non-specific phenomenon, its measurement is clinically useful in disorders associated with an increased production of acute-phase proteins. It provides an index of progress of the disease in rheumatoid arthritis or tuberculosis, and it is of considerable value in diagnosis of temporal arteritis and polymyalgia rheumatica. It is often used if multiple myeloma is suspected, but when the myeloma is non-secretory or light chain, a normal ESR does not exclude this diagnosis.

An elevated ESR may occur as an early feature in myocardial infarction. Although a normal ESR cannot be taken to exclude the presence of organic disease, the vast majority of acute or chronic infections and most neoplastic and degenerative diseases are associated with changes in the plasma proteins that increased ESR values.

The ESR is influenced by age, stage of the menstrual cycle and medications taken (corticosteroids, contraceptive pills). It is especially low (0–1 mm) in polycythaemia, hypofibrinogenaemia and congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis, or sickle cells. In cases of performance enhancing drug intake by athletes the ESR values are generally lower than the usual value for the individual and as a result of the increase in haemoglobin (i.e. the effect of secondary polycythaemia).

- End of Report -

Dr.Ritesh Kharche MD, PGD-HM

Consultant Pathologist and Director of Laboratory Services

RegNo: 2006/03/1680

Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

**Ref. Doctor** : Self **Mobile No** : 9674611213

: OP

**Episode** 

DOB : 13/12/2001

Facility: SEVENHILLS HOSPITAL, MUMBAI



Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

**Episode** : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

## **Biochemistry**

Test Name Result		Result		Unit	Biol	ogical Reference Interval	
Sample No: 00331737B	Collection Date :	14/05/24 08:55	Ack Date :	14/05/2024 09:25	Report Date :	14/05/24 12:21	

GLUCOSE-PLASMA-FASTING			
Glucose,Fasting	76.91	mg/dl	70 - 100

## American Diabetes Association Reference Range:

Normal: < 100 mg/dl

Impaired fasting glucose(Prediabetes): 100 - 126 mg/dl

Diabetes : >= 126 mg/dl

#### References:

1)Pack Insert of Bio system

2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018

## Interpretation :-

Conditions that can result in an elevated blood glucose level include: Acromegaly, Acute stress (response to trauma, heart attack, and stroke for instance), Chronic kidney disease, Cushing syndrome, Excessive consumption of food, Hyperthyroidism, Pancreatitis.

A low level of glucose may indicate hypoglycemia, a condition characterized by a drop in blood glucose to a level where first it causes nervous system symptoms (sweating, palpitations, hunger, trembling, and anxiety), then begins to affect the brain (causing confusion, hallucinations, blurred vision, and sometimes even coma and death). A low blood glucose level (hypoglycemia) may be

seen with:Adrenal insufficiency, Drinking excessive alcohol, Severe liver disease, Hypopituitarism, Hypothyroidism, Severe infections, Severe heart failure, Chronic kidney (renal) failure, Insulin overdose, Tumors that produce insulin (insulinomas), Starvation.

Sample No: 00331737C Collection Date: 14/05/24 08:55 Ack Date: 14/05/2024 09:25 Report Date: 14/05/24 12:21

ALT(SGPT) - SERUM			
SGPT (Alanine Transaminase) - SERUM  Method - IFCC	9.2	IU/L	0 - 34
References :			



Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

**Episode** : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

## 1)Pack Insert of Bio system

2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018

,			
Total Bilirubin - SERUM Method - Diazo	1.14	mg/dl	0 - 2
Direct Bilirubin SERUM  Method - Diazotization	<b>0.5 ▲</b> (H)	mg/dl	0 - 0.4
Indirect Bilirubin - Calculated  Method - Calculated	0.64	mg/dl	
CREATININE-SERUM			
Creatinine - SERUM  Method - Jaffes Kinetic	0.65	mg/dl	0.5 - 1.1

#### References:

- 1)Pack Insert of Bio system
- 2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018

#### Notes :-

Creatinine is a chemical waste molecule that is generated from muscle metabolism. Creatinine is produced from creatine, a molecule of major importance for energy production in muscles. Approximataly 1-2% of the body's creatine is converted to creatinine every day. Creatinine is transported through the bloodstream to the kidneys. The kidneys filter out host of the creatinine and dispose of it in the urine. The kidneys maintain the blood creatinine in a normal ranges. Creatinine has been found to be a fairly reliable indicator of kidney function.

GLUCOSE-PLASMA POST PRANDIAL			
Glucose, Post Prandial	100.53	mg/dl	70 - 140

## American Diabetes Association Reference Range:

Post-Prandial Blood Glucose:

Non- Diabetic: Up to 140mg/dL Pre-Diabetic: 140-199 mg/dL Diabetic: >200 mg/dL

#### References:

- 1)Pack Insert of Bio system
- 2) Tietz Textbook Of Clinical Chemistry And Molecular Diagnostics, 6th Ed, Editors: Rifai et al. 2018



Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

Episode : OP

**Ref. Doctor**: Self **Mobile No**: 9674611213

**DOB** : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

#### Interpretation :-

Conditions that can result in an elevated blood glucose level include: Acromegaly, Acute stress (response to trauma, heart attack, and stroke for instance), Chronic kidney disease, Cushing syndrome, Excessive consumption of food, Hyperthyroidism, Pancreatitis.

A low level of glucose may indicate hypoglycemia, a condition characterized by a drop in blood glucose to a level where first it causes nervous system symptoms (sweating, palpitations, hunger, trembling, and anxiety), then begins to affect the brain (causing confusion, hallucinations, blurred vision, and sometimes even coma and death). A low blood glucose level (hypoglycemia) may be

seen with:Adrenal insufficiency, Drinking excessive alcohol, Severe liver disease, Hypopituitarism, Hypothyroidism, Severe infections, Severe heart failure, Chronic kidney (renal) failure, Insulin overdose, Tumors that produce insulin (insulinomas), Starvation.

End of Report

Dr.Ritesh Kharche MD, PGD-HM

Consultant Pathologist and Director of Laboratory Services

RegNo: 2006/03/1680





Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

**Episode** : OP

**Ref. Doctor** : Self **Mobile No** : 9674611213

**DOB** : 13/12/2001

Facility: SEVENHILLS HOSPITAL, MUMBAI

# Urinalysis

Test Name Res	ult Unit	Bio	ological Reference Interval
Sample No: 00331737D Collection Date: 14/05/24 0	8:55 Ack Date : 14/05/2024 09:26	Report Date :	14/05/24 11:33
Physical Examination			
QUANTITY	20	ml	
Colour	Pale Yellow		
Appearance	Clear		
DEPOSIT	Absent		Absent
рН	Acidic		
Specific Gravity	1.010		
Chemical Examination			
Protein	Trace		Absent
Sugar	Absent		Absent
ketones	POSITIVE ( ++ )		Absent
Occult Blood	Trace		Negative
Bile Salt	Absent		Absent
Bile Pigments	Absent		Absent
Urobilinogen	NORMAL		Normal
NITRATE	Absent		Absent
LEUKOCYTES	Absent		Absent

Patient Name : Ms. SWATI JHUNJHUNWALA Age/Sex : 22 Year(s) / Female

Episode : OP

 Ref. Doctor
 : Self
 Mobile No
 : 9674611213

 DOB
 : 13/12/2001

**Facility**: SEVENHILLS HOSPITAL, MUMBAI

Microscopic Examination			
Pus cells	Occasional	/HPF	
Epithelial Cells	Occasional	/HPF	
RBC	3-5	/HPF	Absent
Cast	ABSENT	/LPF	Absent
Crystal	ABSENT	/HPF	Absent
Amorphous Materials	Absent		Absent
Yeast	Absent		Absent
Bacteria	Absent		Absent

End of Report -

Dr.Pooja Vinod Mishra MD Pathology

Jr Consultant Pathologist, MMC Reg No. 2017052191

RegNo: 2017/05/2191



## **DIAGNOSTICS REPORT**

Facility

: Ms. SWATI JHUNJHUNWALA Patient Name

Age/Sex : 22 Year(s)/Female

: SHHM.94476 UHID Ref. Doctor : Self

Address : KALINA, SANTACRUZ

EAST, Mumbai, Maharastra,

400029

Order Date : 14/05/2024 08:31 Report Date : 14/05/2024 13:30

MUMBAI

: SEVENHILLS HOSPITAL,

: 9674611213 Mobile

# X-RAY CHEST PA VIEW

Both lungs are clear.

The frontal cardiac dimensions are normal.

The pleural spaces are clear.

Both hilar shadows are normal in position and density.

No diaphragmatic abnormality is seen.

The soft tissues and bony thorax are normal.

IMPRESSION: No pleuroparenchymal lesion is seen.

Dr.Bhujang Pai MBBS,MD

Consultant RegNo: 49380