Name	: Mrs. VIJETA SINGH			
PID No.	: MED111491927	Register On	: 11/02/2023 8:13 AM	-
SID No.	: 80067252	Collection On	: 11/02/2023 9:14 AM	
Age / Sex	: 40 Year(s) / Female	Report On	: 11/02/2023 12:48 PM	me
Туре	: OP	Printed On	: 21/02/2023 1:13 PM	DIAGN

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	Unit	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING	'O' 'Positive'		
(Blood/Agglutination)			
Complete Blood Count With - ESR			
Haemoglobin (Blood/Spectrophotometry)	12.1	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (Blood/Numeric Integration of MCV)	36.9	%	37 - 47
RBC Count (Blood/Electrical Impedance)	4.00	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (Blood/ <i>Calculated</i>)	92.2	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (Blood/ <i>Calculated</i>)	30.3	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (Blood/ <i>Calculated</i>)	32.8	g/dL	32 - 36
RDW-CV (Calculated)	14.5	%	11.5 - 16.0
RDW-SD (Calculated)	46.79	fL	39 - 46
Total Leukocyte Count (TC) (Blood/ <i>Electrical Impedance</i>)	5630	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance and absorbance)	55.77	%	40 - 75
Lymphocytes (Blood/Impedance and absorbance)	29.15	%	20 - 45
Eosinophils (Blood/Impedance and absorbance)	5.16	%	01 - 06
Monocytes (Blood/Impedance and absorbance)	9.63	%	01 - 10







APPROVED BY

The results pertain to sample tested.

Page 1 of 7

Name	: Mrs. VIJETA SINGH			
PID No.	: MED111491927	Register On :	1/02/2023 8:13 AM	~
SID No.	: 80067252	Collection On :	11/02/2023 9:14 AM	
Age / Sex	: 40 Year(s) / Female	Report On :	11/02/2023 12:48 PM	medall
Туре	: OP	Printed On :	21/02/2023 1:13 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			
Investiga	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
Basophil (Blood/Imj	ls pedance and absorbance)	0.29	%	00 - 02
INTERP	RETATION: Tests done on Automa	ted Five Part cell coun	ter. All abnormal results ar	e reviewed and confirmed microscopically.
	e Neutrophil count pedance and absorbance)	3.14	10^3 / µl	1.5 - 6.6
Absolute (Blood/Imj	e Lymphocyte Count	1.64	10^3 / µl	1.5 - 3.5
Absolute (Blood/Im)	e Eosinophil Count (AEC) pedance)	0.29	10^3 / µl	0.04 - 0.44
Absolute (Blood/Im	e Monocyte Count	0.54	10^3 / µl	< 1.0
-	e Basophil count	0.02	10^3 / µl	< 0.2
Platelet (Blood/Im)	Count	1.45	lakh/cu.mm	1.4 - 4.5
-	RETATION: Platelet count less that	n 1.5 lakhs will be conf	irmed microscopically.	
MPV (Blood/De	rived from Impedance)	12.51	fL	8.0 - 13.3
PCT (Calculate	<i>d</i>)	0.18	%	0.18 - 0.28
	ythrocyte Sedimentation Rate) tomated ESR analyser)	28	mm/hr	< 20
BUN/C	Creatinine Ratio	8.7		
	Fasting (FBS) F/Glucose oxidase/Peroxidase)	110	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD - POD)	124	mg/dL	70 - 140
P.Y. Pradcep P.Venkata Pradcep Lab Manager VERIFIED BY	国際設計 新設施設 充分に対応 国家体計		APPROVED BY

The results pertain to sample tested.

Page 2 of 7

Name	: Mrs. VIJETA SINGH			
PID No.	: MED111491927	Register On	: 11/02/2023 8:13 AM	~
SID No.	: 80067252	Collection On	: 11/02/2023 9:14 AM	\mathbf{O}
Age / Sex	: 40 Year(s) / Female	Report On	: 11/02/2023 12:48 PM	medall
Туре	: OP	Printed On	: 21/02/2023 1:13 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Calculated)	7.9	mg/dL	7.0 - 21
Creatinine (Serum/ <i>Jaffe ⁻ Alkaline Picrate</i>)	1	mg/dL	0.6 - 1.1
Uric Acid (Serum/Uricase/Peroxidase)	4.9	mg/dL	2.6 - 6.0
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulphanilic acid)	0.7	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulphanilic acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Calculated)	0.50	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC without P-5-P)	33	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC without P-5-P)	26	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC AMP Buffer)	108	U/L	42 - 98
Total Protein (Serum/Biuret)	7.0	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.8	gm/dl	3.5 - 5.2
Globulin (Serum/Calculated)	3.20	gm/dL	2.3 - 3.6







APPROVED BY

The results pertain to sample tested.

Page 3 of 7

Name	: Mrs. VIJETA SINGH		
PID No.	: MED111491927	Register On : 11/02/2023 8:13 AM	
SID No.	: 80067252	Collection On : 11/02/2023 9:14 AM	
Age / Sex	: 40 Year(s) / Female	Report On : 11/02/2023 12:48 PM	medall
Туре	: OP	Printed On : 21/02/2023 1:13 PM	DIAGNOSTICS
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
A : G RATIO (Serum/Calculated)	1.19		1.1 - 2.2
INTERPRETATION: Enclosure : Graph GGT(Gamma Glutamyl Transpeptidase) (Serum/ <i>IFCC / Kinetic</i>)	23	U/L	< 38
Lipid Profile			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	167	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol-phosphate oxidase/Peroxidase)	89	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/ <i>Immunoinhibition</i>)	37	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/ <i>Calculated</i>)	112.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >=190
VLDL Cholesterol (Serum/Calculated)	17.8	mg/dL	< 30
P.V. Pradeep P.Venkata Pradeep Lab Manager Verified By	(1) (1) (1) (1) (1) (1) (1) (1) (1) (1)		K.Nuch out a Dr K. NEEHARIKA MD PATHOLOGY Reg No : 96545
			APPROVED BY

The results pertain to sample tested.

Page 4 of 7

Name	: Mrs. VIJETA SINGH			
PID No.	: MED111491927	Register On	: 11/02/2023 8:13 AM	
SID No.	: 80067252	Collection On	: 11/02/2023 9:14 AM	
Age / Sex	: 40 Year(s) / Female	Report On	: 11/02/2023 12:48 PM	medall
Туре	: OP	Printed On	: 21/02/2023 1:13 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			
Investiga	ation	Observe	<u>d Unit</u>	<u>Biological</u>

<u>Investigation</u>	Value	<u>Om</u>	Reference Interval
Non HDL Cholesterol (Serum/ <i>Calculated</i>)	130.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/ <i>Calculated</i>)	4.5		Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/ <i>Calculated</i>)	2.4		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	3		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
<u>Glycosylated Haemoglobin (HbA1c)</u>			
HbA1C (Whole Blood/ <i>HPLC-Ion exchange</i>)	6.1	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control : 6.1 - 7.0 %, Fair control : 7.1 - 8.0 %, Poor control >= 8.1 %Mean Blood Glucose 128.37 mg/dl

Mean Blood Glucose (Whole Blood)







APPROVED BY

The results pertain to sample tested.

Page 5 of 7

Name	: Mrs. VIJETA SINGH			
PID No.	: MED111491927	Register On : 1	1/02/2023 8:13 AM	
SID No.	: 80067252	Collection On : 1	1/02/2023 9:14 AM	\mathbf{O}
Age / Sex	: 40 Year(s) / Female	Report On : 1	1/02/2023 12:48 PM	medall
Туре	: OP	-	1/02/2023 1:13 PM	DIAGNOSTICS
Ref. Dr	: MediWheel		.,	
Investiga	ition	<u>Observed</u> <u>Value</u>	Unit	<u>Biological</u> Reference Interval
INTERPRETATION: Comments HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations. Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency, hypertriglyceridemia,hyperbilirubinemia,Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values. Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly,Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.				
<u>THYROI</u>	<u>D PROFILE / TFT</u>			
	dothyronine) - Total emiluminescent Immunometric Assay	0.97	ng/ml	0.7 - 2.04
Comment	ariation can be seen in other condition	on like pregnancy, drugs	s, nephrosis etc. In such ca	ases, Free T3 is recommended as it is
T4 (Thyr	oxine) - Total emiluminescent Immunometric Assay	8.91	µg/dl	4.2 - 12.0
Comment	ariation can be seen in other condition	on like pregnancy, drugs	s, nephrosis etc. In such ca	ases, Free T4 is recommended as it is
TSH (Th	yroid Stimulating Hormone) emiluminescence)	5.84	µIU/mL	0.35 - 5.50
Reference 1 st trimes 2 nd trimes 3 rd trimes (Indian Th Comment 1.TSH refe 2.TSH Lev be of the o 3.Values&	erence range during pregnancy depen	n, reaching peak levels b as influence on the mea	between 2-4am and at a m sured serum TSH concent	

Urine Analysis - Routine







APPROVED BY

The results pertain to sample tested.

Page 6 of 7

Name PID No. SID No. Age / Sex Type Ref. Dr	 : Mrs. VIJETA SINGH : MED111491927 : 80067252 : 40 Year(s) / Female : OP : MediWheel 	Collection On : Report On :	11/02/2023 8:13 AM 11/02/2023 9:14 AM 11/02/2023 12:48 PM 21/02/2023 1:13 PM	DIAGNOSTICS
<u>Investiga</u>	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
	croscopy) RETATION:Note: Done with Auto Examination(Urine Routine)	NIL	& microscopy	
Colour	vsical examination)	PALE YELLO	W	Yellow to Amber
Appeara		Clear		Clear
	ul Examination(Urine Routine	<u>e)</u>		
	ostick-Error of indicator/ cylic acid method)	Negative		Negative
	o Stick Method / Glucose Oxidase - e / Benedict š semi quantitative	Negative		Negative
<u>Microsco</u> <u>Routine</u>)	opic Examination(Urine)			
Pus Cells (Urine/Mic	S croscopy exam of urine sediment)	2-3	/hpf	0 - 5
Epithelia		1-2	/hpf	NIL
RBCs (Urine/Mid	croscopy exam of urine sediment)	NIL	/hpf	0 - 5
	P.V. Pradece ab Manager erified by		5870 2007 14-14 15-15 10-15	K.Nukouka Dr.K. NEEHARIKA MD PATHOLOGY Reg No : 96545

VERIFIED BY

-- End of Report --



APPROVED BY

The results pertain to sample tested.

Page 7 of 7



Name	VIJETA SINGH	ID	MED111491927
Age & Gender	40Y/F	Visit Date	Feb 11 2023 8:12AM
Ref Doctor	MediWheel		

ULTRASOUND WHOLE ABDOMEN

Liver :	Normal in size (14.3 cm) with regular outlines and normal echopattern. There is no evidence of IHBR / EHBR dilatation seen. No focal space occupying lesions seen. CBD is normal. PV normal.			
Gall Bladder :	Normal in volume and wall thickness. No e/o intraluminal calculi seen.			
Pancreas :	Head, body and tail are identified with normal echopattern and smooth outlines.			
Spleen :	Measured 10.3 cm, in size with normal echotexture.			
Right kidney :	Measured 9.5 x 4.0 cm in size.			
Left kidney :	Measured 9.8 x 4.1 cm in size. Both kidneys are normal in size, position, with well preserved cortico medullary differentiation and normal pelvicalyceal anatomy. No e/o calculi / space occupying lesion seen. No e/o suprarenal / retroperitoneal masses noted.			
Urinary bladder	: Normal in volume and wall thickness. No e/o intraluminal calculi / masses seen.			
Uterus :	Measured 7.1 x 4.0 x 5.0 cm in size with normal myometrial and endometrial echotexture.			
- Intra uterine contraceptive device seen insitu.				
Right ovary : Left ovary :	Measured $3.1 \ge 2.3$ cm in size. Measured $3.0 \ge 1.5$ cm in size. Both ovaries are normal in size and appearance.			
	No e/o ascites / pleural effusion seen. No e/o detectable bowel pathology seen.			



Name	VIJETA SINGH	ID	MED111491927
Age & Gender	40Y/F	Visit Date	Feb 11 2023 8:12AM
Ref Doctor	MediWheel		

IMPRESSION :

- Intra uterine contraceptive device insitu.
- Rest of the study is essentially normal.

- For clinical correlation.

Dr.Jahn avi Barla, MD (RD)

Consultant Radiologist