

LEELAVATI LAHANE

CLIENT'S NAME AND ADDRESS :







SRL Ltd 74,PASHCHIMI MARG,VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
DRAWN : 08/04/2022 09:13	RECEIVED : 08/04/2022 09:15	REPORTED : 09/04/2022 10:18
ACCESSION NO : 0063VD001703	AGE : 37 Years SEX : Female	
PATIENT NAME : LEELAVATI LAH	IANE	PATIENT ID : LEELF09038563
	Email : cu	istomercare.palammarg@srl.in

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

BLOOD COUNTS,EDTA WHOLE BLOOD				
HEMOGLOBIN	11.1	Low	12.0 - 15.0	g/dL
RED BLOOD CELL COUNT	4.37		3.8 - 4.8	mil/µL
WHITE BLOOD CELL COUNT	6.58		4.0 - 10.0	thou/µL
PLATELET COUNT	237		150 - 410	thou/µL
RBC AND PLATELET INDICES				
HEMATOCRIT	35.8	Low	36 - 46	%
MEAN CORPUSCULAR VOL	82.0	Low	83 - 101	fL
MEAN CORPUSCULAR HGB.	25.4	Low	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN	30.9	Low	31.5 - 34.5	g/dL
CONCENTRATION MENTZER INDEX	18.8			
RED CELL DISTRIBUTION WIDTH	14.1	High	11.6 - 14.0	%
MEAN PLATELET VOLUME	12.6	_	6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT - NLR		_		
SEGMENTED NEUTROPHILS	63		40 - 80	%
ABSOLUTE NEUTROPHIL COUNT	4.15		2.0 - 7.0	thou/µL
LYMPHOCYTES	26		20 - 40	%
ABSOLUTE LYMPHOCYTE COUNT	1.71		1 - 3	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	2.4			
EOSINOPHILS	3		1 - 6	%
ABSOLUTE EOSINOPHIL COUNT	0.20		0.02 - 0.50	thou/µL
MONOCYTES	7		2 - 10	%
ABSOLUTE MONOCYTE COUNT	0.46		0.20 - 1.00	thou/µL
BASOPHILS	1		0 - 2	%
ABSOLUTE BASOPHIL COUNT	0.07		0.02 - 0.10	thou/µL
DIFFERENTIAL COUNT PERFORMED ON:	EDTA SMEAR			
METHOD : AUTOMATED ANALYZER / MICROSCOPY				

DISCLAIMER: THE ABSOLUTE WHITE CELL COUNTS ARE OUTSIDE THE NABL ACCREDITED SCOPE OF THE LABORATORY.


ERYTHRO SEDIMENTATION RATE, BLOOD

SEDIMENTATION RATE (ESR)	30	High 0 - 20
METHOD : MODIFIED WESTERGREN		









METHOD : CALCULATED PARAMETER







LEELAVATI LAHANE

CLIENT'S NAME AND ADDRESS :







LEELF09038563

SRL Ltd 74, PASHCHIMI MARG, VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

REPORTED :

PATIENT NAME : LEELAVATI LAHANE ACCESSION NO : 0063VD001703 AGE : 37 Years SEX : Female

DRAWN : 08/04/2022 09:13 RECEIVED : 08/04/2022 09:15

REFERRING DOCTOR : SELF

Test Report Status <u>Final</u>	Results		Biological Reference Interva	al Units
	2.7	Low	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk	
METHOD : CALCULATED PARAMETER LDL/HDL RATIO	1.6		0.5-3 Desirable/Low risk	
			3.1-6 Borderline/Moderate risk >6.0 High Risk	
VERY LOW DENSITY LIPOPROTEIN	14.0		= 30</td <td>mg/dL</td>	mg/dL
METHOD : CALCULATED PARAMETER * LIVER FUNCTION PROFILE, SERUM				
	0.34		Linta 1 D	m a (dl
BILIRUBIN, TOTAL METHOD : SPECTROPHOTOMETRY	0.34		Upto 1.2	mg/dL
BILIRUBIN, DIRECT	0.15		Upto 0.2	mg/dL
METHOD : SPECTROPHOTOMETRY	0.15		000012	iiig/ dE
BILIRUBIN, INDIRECT	0.19		0.00 - 0.60	mg/dL
METHOD : CALCULATED PARAMETER				5,
TOTAL PROTEIN	8.1		6.4 - 8.3	g/dL
METHOD : SPECTROPHOTOMETRY				
ALBUMIN	4.6		3.70 - 4.94	g/dL
METHOD : SPECTROPHOTOMETRY				
GLOBULIN	3.5		2.0 - 4.0	g/dL
METHOD : CALCULATED PARAMETER				
ALBUMIN/GLOBULIN RATIO	1.3		1.0 - 2.0	RATIO
METHOD : CALCULATED PARAMETER				
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	14		0 - 32	U/L
	10		2 22	
ALANINE AMINOTRANSFERASE (ALT/SGPT)	10		0 - 33	U/L
METHOD : SPECTROPHOTOMETRY ALKALINE PHOSPHATASE	85		35 - 104	11/1
METHOD : SPECTROPHOTOMETRY	60		55 - 104	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT)	13		5 - 36	U/L
METHOD : SPECTROPHOTOMETRY	15		3 - 30	0/L
	184		135 - 214	U/L
METHOD : SPECTROPHOTOMETRY				-, -
* SERUM BLOOD UREA NITROGEN				
BLOOD UREA NITROGEN	10		6 - 20	mg/dL



METHOD : SPECTROPHOTOMETRY



CLIENT PATIENT ID :

PATIENT ID :

09/04/2022 10:18



LEELAVATI LAHANE

CLIENT'S NAME AND ADDRESS :







SRL Ltd 74,PASHCHIMI MARG,VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

	Email :	customercare.palammarg@srl.in	
PATIENT NAME : LEELAVATI LAHANE	E	PATIENT ID :	LEELF09038563
ACCESSION NO : 0063VD001703 AG	E : 37 Years SEX : Female		
DRAWN : 08/04/2022 09:13	RECEIVED : 08/04/2022 09:15	REPORTED : 09/04/2022	2 10:18
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :	
Test Report Status <u>Final</u>	Results	Biological Reference Ir	iterval Units
* CREATININE, SERUM			
CREATININE	0.60	0.5 - 0.9	mg/dL
METHOD : SPECTROPHOTOMETRY			2.
* BUN/CREAT RATIO			
BUN/CREAT RATIO	16.67 H	High 5.00 - 15.00	
METHOD : CALCULATED PARAMETER			
URIC ACID, SERUM			
URIC ACID	3.9	2.4 - 5.7	mg/dL
METHOD : SPECTROPHOTOMETRY			
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	8.1	6.4 - 8.3	g/dL
METHOD : SPECTROPHOTOMETRY			
ALBUMIN, SERUM			
ALBUMIN	4.6	3.97 - 4.94	g/dL
METHOD : SPECTROPHOTOMETRY			
* GLOBULIN			
GLOBULIN	3.5	2.0 - 4.0	g/dL
METHOD : CALCULATED PARAMETER			
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM	144	136 - 145	mmol/L
METHOD : SPECTROPHOTOMETRY			
	4.34	3.3 - 5.1	mmol/L
	102	09 106	mmal/l
	103	98 - 106	mmol/L
METHOD : SPECTROPHOTOMETRY URINALYSIS			
UNINAL I 313			

CHLORIDE	105	98 - 100
METHOD : SPECTROPHOTOMETRY		
URINALYSIS		
COLOR	PALE YELLOW	
METHOD : MACROSCOPY		
APPEARANCE	Clear	
METHOD : VISUAL EXAMINATION		
PH	6.0	4.7 - 7.5
METHOD : PH INDICATOR AND REFLECTANCE, SPECTROPHOTOM	ETRY	
SPECIFIC GRAVITY	1.010	1.003 - 1.035
METHOD : PKA CHANGE WITH REFLECTANCE, SPECTROPHOTOME	ſŖŶ	
GLUCOSE	NOT DETECTED	NOT DETECTED
METHOD : GLUCOSE OXIDASE WITH REFLECTANCE, SPECTROPHC	TOMETRY	





Scan to View Details



LEELAVATI LAHANE

CLIENT'S NAME AND ADDRESS :







SRL Ltd 74,PASHCHIMI MARG,VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

PATIENT NAME : LEELAVATI LAH	IANE	PATIENT ID : LEELF09038563
ACCESSION NO : 0063VD001703	AGE : 37 Years SEX : Female	
DRAWN : 08/04/2022 09:13	RECEIVED : 08/04/2022 09:15	REPORTED : 09/04/2022 10:18
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
PROTEIN	NOT DETECTED	NOT DETECTED
		NOT DETECTED
METHOD : PROTEIN ERROR OF INDICATORS V	ITH REFLECTANCE, SPECTROPHOTOMETRY	
KETONES	NOT DETECTED	NOT DETECTED

PROTEIN	NOT DETECTED	NOT DETECTED	
METHOD : PROTEIN ERROR OF INDICATORS WITH REFLECTANCE, SI	PECTROPHOTOMETRY		
KETONES	NOT DETECTED	NOT DETECTED	
METHOD : ROTHERA'S WITH REFLECTANCE, SPECTROPHOTOMETRY			
BLOOD	NOT DETECTED	NOT DETECTED	
METHOD : PEROXIDASE METHOD WITH REFLECTANCE, SPECTROPHO	DTOMETRY		
BILIRUBIN	NOT DETECTED	NOT DETECTED	
METHOD : DIAZOTIZED WITH REFLECTANCE, SPECTROPHOTOMETR	Y		
UROBILINOGEN	NORMAL	NORMAL	
METHOD : EHRLICH REACTION WITH REFLECTANCE, SPECTROPHOT	OMETRY		
NITRITE	NOT DETECTED	NOT DETECTED	
METHOD : DIAZONIUM COMPOUND WITH REFLECTANCE, SPECTROP	HOTOMETRY		
PUS CELL (WBC'S)	2-3	0-5	/HPF
EPITHELIAL CELLS	2-3	0-5	/HPF
ERYTHROCYTES (RBC'S)	NOT DETECTED	NOT DETECTED	/HPF
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	

NOTE:-MICROSCOPIC EXAMINATION OF URINE PERFORMED BY CENTRIFUGED URINARY SEDIMENT

* THYROID PANEL, SERUM			
T3	117.8	80.00 - 200.00	ng/dl
METHOD : ELECTROCHEMILUMINESCENCE	117.0	80.00 - 200.00	ng/dL
	0.07	5.10 - 14.10	
Τ4	8.07	5.10 - 14.10	µg/dL
METHOD : ELECTROCHEMILUMINESCENCE			
TSH 3RD GENERATION	2.590	0.270 - 4.200	µIU/mL
STOOL: OVA & PARASITE			
COLOUR	SAMPLE NOT RECEIVED		
METHOD : MANUAL			
* ABO GROUP & RH TYPE, EDTA WHOLE BLOOD			
ABO GROUP	TYPE B		
METHOD : MANUAL			
RH TYPE	POSITIVE		



METHOD : MANUAL
* XRAY-CHEST

REMARKS





LEELAVATI LAHANE

CLIENT'S NAME AND ADDRESS :







SRL Ltd 74,PASHCHIMI MARG,VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

PATIENT NAME : LEELAVATI LAHANE PATIENT ID : LEELF09038563 ACCESSION NO : 0063VD001703 AGE: 37 Years SEX : Female DRAWN : 08/04/2022 09:13 RECEIVED : 08/04/2022 09:15 REPORTED : 09/04/2022 10:18 REFERRING DOCTOR : SELF CLIENT PATIENT ID : **Test Report Status** Results **Biological Reference Interval** Units <u>Final</u>

IMPRESSION	NORMAL		
TMT OR ECHO			
TMT OR ECHO	TMT DONE		
ECG			
ECG	SINUS TACHYCARDIA RES	T WITHIN NORMAL LIMITS	
* MEDICAL HISTORY			
RELEVANT PRESENT HISTORY	NOT SIGNIFICANT		
RELEVANT PAST HISTORY	H/O COVID IN APRIL 2021	l.	
RELEVANT PERSONAL HISTORY	UNMARRIED.		
RELEVANT FAMILY HISTORY	FATHER DIABETES. MOTHER DIED OF CARDIA	C ARREST.	
OCCUPATIONAL HISTORY	NOT SIGNIFICANT		
HISTORY OF MEDICATIONS	NOT SIGNIFICANT		
* ANTHROPOMETRIC DATA & BMI			
HEIGHT IN METERS	1.58		mts
WEIGHT IN KGS.	56		Kgs
BMI	22	BMI & Weight Status as follows Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese	s: kg/sqmts
* GENERAL EXAMINATION			
MENTAL / EMOTIONAL STATE	NORMAL		
PHYSICAL ATTITUDE	NORMAL		
GENERAL APPEARANCE / NUTRITIONAL STATUS	HEALTHY		
BUILT / SKELETAL FRAMEWORK	AVERAGE		
FACIAL APPEARANCE	NORMAL		
SKIN	NORMAL		
UPPER LIMB	NORMAL		
LOWER LIMB	NORMAL		
NECK	NORMAL		
NECK LYMPHATICS / SALIVARY GLANDS	NOT ENLARGED OR TENDI	ER	
THYROID GLAND	NOT ENLARGED		
CAROTID PULSATION	NORMAL		
BREAST (FOR FEMALES)	NORMAL		











PATIENT ID :

CLIENT PATIENT ID :

09/04/2022 10:18



LEELF09038563

SRL Ltd 74,PASHCHIMI MARG,VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

REPORTED :

CLIENT'S NAME AND ADDRESS : LEELAVATI LAHANE

CLIENT CODE : C000138377

 PATIENT NAME : LEELAVATI LAHANE

 ACCESSION NO :
 0063VD001703
 AGE : 37 Years
 SEX : Female

 DRAWN :
 08/04/2022 09:13
 RECEIVED : 08/04/2022 09:15

 REFERRING DOCTOR :
 SELF

Test Report Status Results **Biological Reference Interval** Units **Final** NORMAL TEMPERATURE REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID BRUIT PULSE RESPIRATORY RATE NORMAL * CARDIOVASCULAR SYSTEM BΡ 144/84 MM HG mm/Hg (SITTING) PERICARDIUM NORMAL APEX BEAT NORMAL HEART SOUNDS NORMAL MURMURS ABSENT *** RESPIRATORY SYSTEM** SIZE AND SHAPE OF CHEST NORMAL MOVEMENTS OF CHEST SYMMETRICAL BREATH SOUNDS INTENSITY NORMAL BREATH SOUNDS QUALITY VESICULAR (NORMAL) ADDED SOUNDS ABSENT *** PER ABDOMEN** APPEARANCE NORMAL VENOUS PROMINENCE ABSENT NOT PALPABLE LIVER NOT PALPABLE SPLEEN *** CENTRAL NERVOUS SYSTEM** HIGHER FUNCTIONS NORMAL CRANIAL NERVES NORMAL CEREBELLAR FUNCTIONS NORMAL SENSORY SYSTEM NORMAL MOTOR SYSTEM NORMAL REFLEXES NORMAL * MUSCULOSKELETAL SYSTEM SPINE NORMAL JOINTS NORMAL *** BASIC EYE EXAMINATION** CONJUNCTIVA NORMAL **EYELIDS** NORMAL







LEELAVATI LAHANE

CLIENT'S NAME AND ADDRESS :







SRL Ltd 74, PASHCHIMI MARG, VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

REPORTED :

PATIENT NAME : LEELAVATI LAHANE

PATIENT ID : LEELF09038563

09/04/2022 10:18

CLIENT PATIENT ID :

ACCESSION NO : 0063VD001703 AGE: 37 Years SEX : Female

DRAWN: 08/04/2022 09:13 RECEIVED : 08/04/2022 09:15

REFERRING DOCTOR : SELF

CLIENT FATIENT ID .			
Test Report Status <u>Final</u>	Results	Biological Reference Interval	Units
EYE MOVEMENTS	NORMAL		
CORNEA	NORMAL		
DISTANT VISION RIGHT EYE WITH GLASSES	6/6		
DISTANT VISION LEFT EYE WITH GLASSES	6/6		
NEAR VISION RIGHT EYE WITH GLASSES	N6		
NEAR VISION LEFT EYE WITH GLASSES	N6		
COLOUR VISION	NORMAL		
* BASIC ENT EXAMINATION			
EXTERNAL EAR CANAL	NORMAL		
TYMPANIC MEMBRANE	NORMAL		
NOSE	NO ABNORMALITY D	DETECTED	
SINUSES	NORMAL		
THROAT	NO ABNORMALITY D	DETECTED	
TONSILS	NOT ENLARGED		
* SUMMARY			
RELEVANT HISTORY	NOT SIGNIFICANT		
RELEVANT GP EXAMINATION FINDINGS	NOT SIGNIFICANT		
RELEVANT LAB INVESTIGATIONS	LOW HB INCRASE IN ESR		
RELEVANT NON PATHOLOGY DIAGNOSTICS	ATTACHED		
REMARKS / RECOMMENDATIONS	PHYSICIAN CONSUL	.T	
* FITNESS STATUS			
FITNESS STATUS	FIT (WITH MEDICAL	ADVICE) (AS PER REQUESTED PANEL OF	TESTS)

FITNESS STATUS

FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

Interpretation(s) BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-

Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT - NLR-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease. (Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients ; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504

This ratio element is a calculated parameter and out of NABL scope. ERYTHRO SEDIMENTATION RATE, BLOOD-

Erythrocyte sedimentation rate (ESR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by







Scan to View Details



I FEI AVATT I AHANE

CLIENT'S NAME AND ADDRESS :







SRL Ltd 74, PASHCHIMI MARG, VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.palammarg@srl.in **PATIENT NAME : LEELAVATI LAHANE** PATIENT ID : LEELF09038563 ACCESSION NO : 0063VD001703 AGE: 37 Years SEX : Female DRAWN: 08/04/2022 09:13 RECEIVED : 08/04/2022 09:15 **REPORTED** : 09/04/2022 10:18 REFERRING DOCTOR : SELF CLIENT PATIENT ID : Test Report Status Results Biological Reference Interval Units **Final**

age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0 -1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

Reference :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin

3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition"

GLUCOSE, FASTING, PLASMA-ADA 2021 guidelines for adults, after 8 hrs fasting is as follows: Pre-diabetics: 100 - 125 mg/dL Diabetic: > or = 126 mg/dL GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-

Glycosylated hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood, the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells. Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia, increased red cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of the testing average decide the some decidered

"Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations.'

References

1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006, 879-884.

 Forsham PH. Diabetes Mellitus: A rational plan for management. Postgrad Med 1982, 71,139-154.
 Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184. GLUCOSE, POST-PRANDIAL, PLASMA-ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75 grams of glucose in 300 ml water, over a period of 5 minutes.

CORONARY RISK PROFILE (LIPID PROFILE), SERUM.-

Serum cholesterol is a blood test that can provide valuable information for the risk of coronary artery disease This test can help determine your risk of the build up of plaques in your arteries that can lead to narrowed or blocked arteries throughout your body (atherosclerosis). High cholesterol levels usually don't cause any signs or symptoms, so a cholesterol test is an important tool. High cholesterol levels often are a significant risk factor for heart disease and important for diagnosis of hyperlipoproteinemia, atherosclerosis, hepatic and thyroid diseases.

Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn't need into triglycerides, which are stored in fat cells. High triglyceride levels are associated with several factors, including being overweight, eating too many sweets or drinking too much alcohol, smoking, being sedentary, or having diabetes with elevated blood sugar levels. Analysis has proven useful in the diagnosis and treatment of patients with diabetes mellitus, nephrosis, liver obstruction, other diseases involving lipid metabolism, and various endocrine disorders. In conjunction with high density lipoprotein and total serum cholesterol, a triglyceride determination provides valuable information for the assessment of coronary heart disease risk. It is done in fasting state.

High-density lipoprotein (HDL) cholesterol. This is sometimes called the ""good"" cholesterol because it helps carry away LDL cholesterol, thus keeping arteries open and blood flowing more freely. HDL cholesterol is inversely related to the risk for cardiovascular disease. It increases following regular exercise, moderate alcohol consumption and with oral estrogen therapy. Decreased levels are associated with obesity, stress, cigarette smoking and diabetes mellitus.

SERUM LDL The small dense LDL test can be used to determine cardiovascular risk in individuals with metabolic syndrome or established/progressing coronary artery disease, individuals with triglyceride levels between 70 and 140 mg/dL, as well as individuals with a diet high in trans-fat or carbohydrates. Elevated sdLDL levels are associated with metabolic syndrome and an 'atherogenic lipoprotein profile', and are a strong, independent predictor of cardiovascular disease. Elevated levels of LDL arise from multiple sources. A major factor is sedentary lifestyle with a diet high in saturated fat. Insulin-resistance and pre-diabetes have also been implicated, as has genetic predisposition. Measurement of sdLDL allows the clinician to get a more comprehensive picture of lipid risk factors and tailor treatment accordingly. Reducing LDL levels will reduce the risk of CVD and MI.

Non HDL Cholesterol - Adult treatment panel ATP III suggested the addition of Non-HDL Cholesterol as an indicator of all atherogenic lipoproteins (mainly LDL and VLDL). NICE guidelines recommend Non-HDL Cholesterol measurement before initiating lipid lowering therapy. It has also been shown to be a better marker of risk in both primary and secondary prevention studies.

Recommendations:

Results of Lipids should always be interpreted in conjunction with the patient's medical history, clinical presentation and other findings.

NON FASTING LIPID PROFILE includes Total Cholesterol, HDL Cholesterol and calculated non-HDL Cholesterol. It does not include triglycerides and may be best used in patients for whom fasting is difficult.













CLIENT'S NAME AND ADDRESS : I FEI AVATT I AHANE

SRL Ltd 74, PASHCHIMI MARG, VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

PATIENT NAME : LEELAVATI LAH	ANE	PATIENT ID : LEELF09038563
ACCESSION NO : 0063VD001703	AGE : 37 Years SEX : Female	
DRAWN : 08/04/2022 09:13	RECEIVED : 08/04/2022 09:15	REPORTED : 09/04/2022 10:18
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
Test Report Status Final	Results	Biological Reference Interval Units

LIVER FUNCTION PROFILE, SERUM-

LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver,liver cancer,kidney failure,hemolytic anemia,pancreatitis,hemochromatosis. AST levels may also increase after a heart attack or strenuous activity.ALT test measures the amount of this enzyme in the blood.ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis. ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction,

Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease,Rickets,Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia,Malnutrition,Protein deficiency,Wilson's disease.GGT is an enzyme found in cell membranes of many tissues mainly in the liver,kidney and pancreas.It is also found in other tissues including intestine,spleen,heart, brain and seminal vesicles.The highest concentration is in the kidney,but the liver is considered the source of normal enzyme activity.Serum GGT has been widely used as an index of liver dysfunction.Elevated serum GGT activity can be found in diseases of the liver,biliary system and pancreas.Conditions that increase serum GGT are obstructive liver disease,high alcohol consumption and use of enzyme-inducing drugs etc.Serum total protein,also known as total protein,is a biochemical test for measuring the total amount of protein in serum.Protein in the plasma is made up of albumin and globulin.Higher-than-normal levels may be due to:Chronic inflammation or infection,including HIV and hepatitis B or C,Multiple myeloma,Waldenstrom's disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome,Protein-losing enteropathy etc.Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc SERUM BLOOD UREA NITROGEN-

Causes of Increased levels

Pre renal

High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal

Renal Failure

Post Renal • Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

Liver diseaseSIADH.

CREATININE, SERUM-

- Higher than normal level may be due to:
 Blockage in the urinary tract
 Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

 Mvasthenia Gravis Muscular dystrophy URIC ACID, SERUM-Causes of Increased levels Dietary • High Protein Intake. Prolonged Fasting, Rapid weight loss Gout

Lesch nyhan syndrome. Type 2 DM. Metabolic syndrome.

Causes of decreased levels

Low Zinc Intake
OCP's

Multiple Sclerosis













CLIENT'S NAME AND ADDRESS : I FEI AVATT I AHANE

SRL Ltd 74, PASHCHIMI MARG, VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

	CLIENT PATIENT ID :
RECEIVED : 08/04/2022 09:15	REPORTED : 09/04/2022 10:18
3 AGE : 37 Years SEX : Female	
AHANE	PATIENT ID : LEELF09038563

Nutritional tips to manage increased Uric acid levels

Drink plenty of fluids

Limit animal proteins

High Fibre foods

Vit C IntakeAntioxidant rich foods

TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum.. Protein in the plasma is made up of albumin and alobulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. ALBUMIN, SERUM-

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

ELECTROLYTES (NA/K/CL), SERUM-Sodium levels are Increased in dehydration, cushing's syndrome, aldosteronism & decreased in Addison's disease, hypopituitarism, liver disease. Hypokalemia (low K) is Addison's disease, metabolic acidosis, acute starvation, dehydration, and with rapid K infusion.Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bicarbonate, diabetes insipidus, adrenocortical hyperfuction, salicylate intoxication and with excessive infusion of isotonic saline or extremely high dietary intake of salt.Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and metabolic acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and prolonged vomiting,

URINALYSIS-Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus. Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia

THYROID PANEL, SERUM-

Trilodo FARLE, Section and the section of T3 is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3

Below mentioned	are the guidelines f	or Pregnancy relate	d reference ranges for Tota
Levels in	TOTAL T4	TSH3G	TOTAL T3
Pregnancy	(µg/dL)	(µIU/mL)	(ng/dL)
First Trimester	6.6 - 12.4	0.1 - 2.5	81 - 190
2nd Trimester	6.6 - 15.5	0.2 - 3.0	100 - 260
3rd Trimester	6.6 - 15.5	0.3 - 3.0	100 - 260
Below mentioned	are the guidelines f	or age related refer	ence ranges for T3 and T4.
Т3		T4	
(na/dL)		(ug/dL)	

(ng/dL) New Born: 75 - 260 (µg/dL) 1-3 day: 8.2 - 19.9 1 Week: 6.0 - 15.9

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group.

Kindly note: Method specific reference ranges are appearing on the report under biological reference range.













CLIENT'S NAME AND ADDRESS : I FEI AVATT I AHANE

SRL Ltd 74, PASHCHIMI MARG, VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

PATIENT NAME : LEELAVATI LAH	ANE	PATIENT ID : LEELF09038563
ACCESSION NO : 0063VD001703	AGE : 37 Years SEX : Female	
DRAWN : 08/04/2022 09:13	RECEIVED : 08/04/2022 09:15	REPORTED : 09/04/2022 10:18
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
Test Report Status Final	Results	Biological Reference Interval Units

Reference

1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.

Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.

3. Behrman R.E. Kilegman R.M., Jenson H. B. Nelson Text Book of Pediatrics, 17th Edition STOOL: OVA & PARASITE-

Acute infective diarrhoea and gastroenteritis (diarrhoea with vomiting) are major causes of ill health and premature death in developing countries. Loss of water and electrolytes from the body can lead to severe dehydration which if untreated, can be rapidly fatal in young children, especially that are malnourished, hypoglycaemic, and generally in poor health.

Laboratory diagnosis of parasitic infection is mainly based on microscopic examination and the gross examination of the stool specimen. Depending on the nature of the parasite, the microscopic observations include the identification of cysts, ova, trophozoites, larvae or portions of adult structure. The two classes of parasites that cause human infection are the Protozoa and Helminths. The protozoan infections include amoebiasis mainly caused by Entamoeba histolytica and giardiasis caused by Giardia lamblia. The common helminthic parasites are Trichuris trichiura, Ascaris lumbricoides, Strongyloides stercoralis, Taenia sp. etc

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

MEDICAL THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

FITNESS STATUS-

Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history; as well as the comprehensiveness of the diagnostic panel which has been requested for . These are then further correlated with details of the job under consideration to eventually fit the right man to the right job. Basis the above, SRL classifies a candidate's Fitness Status into one of the following categories:

• Fit (As per requested panel of tests) – SRL Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.

specific test panel requested for.
Fit (with medical advice) (As per requested panel of tests) - This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician's consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.
Fitness on Hold (Temporary Unfit) (As per requested panel of tests) - Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.

elevated blood sugars, etc. • Unfit (As per requested panel of tests) - An unfit report by SRL Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color

blindness in color related jobs













CLIENT CODE : C000138377 CLIENT'S NAME AND ADDRESS :

LEELAVATI LAHANE

SRL Ltd 74,PASHCHIMI MARG,VASANT VIHAR NEW DELHI, 110057 NEW DELHI, INDIA Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.palammarg@srl.in

ACCESSION NO : 0063VD001703	AGE: 37 Years SEX : Female	
DRAWN : 08/04/2022 09:13	RECEIVED : 08/04/2022 09:15	REPORTED : 09/04/2022 10:18
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Units

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

* ULTRASOUND ABDOMEN

ULTRASOUND ABDOMEN REPORTS ENCLOSED.

> **End Of Report** Please visit www.srlworld.com for related Test Information for this accession TEST MARKED WITH '*' ARE OUTSIDE THE NABL ACCREDITED SCOPE OF THE LABORATORY.

K. I. Prejapati

Dr. Kamlesh I Prajapati Consultant Pathologist

CONDITIONS OF LABORAT	ORY TESTING & REPORTING
 It is presumed that the test sample belongs to the patient named or identified in the test requisition form. All Tests are performed and reported as per the turnaround time stated in the SRL Directory of services (DOS). SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity. A requested test might not be performed if: a. Specimen received is insufficient or inappropriate specimen quality is unsatisfactory b. Incorrect specimen type c. Request for testing is withdrawn by the ordering doctor or patient d. There is a discrepancy between the label on the specimen container and the name on the test requisition form 	 The results of a laboratory test are dependent on the quality of the sample as well as the assay technology. Result delays could be because of uncontrolled circumstances. e.g. assay run failure. Tests parameters marked by asterisks are excluded from the "scope" of NABL accredited tests. (If laboratory is accredited). Laboratory results should be correlated with clinical information to determine Final diagnosis. Test results are not valid for Medico- legal purposes. In case of queries or unexpected test results please call at SRL customer care (Toll free: 1800-222-000). Post proper investigation repeat analysis may be carried out.
	SRL Limited

Fortis Hospital, Sector 62, Phase VIII,

Mohali 160062



