PID No.
 : MED121024252
 Register On
 : 03/05/2022 10:12 AM

 SID No.
 : 522215958
 Collection On
 : 03/05/2022 11:33 AM

 Age / Sex
 : 44 Year(s) / Male
 Report On
 : 03/05/2022 6:30 PM

Printed On

: 16/05/2022 5:51 PM



Ref. Dr : MediWheel

: OP

Type

Investigation  II A FINATION OCT	Observed Value	<u>Unit</u>	Biological Reference Interval
<b>HAEMATOLOGY</b>			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	15.2	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	45.7	%	42 - 52
RBC Count (EDTA Blood)	5.06	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	90.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	30.1	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.3	g/dL	32 - 36
RDW-CV	13.5	%	11.5 - 16.0
RDW-SD	42.71	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	5000	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	41.7	%	40 - 75
Lymphocytes (Blood)	42.6	%	20 - 45
Eosinophils (Blood)	7.4	%	01 - 06



**VERIFIED BY** 



 PID No.
 : MED121024252
 Register On
 : 03/05/2022 10:12 AM

 SID No.
 : 522215958
 Collection On
 : 03/05/2022 11:33 AM

 Age / Sex
 : 44 Year(s) / Male
 Report On
 : 03/05/2022 6:30 PM

**Printed On** 



Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood)	7.9	%	01 - 10
Basophils (Blood)	0.4	%	00 - 02
INTERPRETATION: Tests done on Automated Five Pa	art cell counter. All	abnormal results are	reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	2.08	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.13	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.37	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.40	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.02	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	267	10 <b>^</b> 3 / μl	150 - 450
MPV (Blood)	7.7	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.21	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate)	7	mm/hr	< 15

: 16/05/2022 5:51 PM



(Citrated Blood)

**VERIFIED BY** 



 PID No.
 : MED121024252
 Register On
 : 03/05/2022 10:12 AM

 SID No.
 : 522215958
 Collection On
 : 03/05/2022 11:33 AM

 Age / Sex
 : 44 Year(s) / Male
 Report On
 : 03/05/2022 6:30 PM

**Printed On** 



Ref. Dr : MediWheel

: OP

Type

Investigation BIOCHEMISTRY	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.68	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.22	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.46	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	18.94	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	16.83	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	19.13	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	64.9	U/L	53 - 128
Total Protein (Serum/Biuret)	7.50	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.83	gm/dl	3.5 - 5.2
Globulin	2.67	gm/dL	2.3 - 3.6

1.81

: 16/05/2022 5:51 PM



(Serum/Derived)

A : G RATIO (Serum/Derived)



1.1 - 2.2

 PID No.
 : MED121024252
 Register On
 : 03/05/2022 10:12 AM

 SID No.
 : 522215958
 Collection On
 : 03/05/2022 11:33 AM

**Printed On** 

Type : OP

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	218.83	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	87.36	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 16/05/2022 5:51 PM

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	53.53	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	147.8	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol	17.5	mg/dL	< 30



(Serum/Calculated)



PID No. : MED121024252 Register On : 03/05/2022 10:12 AM

: 522215958 SID No. Collection On : 03/05/2022 11:33 AM

Age / Sex : 44 Year(s) / Male Report On 03/05/2022 6:30 PM **Type** 

Ref. Dr : MediWheel

: OP **Printed On** : 16/05/2022 5:51 PM

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

Non HDL Cholesterol 165.3 mg/dL

(Serum/Calculated)

Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio 4.1 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated) Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 1.6 Optimal: < 2.5

Mild to moderate risk: 2.5 - 5.0 (TG/HDL)

High Risk: > 5.0(Serum/Calculated)

Optimal: 0.5 - 3.0 LDL/HDL Cholesterol Ratio 2.8 Borderline: 3.1 - 6.0 (Serum/Calculated)

High Risk: > 6.0





**Type** : OP **Printed On** : 16/05/2022 5:51 PM

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i> )	5.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 105.41 mg/dL

(Whole Blood)

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



**VERIFIED BY** 



**Printed On** 

**Age / Sex** : 44 Year(s) / Male **Report On** : 03/05/2022 6:30 PM

Ref. Dr : MediWheel

: OP

**Type** 

**(\*)** MEDALI

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

: 16/05/2022 5:51 PM

### **IMMUNOASSAY**

#### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.22 ng/ml 0.7 - 2.04

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Tyroxine) - Total 9.33 µg/dl 4.2 - 12.0

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.76 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

#### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

#### **Comment:**

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.





 PID No.
 : MED121024252
 Register On
 : 03/05/2022 10:12 AM

 SID No.
 : 522215958
 Collection On
 : 03/05/2022 11:33 AM

 Age / Sex
 : 44 Year(s) / Male
 Report On
 : 03/05/2022 6:30 PM



Type : OP

Printed On : 16/05/2022 5:51 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

# **CLINICAL PATHOLOGY**

# PHYSICAL EXAMINATION (URINE COMPLETE)

Colour Pale Yellow Yellow to Amber

(Urine)

Appearance Clear Clear

(Urine)

Volume(CLU) 20

(Urine)

# CHEMICAL EXAMINATION (URINE

<u>COMPLETE</u>)

pH 6.0 4.5 - 8.0

(Urine)

Specific Gravity 1.025 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)

Nitrite Negative Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)



**VERIFIED BY** 



**APPROVED BY** 

The results pertain to sample tested.

Page 8 of 12

 PID No.
 : MED121024252
 Register On
 : 03/05/2022 10:12 AM

 SID No.
 : 522215958
 Collection On
 : 03/05/2022 11:33 AM

 Age / Sex
 : 44 Year(s) / Male
 Report On
 : 03/05/2022 6:30 PM

MEDALL

Type : OP

Printed On : 16/05/2022 5:51 PM

Ref. Dr : MediWheel

InvestigationObserved<br/>ValueUnit<br/>Reference Interval

Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

(Urine)

MICROSCOPIC EXAMINATION (URINE COMPLETE)

Pus Cells 1-3 /hpf NIL

(Urine)

Epithelial Cells 2-4 /hpf NIL

(Urine)

RBCs NIL /hpf NIL

(Urine)

Others

(Urine)

**INTERPRETATION:** Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.



**VERIFIED BY** 



**Age / Sex** : 44 Year(s) / Male **Report On** : 03/05/2022 6:30 PM

Ref. Dr : MediWheel

PID No.

**Type** 

: OP **Printed On** : 16/05/2022 5:51 PM



<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u>
<u>Value</u> <u>Reference Interval</u>

# **IMMUNOHAEMATOLOGY**

BLOOD GROUPING AND Rh TYPING 'B' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.





PID No. Register On : 03/05/2022 10:12 AM : MED121024252

**Printed On** 

: 522215958 SID No. Collection On : 03/05/2022 11:33 AM

Age / Sex : 44 Year(s) / Male Report On 03/05/2022 6:30 PM **Type** 

: OP

Ref. Dr : MediWheel



Investigation  PLOCULED (ICEDA)	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY  BUN / Creatinine Ratio	9		6.0 - 22.0
Glucose Fasting (FBS)	83.83	mg/dL	Normal: < 100
(Plasma - F/GOD-PAP)			Pre Diabetic: 100 - 125 Diabetic: >= 126

: 16/05/2022 5:51 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	100.23	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	9.8	mg/dL	7.0 - 21
(Serum/ <i>Urease UV / derived</i> )			
Creatinine	1.03	mg/dL	0.9 - 1.3
(Serum/Modified Jaffe)			

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine

Uric Acid 6.05 3.5 - 7.2mg/dL (Serum/Enzymatic)





**APPROVED BY** 

Page 11 of 12

**Printed On** 

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Prostate specific antigen - Total(PSA) (Serum/Manometric method)	0.574	ng/ml	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0

: 16/05/2022 5:51 PM





**APPROVED BY** 

-- End of Report --