Age / Sex : 32 Year(s) / Male

: OP

PID No. : MED111179628 Register On : 422051814 SID No.

: 28/06/2022 8:59 AM

Collection On : 28/06/2022 9:27 AM

: 06/07/2022 11:37 AM

Report On : 28/06/2022 3:55 PM

Printed On

Ref. Dr : MediWheel

Type



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	16.1	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	48.9	%	42 - 52
RBC Count (EDTA Blood)	5.69	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	85.9	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	28.3	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.0	g/dL	32 - 36
RDW-CV (EDTA Blood)	13.4	%	11.5 - 16.0
RDW-SD (EDTA Blood)	40.7	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	7700	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood)	52.5	%	40 - 75
Lymphocytes (EDTA Blood)	35.7	%	20 - 45
Eosinophils (EDTA Blood)	4.4	%	01 - 06



PID No. : MED111179628 Register On : 28/06/2022 8:59 AM : 422051814 Collection On : 28/06/2022 9:27 AM SID No. Age / Sex : 32 Year(s) / Male Report On : 28/06/2022 3:55 PM



Type : OP

Printed On : 06/07/2022 11:37 AM Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (EDTA Blood)	7.0	%	01 - 10
Basophils (Blood)	0.4	%	00 - 02
INTERPRETATION: Tests done on Automated Five	Part cell counter. All	abnormal results are	e reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	4.1	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.8	10^3 / μ1	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.3	10^3 / μ1	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.5	10^3 / μ1	< 1.0
Absolute Basophil count (EDTA Blood)	0.0	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	201	10^3 / µl	150 - 450
MPV (EDTA Blood)	13.2	fL	7.9 - 13.7
PCT (EDTA Blood/Automated Blood cell Counter)	0.265	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	6	mm/hr	< 15



Age / Sex : 32 Year(s) / Male

PID No. : MED111179628 Register On : 28/06/2022 8:59 AM : 422051814 SID No. Collection On : 28/06/2022 9:27 AM

Type : OP

Ref. Dr : MediWheel

Printed On : 06/07/2022 11:37 AM

Report On

: 28/06/2022 3:55 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.41	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.18	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.23	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	38.13	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	79.56	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	45.11	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	86.4	U/L	53 - 128
Total Protein (Serum/Biuret)	7.50	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.67	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.83	gm/dL	2.3 - 3.6
A: GRATIO (Serum/Derived)	1.65		1.1 - 2.2





Age / Sex : 32 Year(s) / Male

 PID No.
 : MED111179628
 Register On
 : 28/06/2022 8:59 AM

 SID No.
 : 422051814
 Collection On
 : 28/06/2022 9:27 AM

Report On

Printed On

28/06/2022 9:27 AM 28/06/2022 3:55 PM (*) MEDALL

Type : OP

Ref. Dr : MediWheel

: 06/07/2022 11:37 AM

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	208.55	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	249.61	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	37.41	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	121.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	49.9	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	171.1	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220





 PID No.
 : MED111179628
 Register On
 : 28/06/2022 8:59 AM

 SID No.
 : 422051814
 Collection On
 : 28/06/2022 9:27 AM

 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 28/06/2022 3:55 PM

Printed On



Type : OP

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u> <u>Value</u> <u>Reference Interval</u>

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 06/07/2022 11:37 AM

Total Cholesterol/HDL Cholesterol Ratio 5.6 Optimal: < 3.3 (Serum/Calculated) Low Risk: 3.4 - 4.4

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Mild to moderate risk: 2.5 - 5.0

High Risk: > 5.0

Triglyceride/HDL Cholesterol Ratio 6.7 Optimal: < 2.5

(TG/HDL)

(Serum/Calculated)

Consultant Pathologist

Reg No: 99049

VERIFIED BY

LDL/HDL Cholesterol Ratio

3.2

Optimal: 0.5 - 3.0

(Serum/Calculated)
Borderline: 3.1 - 6.0
High Risk: > 6.0



Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	6.0	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 06/07/2022 11:37 AM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Printed On

Estimated Average Glucose 125.5 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.





: MediWheel

PID No. Register On : 28/06/2022 8:59 AM : MED111179628

: 422051814 SID No. Collection On : 28/06/2022 9:27 AM Age / Sex : 32 Year(s) / Male

Type : OP

Ref. Dr

Report On : 28/06/2022 3:55 PM

Printed On : 06/07/2022 11:37 AM

<u>Unit</u> **Biological** <u>Investigation</u> <u>Observed</u> <u>Value</u>

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.60 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Tyroxine) - Total 9.15 μg/dl 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is

Metabolically active.

 $\mu IU/mL$ 0.35 - 5.50TSH (Thyroid Stimulating Hormone) 2.05

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



Reference Interval

 PID No.
 : MED111179628
 Register On
 : 28/06/2022 8:59 AM

 SID No.
 : 422051814
 Collection On
 : 28/06/2022 9:27 AM

 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 28/06/2022 3:55 PM

Printed On



Type : OP

Ref. Dr : MediWheel

: 06/07/2022 11:37 AM

InvestigationObservedUnitBiologicalValueReference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION (URINE COMPLETE)

Colour Yellow Yellow Yellow to Amber

(Urine)

Appearance clear Clear

(Urine)

Volume(CLU) 20

(Urine)

CHEMICAL EXAMINATION (URINE

<u>COMPLETE</u>)

pH 5.5 4.5 - 8.0

(Urine)

Specific Gravity 1.016 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)
Nitrite Negative

Nitrite Negative Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)





 PID No.
 : MED111179628
 Register On
 : 28/06/2022 8:59 AM

 SID No.
 : 422051814
 Collection On
 : 28/06/2022 9:27 AM

 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 28/06/2022 3:55 PM

Printed On



Type : OP

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u>
<u>Value</u> <u>Reference Interval</u>

: 06/07/2022 11:37 AM

Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

(Urine)

MICROSCOPIC EXAMINATION (URINE COMPLETE)

Pus Cells 0-1 /hpf NIL

(Urine)

Epithelial Cells **0-1** /hpf NIL

(Urine)

RBCs NIL /HPF NIL

(Urine)

Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

Casts NIL /hpf NIL

(Urine)

Crystals NIL /hpf NIL

(Urine)



APPROVED BY

Consultant Pathologist

Reg No: 99049

VERIFIED BY

PID No. **Register On** : 28/06/2022 8:59 AM : MED111179628 : 422051814 SID No.

Age / Sex : 32 Year(s) / Male Report On

Printed On

Type : OP

Ref. Dr : MediWheel

Collection On : 28/06/2022 9:27 AM

: 28/06/2022 3:55 PM

: 06/07/2022 11:37 AM



Investigation <u>Unit</u> <u>Observed</u> **Biological** <u>Value</u> Reference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'O' 'Positive'

(EDTA Blood/Agglutination)



VERIFIED BY



PID No. Register On : MED111179628 : 422051814 SID No.

Age / Sex : 32 Year(s) / Male Report On 28/06/2022 3:55 PM

Printed On

Type : OP

Ref. Dr : MediWheel : 28/06/2022 8:59 AM

Collection On : 28/06/2022 9:27 AM

: 06/07/2022 11:37 AM



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	12.89		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	89.18	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	105.22	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	9.8	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.76	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists, N-acetylcyteine , chemotherapeutic agent such as flucytosine

Uric Acid 6.61 3.5 - 7.2mg/dL

(Serum/Enzymatic)





APPROVED BY

-- End of Report --