

| Name | Mr.UDAYA KUMAR R V | ID | MED111193309 |
|-----------------|--------------------|------------|--------------|
| Age & Gender | 32/MALE | Visit Date | 09/07/2022 |
| Ref Doctor Name | MediWheel | | |

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size and shows diffuse fatty changes. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER shows normal shape and has clear contents. Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

No demonstrable Para -aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

| | Bipolar length (cms) | Parenchymal thickness (cms) |
|--------------|----------------------|-----------------------------|
| Right Kidney | 10.7 | 1.3 |
| Left Kidney | 10.9 | 1.3 |

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern. It measures 2.9 x 2.4 x 2.9cms (Vol: 11cc).

No evidence of ascites / pleural effusion.

IMPRESSION:

- > FATTY LIVER.
- > NO OTHER SIGNIFICANT ABNORMALITY DETECTED.

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



| Name | Mr.UDAYA KUMAR R V | ID | MED111193309 |
|-----------------|--------------------|------------|--------------|
| Age & Gender | 32/MALE | Visit Date | 09/07/2022 |
| Ref Doctor Name | MediWheel | | |

DR. APARNA CONSULTANT RADIOLOGIST

A/an

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11. Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

PID No. : MED111193309 **SID No.** : 422053993

Age / Sex : 32 Year(s) / Male

- 00

Type

: OP

Register On : 09/07/2022 10:12 AM

Collection On : 09/07/2022 10:43 AM

Report On : 10/07/2022 9:57 AM

Printed On : 15/07/2022 11:34 AM

Ref. Dr : MediWheel

| (7) | |
|------|----|
| MEDA | LL |

| Investigation | Observed Value | <u>Unit</u> | <u>Biological</u> <u>Reference Interval</u> |
|---|-------------------|-------------|--|
| HAEMATOLOGY | | | |
| Complete Blood Count With - ESR | | | |
| Haemoglobin (EDTA Blood/Spectrophotometry) | 15.64 | g/dL | 13.5 - 18.0 |
| Packed Cell Volume(PCV)/Haematocrit (EDTA Blood) | 48.3 | % | 42 - 52 |
| RBC Count (EDTA Blood) | 5.74 | mill/cu.mm | 4.7 - 6.0 |
| Mean Corpuscular Volume(MCV) (EDTA Blood) | 84.2 | fL | 78 - 100 |
| Mean Corpuscular Haemoglobin(MCH) (EDTA Blood) | 27.2 | pg | 27 - 32 |
| Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood) | 32.4 | g/dL | 32 - 36 |
| RDW-CV (EDTA Blood) | 14.3 | % | 11.5 - 16.0 |
| RDW-SD (EDTA Blood) | 42.14 | fL | 39 - 46 |
| Total Leukocyte Count (TC) (EDTA Blood) | 5800 | cells/cu.mm | 4000 - 11000 |
| Neutrophils (EDTA Blood) | 56.73 | % | 40 - 75 |
| Lymphocytes (EDTA Blood) | 35.73 | % | 20 - 45 |
| Eosinophils (EDTA Blood) | 2.30 | % | 01 - 06 |



VERIFIED BY



 PID No.
 : MED111193309
 Register On
 : 09/07/2022 10:12 AM

 SID No.
 : 422053993
 Collection On
 : 09/07/2022 10:43 AM

 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 10/07/2022 9:57 AM

Printed On



Ref. Dr : MediWheel

: OP

Type

| <u>Investigation</u> | Observed Value | <u>Unit</u> | <u>Biological</u> Reference Interval |
|---|-----------------------|----------------------|---|
| Monocytes (EDTA Blood) | 5.12 | % | 01 - 10 |
| Basophils (Blood) | 0.12 | % | 00 - 02 |
| INTERPRETATION: Tests done on Automated Five Pa | art cell counter. All | abnormal results are | reviewed and confirmed microscopically. |
| Absolute Neutrophil count (EDTA Blood) | 3.29 | 10^3 / μl | 1.5 - 6.6 |
| Absolute Lymphocyte Count (EDTA Blood) | 2.07 | 10^3 / μl | 1.5 - 3.5 |
| Absolute Eosinophil Count (AEC) (EDTA Blood) | 0.13 | 10^3 / μl | 0.04 - 0.44 |
| Absolute Monocyte Count (EDTA Blood) | 0.30 | 10^3 / μl | < 1.0 |
| Absolute Basophil count (EDTA Blood) | 0.01 | 10^3 / μl | < 0.2 |
| Platelet Count (EDTA Blood) | 236.8 | 10^3 / μl | 150 - 450 |
| MPV (EDTA Blood) | 7.34 | fL | 7.9 - 13.7 |
| PCT (EDTA Blood/Automated Blood cell Counter) | 0.17 | % | 0.18 - 0.28 |
| ESR (Erythrocyte Sedimentation Rate) (Citrated Blood) | 4 | mm/hr | < 15 |

: 15/07/2022 11:34 AM



VERIFIED BY



 PID No.
 : MED111193309
 Register On
 : 09/07/2022 10:12 AM

 SID No.
 : 422053993
 Collection On
 : 09/07/2022 10:43 AM

 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 10/07/2022 9:57 AM

Printed On

MEDALL

Type : OP

| Ref. Dr | : MediWheel | | | |
|---------|-------------|--|--|--|
| | | | | |

: 15/07/2022 11:34 AM

| Investigation | Observed Value | <u>Unit</u> | <u>Biological</u> <u>Reference Interval</u> |
|---|-------------------|-------------|--|
| BIOCHEMISTRY | | | |
| Liver Function Test | | | |
| Bilirubin(Total) (Serum/DCA with ATCS) | 1.44 | mg/dL | 0.1 - 1.2 |
| Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid) | 0.39 | mg/dL | 0.0 - 0.3 |
| Bilirubin(Indirect) (Serum/Derived) | 1.05 | mg/dL | 0.1 - 1.0 |
| SGOT/AST (Aspartate Aminotransferase) (Serum/ <i>Modified IFCC</i>) | 26.85 | U/L | 5 - 40 |
| SGPT/ALT (Alanine Aminotransferase) (Serum/ <i>Modified IFCC</i>) | 31.62 | U/L | 5 - 41 |
| GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic) | 21.63 | U/L | < 55 |
| Alkaline Phosphatase (SAP) (Serum/ <i>Modified IFCC</i>) | 79.4 | U/L | 53 - 128 |
| Total Protein (Serum/Biuret) | 7.65 | gm/dl | 6.0 - 8.0 |
| Albumin (Serum/Bromocresol green) | 4.87 | gm/dl | 3.5 - 5.2 |
| Globulin (Serum/Derived) | 2.78 | gm/dL | 2.3 - 3.6 |
| A : G RATIO (Serum/Derived) | 1.75 | | 1.1 - 2.2 |



VERIFIED BY



PID No. : MED111193309 Register On : 09/07/2022 10:12 AM : 422053993 SID No. Collection On : 09/07/2022 10:43 AM Age / Sex : 32 Year(s) / Male Report On : 10/07/2022 9:57 AM

Printed On



Type : OP

Ref. Dr : MediWheel

: 15/07/2022 11:34 AM

| Investigation | <u>Observed</u> <u>Value</u> | <u>Unit</u> | <u>Biological</u> <u>Reference Interval</u> |
|---|---------------------------------|-------------|---|
| <u>Lipid Profile</u> | | | |
| Cholesterol Total (Serum/CHOD-PAP with ATCS) | 154.41 | mg/dL | Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240 |
| Triglycerides (Serum/GPO-PAP with ATCS) | 134.81 | mg/dL | Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500 |

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_ circulating level of triglycerides during most part of the day.

| r · · · · · · · · · · · · · · · · · · · | | | |
|--|-------|-------|--|
| HDL Cholesterol (Serum/Immunoinhibition) | 28.78 | mg/dL | Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40 |
| LDL Cholesterol (Serum/Calculated) | 98.6 | mg/dL | Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190 |
| VLDL Cholesterol (Serum/Calculated) | 27 | mg/dL | < 30 |
| Non HDL Cholesterol (Serum/Calculated) | 125.6 | mg/dL | Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220 |





PID No. : MED111193309 Register On : 09/07/2022 10:12 AM : 422053993 SID No. Collection On : 09/07/2022 10:43 AM Age / Sex : 32 Year(s) / Male Report On : 10/07/2022 9:57 AM

Printed On



Type : OP

Ref. Dr : MediWheel

Investigation Observed **Unit Biological** <u>Value</u> Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 15/07/2022 11:34 AM

Total Cholesterol/HDL Cholesterol Ratio 5.4 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated) Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0

High Risk: > 11.0 Triglyceride/HDL Cholesterol Ratio Optimal: < 2.54.7

Mild to moderate risk: 2.5 - 5.0 (TG/HDL) High Risk: > 5.0(Serum/Calculated)

LDL/HDL Cholesterol Ratio 3.4 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated) High Risk: > 6.0







 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 10/07/2022 9:57 AM

 Type
 : OP
 Printed On
 : 15/07/2022 11:34 AM

Ref. Dr : MediWheel



| Investigation | Observed Value | <u>Unit</u> | <u>Biological</u> <u>Reference Interval</u> |
|----------------------------------|-------------------|-------------|---|
| Glycosylated Haemoglobin (HbA1c) | | | |
| HbA1C (Whole Blood/HPLC) | 5.6 | % | Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5 |

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 114.02 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



 PID No.
 : MED111193309
 Register On
 : 09/07/2022 10:12 AM

 SID No.
 : 422053993
 Collection On
 : 09/07/2022 10:43 AM

Report On

MEDALI

Type : OP

Ref. Dr

: MediWheel

Age / Sex : 32 Year(s) / Male

Printed On : 15/07/2022 11:34 AM

: 10/07/2022 9:57 AM

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.29 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 8.60 µg/dl 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.27 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



PID No. : MED111193309 : 09/07/2022 10:12 AM Register On : 422053993 SID No. Collection On : 09/07/2022 10:43 AM Age / Sex : 32 Year(s) / Male Report On : 10/07/2022 9:57 AM

Printed On



Type : OP Ref. Dr : MediWheel

: 15/07/2022 11:34 AM

| <u>Investigation</u> | <u>Observed</u> <u>l</u> | <u>Unit</u> | <u>Biological</u> |
|----------------------|--------------------------|-------------|--------------------|
| - | Value | | Reference Interval |

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION (URINE **COMPLETE**)

| Colour | Pale yellow | Yellow to Amber |
|---------|-------------|-----------------|
| (Urine) | | |

(Urine)

Clear Clear Appearance

(Urine)

25 Volume(CLU)

(Urine)

CHEMICAL EXAMINATION (URINE

COMPLETE)

| pH 5.5 4.5 - | 8.0 |
|--------------|-----|
|--------------|-----|

(Urine)

1.012 1.002 - 1.035 Specific Gravity

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)

Negative Negative Nitrite

(Urine)

Bilirubin Negative Negative

(Urine)

(Urine)

Negative Protein Negative

Reg No:KMC 89655

VERIFIED BY



 PID No.
 : MED111193309
 Register On
 : 09/07/2022 10:12 AM

 SID No.
 : 422053993
 Collection On
 : 09/07/2022 10:43 AM

 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 10/07/2022 9:57 AM

Printed On



Type : OP

Ref. Dr : MediWheel

| <u>Investigation</u> | Observed Unit | <u>Biological</u> |
|----------------------|---------------|--------------------|
| | <u>Value</u> | Reference Interval |

: 15/07/2022 11:34 AM

Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

(Urine)

MICROSCOPIC EXAMINATION (URINE COMPLETE)

Pus Cells 0-1 /hpf NIL

(Urine)

Epithelial Cells **0-1** /hpf NIL

(Urine)

RBCs NIL /HPF NIL

(Urine)

Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

Casts NIL /hpf NIL

(Urine)

Crystals NIL /hpf NIL

(Urine)



VERIFIED BY



 PID No.
 : MED111193309
 Register On
 : 09/07/2022 10:12 AM

 SID No.
 : 422053993
 Collection On
 : 09/07/2022 10:43 AM

 Age / Sex
 : 32 Year(s) / Male
 Report On
 : 10/07/2022 9:57 AM

Printed On



Type : OP

Ref. Dr : MediWheel

| <u>Investigation</u> | <u>Observed</u> <u>U</u> i | <u>Biological</u> |
|----------------------|----------------------------|--------------------|
| | Value | Reference Interval |

: 15/07/2022 11:34 AM

<u>PHYSICAL EXAMINATION(STOOL</u> <u>COMPLETE)</u>

Mucus Absent Absent

(Stool)

Consistency Semi Solid Semi Solid to Solid

(Stool)

Colour Brownish Brown

(Stool)

Blood Absent Absent

(Stool)

<u>MICROSCOPIC EXAMINATION(STOOL</u> <u>COMPLETE)</u>

Ova NIL NIL

(Stool)

Cysts NIL NIL

(Stool)

Trophozoites NIL NIL

(Stool)

RBCs NIL /hpf Nil

(Stool)

Pus Cells 1-2 /hpf NIL

(Stool)

Others

(Stool)

<u>CHEMICAL EXAMINATION(STOOL</u> <u>ROUTINE)</u>



VERIFIED BY



PID No. : MED111193309

: 422053993 SID No.

Age / Sex : 32 Year(s) / Male

Type : OP

Ref. Dr : MediWheel Register On : 09/07/2022 10:12 AM

Collection On : 09/07/2022 10:43 AM

: 10/07/2022 9:57 AM

: 15/07/2022 11:34 AM **Printed On**



Negative

| <u>Investigation</u> | <u>Observed</u> <u>Unit</u> <u>Value</u> | <u>Biological</u> <u>Reference Interval</u> |
|----------------------|---|--|
| Reaction | Alkaline | Alkaline |
| (Stool) | | |

Negative

Report On

Reducing Substances (Stool/Benedict's)



VERIFIED BY



PID No. : MED111193309 Register On : 422053993

SID No. : 10/07/2022 9:57 AM

Age / Sex : 32 Year(s) / Male Report On **Type** : OP

Ref. Dr : MediWheel : 09/07/2022 10:12 AM

Collection On : 09/07/2022 10:43 AM

Printed On : 15/07/2022 11:34 AM

Investigation <u>Unit</u> **Biological** <u>Observed</u> Reference Interval Value

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING

(EDTA Blood/Agglutination)

Remark: Kindly confirm Rh status by Du Testing

'O' 'Negative'



VERIFIED BY



PID No. : MED111193309

- 400050000

: 422053993

Age / Sex : 32 Year(s) / Male

Type : OP

SID No.

Ref. Dr : MediWheel

Register On : 09/07/2022 10:12 AM

Collection On : 09/07/2022 10:43 AM

Report On : 10/07/2022 9:57 AM

Printed On : 15/07/2022 11:34 AM



| Investigation | <u>Observed</u> <u>Value</u> | <u>Unit</u> | <u>Biological</u> <u>Reference Interval</u> |
|---|---------------------------------|-------------|--|
| BIOCHEMISTRY | | | |
| BUN / Creatinine Ratio | 10 | | 6.0 - 22.0 |
| Glucose Fasting (FBS) (Plasma - F/GOD-PAP) | 78.73 | mg/dL | Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126 |

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)

(Urine - F/GOD - POD)

Negative

Glucose Postprandial (PPBS) 67.43 mg/dL 70 - 140

(Plasma - PP/GOD-PAP)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN) 5.6 mg/dL 7.0 - 21 (Serum/Urease UV / derived)

Creatinine 0.71 mg/dL 0.9 - 1.3 (Serum/Modified Jaffe)

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine

Uric Acid 5.23 mg/dL 3.5 - 7.2

(Serum/Enzymatic)



VERIFIED BY



APPROVED BY

-- End of Report --