

Dr. GOYAL'S

PATH LAB & MAGING CENTER

PATH LAB & MAGING CENTER

B-51, Genesh Nagar-B

New Sangarer Koad, JAJPUR



Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs

Company :- MediWheel

Tele: 0141-2293346, 4049787, 9887049787

Patient ID :-122127686

Ref. By Dr.- BOB

Lab/Hosp :-



Sample Type :-

Sample Collected Time

Final Authentication: 13/03/2022 14:30:03

BOB PACKAGEFEMALE BELOW 40

X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression :- Normal Study

(Please correlate clinically and with relevant further investigations)

*** End of Report ***

ANITASHARMA

Page No: 1 of 1

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996 Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495 Dr. Tej Prakash Gupta DMRD (RADIO DIAGNOSIS) RMC No. 24436

Dr. Hitesh Kumar Sharma M.B.B.S.,D.M.R.D. RMC Reg No. 27380

Transcript by.

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





:- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Sample Type :- EDTA

Sex / Age :- Female

29 Yrs

Company:- MediWheel

Patient ID: -122127686

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 13/03/2022 13:16:01 **HAEMATOLOGY**

Test Name Value

Unit Biological Ref Interval

BOB PACKAGEFEMALE BELOW 40

GLYCOSYLATED HEMOGLOBIN (HbA1C)

5.5

Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0

Final Authentication: 13/03/2022 14:45:17

Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb.High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1cmeasurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method. Ref by ADA 2020

MEAN PLASMA GLUCOSE

Method:- Calculated Parameter

111

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100-125 mg/dL Diabetic 126 mg/dL or Higher

BANWARI **Technologist**

Page No: 1 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019
Tele: 0141-2293346, 4049787, 9887049787
Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 13/03/2022 13:02:44

NAME: - Mrs. GARIMA SAINI
Sex / Age: - Female 29 Yrs
Company: - MediWheel

Patient ID :-122127686 Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type :- EDTA

Sample Collected Time 13/03/2022 13:16:01

Final Authentication: 13/03/2022 14:45:17

H	A	F	M	IA	TC	M	0	CI	J
	$\overline{}$		14	H		,,			r

Test Name	Value	Unit	Biological Ref Interval
HAEMOGARAM			
HAEMOGLOBIN (Hb)	11.6 L	g/dL	12.0 - 15.0
TOTAL LEUCOCYTE COUNT	6.28	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	55.6	%	40.0 - 80.0
LYMPHOCYTE	39.8	%	20.0 - 40.0
EOSINOPHIL	1.5	%	1.0 - 6.0
MONOCYTE	2.9	%	2.0 - 10.0
BASOPHIL	0.2	%	0.0 - 2.0
NEUT#	3.42	10^3/uL	1.50 - 7.00
LYMPH#	2.58	10^3/uL	1.00 - 3.70
EO#	0.09	10^3/uL	0.00 - 0.40
MONO#	0.18	10^3/uL	0.00 - 0.70
BASO#	0.01	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	4.18	x10^6/uL	3.80 - 4.80
HEMATOCRIT (HCT)	34.20 L	%	36.00 - 46.00
MEAN CORP VOLUME (MCV)	81.7 ∟	fL	83.0 - 101.0
MEAN CORP HB (MCH)	27.7	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	33.9	g/dL	31.5 - 34.5
PLATELET COUNT	323	x10^3/uL	150 - 410
RDW-CV	14.0	%	11.6 - 14.0
MENTZER INDEX	19.55		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

BANWARI Technologist

Page No: 2 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

NAME :- Mrs. GARIMA SAINI

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

29 Yrs



Sex / Age :- Female Company:- MediWheel Sample Type :- EDTA

Sample Collected Time 13/03/2022 13:16:01

Final Authentication: 13/03/2022 14:45:17

00 - 20

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval

Lab/Hosp:-

Ref. By Dr:- BOB

mm/hr.

Erythrocyte Sedimentation Rate (ESR) (ESR) Methodology: Measurment of ESR by cells aggregation.

Instrument Name: Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

: ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia for Bornetthe 1882 Clarent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance, and MCH, MCV, MCHC, MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L. Japan

BANWARI Technologist

Page No: 3 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





MC - 2300

Date

:- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs

Company:- MediWheel

Patient ID: -122127686

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 13/03/2022 13:16:01

Final Authentication: 13/03/2022 14:08:50

BIOCHEMISTRY

Test Name Value

Unit

Biological Ref Interval

LIPID PROFILE

TRIGLYCERIDES

Method:- GPO-PAP

TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method

Sample Type :- PLAIN/SERUM

267.06 H

mg/dl

mg/dl

Desirable <200

428.81 H

Borderline 200-239

High> 240

Normal

<150 Borderline high 150-199

High

200-499

Very high

>500

Rechecked, kindly correlate clinically.

Comment-

- (1) Serum sample is highly lipemic .
- (2) In case of serum Triglyceride value > 400 mg/dl, VLDL can not be calculated

by Formula TG/5 as it leads to errors in calculated parameter. In such cases VLDL is cancelled.

Kindly correlate clinically.

MUKESHSINGH

Page No: 4 of 14




Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs Company :- MediWheel

Sample Type :- PLAIN/SERUM

diWheel

Sample Collected Time 13/03/2022 13:16:01

Time 13/03/2022 13:16:01 Final Authentication: 13/03/2022 14:08:50

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	44.84	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	174.00 H	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190

Patient ID: -122127686

Ref. By Dr:- BOB

Lab/Hosp :-

TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDLCHOLESTERO InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROLInstrumentName: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.

TOTAL LIPID AND VLDL ARE CALCULATED

MUKESHSINGH

Page No: 5 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Sample Type :- PLAIN/SERUM

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date

:- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Company :- MediWheel

Sex / Age :- Female 29 Yrs

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 13/03/2022 13:16:01

Patient ID: -122127686

Final Authentication: 13/03/2022 14:08:50

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.53	mg/dl	Up to - 1.0 Cord blood <2 mg/dL Premature < 6 days <16mg/dL Full-term < 6 days= 12 mg/dL 1month - <12 months <2 mg/dL 1-19 years <1.5 mg/dL Adult - Up to - 1.2 Ref-(ACCP 2020)
SGOT Method:- IFCC	26.9	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	34.0 H	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	82.80	IU/L	30.00 - 120.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.81	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.51	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	3.30	gm/dl	2.20 - 3.50
A/G RATIO	1.37		1.30 - 2.50

MUKESHSINGH

Page No: 6 of 14



Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Sample Type :- PLAIN/SERUM

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs

Company:- MediWheel

Patient ID :-122127686

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 13/03/2022 13:16:01

Final Authentication: 13/03/2022 14:08:50

BIOCHEMISTRY

Value	Unit	Biological Ref Interval
0.26	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL
0.27	mg/dl	0.30-0.70
59.50 H	U/L	7.00 - 32.00
	0.26	0.26 mg/dL 0.27 mg/dl

Total BilirubinMethodology:Colorimetric method InstrumentName:Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName:Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain. fiver, gastric mucosa, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

Alkaline Phosphatase Methodology:AMP Buffer InstrumentName:Randox Rx Imola Interpretation:Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology:Biuret Reagent InstrumentName:Randox Rx Imola Interpretation: Measurements obtained by this method are used in the

TOTAL PROTEIN Methodology: Biuret Reagent InstrumentName: Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName: Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

MUKESHSINGH

Page No: 7 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date

:- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID: -122127686

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 13/03/2022 13:16:01 **IMMUNOASSAY**

Final Authentication: 13/03/2022 14:30:28

Test Name Value Biological Ref Interval

TOTAL THYROID PROFILE

SERUM TSH

Method:- Enhanced Chemiluminescence Immunoassay

1.880

μIU/mL

Unit

0.465 - 4.680

ANANDSHARMA Technologist

Page No: 8 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

29 Yrs

Date :- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Company :- MediWheel

Sex / Age :- Female

Sample Type :- PLAIN/SERUM

Patient ID :-122127686

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Collected Time 13/03/2022 13:16:01

Final Authentication: 13/03/2022 14:30:28



Value	Unit	Biological Ref Interval	
1.240	ng/ml	0.970 - 1.690	
6.760	ug/dl	5.500 - 11.000	
	1.240	1.240 ng/ml	

InstrumentName: VITROS ECI Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

InstrumentName: VITROS ECI Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4.Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

InstrumentName: VITROS ECI Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid
20	Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

ANANDSHARMA **Technologist**

Page No: 9 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Sex / Age :- Female 29 Yrs

Company :- MediWheel

Sample Type :- URINE





Patient ID :-122127686 Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 13/03/2022 13:16:01

Final Authentication: 13/03/2022 14:05:32

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	3-5	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		

POOJABOHRA Technologist

Page No: 10 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

:- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs

Sample Type :- URINE

Company :- MediWheel

Patient ID :-122127686

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 13/03/2022 14:05:32

Sample Collected Time 13/03/2022 13:16:01 **CLINICAL PATHOLOGY**

Test Name	Value Unit	Biological Ref Interval
PHYSICAL EXAMINATION COLOUR APPEARANCE CHEMICAL EXAMINATION	PALE YELLOW Clear	PALE YELLOW Clear
REACTION(PH) SPECIFIC GRAVITY PROTEIN SUGAR BILIRUBIN UROBILINOGEN KETONES NITRITE	5.5 1.025 NIL NIL NEGATIVE NORMAL NEGATIVE NEGATIVE	5.0 - 7.5 1.010 - 1.030 NIL NIL NEGATIVE NORMAL NEGATIVE NEGATIVE

POOJABOHRA Technologist

Page No: 11 of 14



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date :- 13/03/2022 13:02:44

NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs

Company :- MediWheel

Patient ID :-122127686

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Type: KOx/Na FLUORIDE-F, KOx/Na Sabbor IDE PRED TANN SER 12022 16:01:27

Final Authentication: 13/03/2022 16:50:08

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interva
FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP	103.5	mg/dl	75.0 - 115.0
Impaired glucose tolerance (IGT)	1	11 - 125 mg/dL	
Diabetes Mellitus (DM)	>	126 mg/dL	

Instrument Name: Randox Rx Imola **Interpretation:** Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

BLOOD SUGAR PP (Plasma)

Method:- GOD PAP

114.8 mg/dl

70.0 - 140.0

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

SERUM CREATININE Method:- Colorimetric Method	0.84	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20
SERUM URIC ACID Method:- Enzymatic colorimetric	5.43	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7

MUKESHSINGH

Page No: 12 of 14



Dr. Chandrika Gupta MBBS.MD (Path) RMC NO. 21021/008037 DR.TANURUNGTA

Dr. Goyal's Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website : www.drgoyalspathlab.com \mid E-mail : drgoyalpiyush@gmail.com

:- 13/03/2022 13:02:44 NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs

Company :- MediWheel

Patient ID: -122127686

Ref. By Dr:- BOB

Lab/Hosp:-



Sample Type :- EDTA, PLAIN/SERUM, URINE, SERINGE-POLICED Time 13/03/2022 16:01:08

Final Authentication: 13/03/2022 16:18:15

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BLOOD GROUP ABO	"O"POSIT	IVE	
BLOOD GROUP ABO Methodology: Haemagg	glutination reaction Kit	Name: Monoclonal aggl	utinating antibodies (Span clone).
URINE SUGAR (FASTING) Collected Sample Received	Nil		Nil
URINE SUGAR PP Collected Sample Received	Nil		Nil
BLOOD UREA NITROGEN (BUN)	12.0	mg/dl	0.0 - 23.0

*** End of Report ***

BANWARI, MUKESHSINGH, POOJABOHRA **Technologist**

Page No: 14 of 14

Dr. Piyush Goyal

M.B.B.S., D.M.R.D.

RMC Reg No. 017996

MBBS, MD RMC No. 32495

Dr. Tej Prakash Gupta DMRD (RADIO DIAGNOSIS) RMC No. 24436

Dr. Hitesh Kumar Sharma M.B.B.S., D.M.R.D. **RMC Reg No. 27380**

Transcript by.

Dr. Piyush Goyal (D.M.R.D.)

Dr. Chandrika Gupta DR.TANURUNGTA

Dr. Goyal Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 13/03/2022 13:02:44 NAME :- Mrs. GARIMA SAINI

Sex / Age :- Female 29 Yrs Company :- MediWheel

Date

Patient ID :-122127686 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 13/03/2022 15:54:39

BOB PACKAGEFEMALE BELOW 40

ULTRA SOUND SCAN OF ABDOMEN

Liver is of normal size. Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is contracted (Post prandial). Common bile duct is not dilated.

Pancreas is obscured by gases.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Urinary Bladder: is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Uterus is anteverted and normal in size and measures 71x37 mm. Myometrium shows normal echo - pattern. No focal space occupying lesion is seen. Endometrial echo is normal. Endometrial thickness is 12 mm.

Both ovaries are visualised and are normal. No adnexal mass is seen.

No enlarged nodes are visualised. No retro-peritoneal lesion is identified. No significant free fluid is seen in pouch of douglas.

IMPRESSION:

Normal Study.

Needs clinical correlation & further evaluation

Page No: 1 of 1

*** End of Report ***

KOMAL

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Tej Prakash Gupta DMRD (RADIO DIAGNOSIS) RMC No. 24436

Dr. Hitesh Kumar Sharma M.B.B.S., D.M.R.D. **RMC Reg No. 27380**

Transcript by.

DR.GOYALS PATH & IMAGING CENTER





