

Dr. II C

Dr. U. C. GUPTA MBBS, MD (Physician) RMC No. 291





(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023 9 +91 141 4824885 maxcarediagnostics1@gmail.com



General Physical Examination

	Date of Examination: 08/11/22
	Name: DINESH KUMAR SAINE Age: 32 DOB: 01.02.1990 Sex: Male
	Referred By: Rank of Barada
	Photo ID: ADHAR CARO ID #: 1039
	Ht: 167 (cm) Wt: 58 (Kg)
	Chest (Expiration): 28 (cm) Abdomen Circumference: 19 (cm)
	Blood Pressure: 120 80 mm Hg PR: 73 min RR: 17 min Temp: Afelosico
	BMI 20.8.
	With Glass RIE 7 6/6, N/6, NCB Eye Examination: 46) 6/6, N/6, NCB
	Eye Examination: 46)6/6, N/6, NCB
	Other:
	On examination he/she appears physically and mentally fit: Yes/No
	Signature Of Examine: Name of Examinee: DIHESH KUMAR SAINS
5	Signature Medical Examiner: Dr. U. C. Gruph Name Medical Examiner Dr. U. C. Gruph
	in the second se
	Dr. U. C. GUPTA
	MBBS, MD (Physician) RMC No. 291



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

9 +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age:- 32 Yrs 9 Mon 7 Days

Sex :- Male



Patient ID: -12222381

Date :- 08/11/2022

09:07:14

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
Test Name Value Unit FULL BODY HEALTH CHECKUP BELOW 40 MALE HAEMOGARAM HAEMOGLOBIN (Hb) 14.5 g/dL			
	IVIALL		
	145	a/dI	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	5.20	/cumm	4.00 - 10.00
	3.20	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT	50 0	0.4	40.0 00.0
NEUTROPHIL	52.0	%	40.0 - 80.0
LYMPHOCYTE	40.0	%	20.0 - 40.0
EOSINOPHIL	3.0	%	1.0 - 6.0
MONOCYTE	5.0	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	5.41	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	46.60	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	86.0	n.	83.0 - 101.0
MEAN CORP HB (MCH)	26.7 L	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	31.0 L	g/dL	31.5 - 34.5
PLATELET COUNT	220	x10^3/uL	150 - 410
RDW-CV	13.4	%	11.6 - 14.0
MENTZER INDEX A complete blood picture (CBP) is a kind of blood test the	15.90 H	ss a person's overall heal	0.00 - 0.00 th and diagnose a wide range of health

A complete blood picture (CBP) is a kind of blood test that is done to assess a person's overall health and diagnose a wide range of health disorders like leukemia, anemia and other infections.

A complete blood count (CBC) is a complete blood test that diagnose many components and features of a persons blood which includes: -

(CBC): Methodology: TLC,TRBC,PCV,PLT Impedance method, HB Calorimetric method, and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: MINDRAY BC-3000 Plus 3 part automatic analyzer,

VIKARANTJI

Technologist Page No: 1 of 16 DR.TANU RUNGTA MD (Pathology) RMC No. 17226

This report is not valid for medico legal purpose

^{*}Red Blood Cells (RBC), which carry oxygen -

^{*}White Blood Cells (WBC), which help in fighting against infections -

^{*}Hemoglobin, which is the oxygen carrying protein in the red blood cells -

^{*}Hematocrit (HCT), the proportion of RBC to the fluid component, or plasma present in blood -

^{*}Platelets, which aid in blood clotting



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

© +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age:- 32 Yrs 9 Mon 7 Days

Sex :- Male

Patient ID :-12222381 Date :- 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

09

mm in 1st hr

00 - 15

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases.ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein.ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



VIKARANTJI

Technologist

Page No: 2 of 16

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

9 +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age:- 32 Yrs 9 Mon 7 Days

Sex :- Male



Patient ID :-12222381 Date :- 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



VIKARANTJI

Page No: 3 of 16



 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

● +91 141 4824885 🖨 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age :-Sex :-Male

Patient ID: -12222381 32 Yrs 9 Mon 7 Days

Date: - 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company:-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

BIOCHEMISTRY

	DIOCI	I DITTALO A A A A	
Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Methord:- GOD POD	79.4	mg/dl	70.0 - 115.0
Impaired glucose tolerance (IGT)		111 - 125 mg/dL	
Diabetes Mellitus (DM)		> 126 mg/dL	

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm,

hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin

therapy or various liver diseases.

BLOOD SUGAR PP (Plasma) Methord:- GOD PAP

118.0

mg/dl

70.0 - 140.0

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases .

VIKARANTJI

Technologist Page No: 4 of 16 DR.TANU RUNGTA

MD (Pathology) RMC No. 17226



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

O B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

♦ +91 141 4824885
maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

32 Yrs 9 Mon 7 Days Age :-

Sex :-Male



Patient ID :-12222381

Date :- 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company:-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (HbA16 Methord:- CAPILLARY with EDTA	C) 5.5	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Methord:- Calculated Parameter	111	mg/dL	68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA) Reference Group HbA1c in % Non diabetic adults >=18 years < 5.7 At risk (Prediabetes) 5.7 - 6.4 Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings. Some of the factors that influence HbA1c and its measurement (Adapted from Gallagher et al.)

1. Erythropoiesis

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropolesis.
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease
- 2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH

4. Erythrocyte destruction

- Increased HbA1c: increased erythrocyte life span: Splenectomy.
 Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
 Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Note:

1. Shortened RBC life span -HbA1c test will not be accurate when a person has a condition that affects the average lifespan of red blood cells (RBCs), such as hemolytic anemia or blood loss. When the lifespan of RBCs in circulation is shortened, the A1c result is falsely low and is an unreliable measurement of a person's average glucose over time.

2.Abnormal forms of hemoglobin – The presence of some hemoglobin variants, such as hemoglobin S in sickle cell anemia, may affect certain methods for measuring A1c. In these cases, fructosamine can be used to monitor glucose control.

1.To follow patient for glycemic control test like fructosamine or glycated albumin may be performed instead.
2.Hemoglobin HPLC screen to analyze abnormal hemoglobin variant.

estimated Average Glucose (eAG); based on value calculated according to National Glycohemoglobin Standardization Program (NGSP) criteria

VIKARANTJI

Technologist

Page No: 5 of 16



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

O B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

S +91 141 4824885 S maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

32 Yrs 9 Mon 7 Days Age :-

Sex :-Male



Patient ID: -12222381

Date :- 08/11/2022

09:07:14

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

HAEMATOLOGY

BLOOD GROUP ABO Methord:- Haemagglutination reaction

"A" POSITIVE



VIKARANTJI

Technologist Page No: 6 of 16



O B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

O +91 141 4824885 ☐ maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age :-

32 Yrs 9 Mon 7 Days

Male

Sex :-



Patient ID: -12222381

Date: - 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company:-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE TOTAL CHOLESTEROL Methord:- CHOD-PAP methodology	190.00	mg/dl	Desirable <200 Borderline 200-239 High> 240
InstrumentName: MISPA PLUS Interpretation: (disorders.	Cholesterol measurements	s are used in the diagnosis	and treatments of lipid lipoprotein metabolism
TRIGLYCERIDES Methord:- GPO-TOPS methodology	104.00	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
	The state of the s	2	

InstrumentName: MISPA PLUS Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL Methord:- Selective inhibition Method

Male 35-80 **Female 42-88**

An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to

precipitation methods. LDL CHOLESTEROL Methord:- Calculated Method

104.67

mg/dl

Optimal <100

Near Optimal/above optimal 100-129 Borderline High 130-159

High 160-189 Very High > 190

VLDL CHOLESTEROL

Methord:- Calculated

20.80

mg/dl

0.00 - 80.00

T.CHOLESTEROL/HDL CHOLESTEROL RATIO

Instrument Name: MISPA PLUS Interpretation:

2.79

0.00 - 4.90

LDL / HDL CHOLESTEROL RATIO

1.54

0.00 - 3.50

TOTAL LIPID

552.74

mg/dl

400.00 - 1000.00

- 1. Measurements in the same patient can show physiological& analytical variations. Three serialsamples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.
- 2. As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is
- 3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated fromperipheral tissues.

Comments: 1- ATP III suggested the addition of Non HDL Cholesterol (Total Cholesterol - HDL Cholesterol) as an indicator of all VIKARANTJI Janu

Technologist

Page No: 7 of 16

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

● +91 141 4824885 🖨 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age:- 32 Yrs 9 Mon 7 Days

Sex :- Male



Patient ID :-12222381 Dat

Date :- 08/11/2022

09:07:14

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

BIOCHEMISTRY

atherogenic lipoproteins (mainly LDL & VLDL). The Non HDL Cholesterolis used as a secondary target of therapy in persons with triglycerides >=200 mg/dL. The goal for Non HDL Cholesterol in those with increased triglyceride is 30 mg/dL above that set for LDL Cholesterol.

2 -For calculation of CHD risk, history of smoking, any medication for hypertension & current B.P. levels are required.



VIKARANTJI

Technologist Page No: 8 of 16



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

© +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age:- 32 Yrs 9 Mon 7 Days

Sex :- Male



Patient ID: -12222381

Date :- 08/11/2022

09:07:14

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

BIOCHEMISTRY

LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Methord:- DMSO/Diazo	0.62	mg/dL	Infants: 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Methord:- DMSO/Diazo	0.21	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Methord:- Calculated	0.41	mg/dl	0.30-0.70
SGOT Methord:- IFCC	18.0	U/L	Men- Up to - 37.0 Female - Up to - 31.0
SGPT Methord:- IFCC	24.9	U/L	Men- Up to - 40.0 Female- Up to - 31.0
SERUM ALKALINE PHOSPHATASE Methord:- DGKC - SCE	68.60	U/L	53.00 - 141.00
SERUM GAMMA GT Methord:- Szasz methodology Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than	18.50	U/L ymes in cases of obstructive jaundice and	10.00 - 45.00
metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post- hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 ti	imes normal)are observed w	with infectious hepatitis.	
SERUM TOTAL PROTEIN Methord:- Direct Biuret Reagent	6.76	g/dl	5.10 - 8.00
SERUM ALBUMIN Methord:- Bromocresol Green	4.06	g/dl	2.80 - 4.50
SERUM GLOBULIN Methord:- CALCULATION	2.70	gm/dl	2.20 - 3.50
The state of the s		A STATE OF THE PARTY OF THE PAR	

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note:- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B,C, paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.

VIKARANTJI

A/G RATIO

Technologist Page No: 9 of 16



O B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

🕲 +91 141 4824885 🖨 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI 32 Yrs 9 Mon 7 Days Age :-

Sex :-

Patient ID: -12222381

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company:-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA Methord:- Urease/GLDH 31.10

mg/dl

10.00 - 50.00

InstrumentName: MISPA PLUS Interpretation: Urea measurements are used in the diagnosis and treatment of certain renal and metabolic

SERUM CREATININE Methord:- Jaffe's Method

0.95

mg/dl

Males: 0.6-1.50 mg/dl Females: 0.6 -1.40 mg/dl

Interpretation:

Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not

clinically significant. SERUM URIC ACID

5.65

mg/dl

2.40 - 7.00

InstrumentName: HORIBA YUMIZEN CA60 Daytona plus Interpretation: Elevated Urate: High purine dict, Alcohol. Renal insufficiency, Drugs, Polycythaemia vera, Malignancies, Hypothyroidism, Rare enzyme defects, Downs syndrome, Metabolic syndrome, Pregnancy, Gout.

SODIUM Methord:- ISE

Interpretation: Decreased sodium - Hyponatraemia Causes include: fluid or electrolyte loss, Drugs, Oedematous states, Legionnaire's disease and other chest infections, pseudonatremia, Hyperlipidaemias and paraproteinaemias, endocrine diseases, SIADH.

POTASSIUM

4.13

mmol/L

3.50 - 5.50

A. Elevated potassium (hyperkalaemia). Artefactual, Physiologidal vation, Drugs, Pathological states, Renal failure Adrenocortical insufficiency, metabolic acidoses, very high platelet or white cell counts B. Decreased potassium (hypokalaemia)Drugs, Liquoric, Diarrhoea and vomiting, Metabolic alkalosis, Corticosteroid excess, Oedematous state, Anorexia nervosa/bulimia

CHLORIDE

105.9

mmol/L

94.0 - 110.0

Interpretation: Used for Electrolyte monitoring.

SERUM CALCIUM Methord:- Colorimetric method 11.10

mg/dl

g/dl

8.10 - 11.50

InstrumentName: Rx Daytona plus Interpretation: Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia. Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN VNEDARIA RITCIBiuret Reagent

6.76

5.10 - 8.00

Technologist

Page No: 10 of 16

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226

fanu



 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

© +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. DINESH KUMAR SAINI

Age:- 32 Yrs 9 Mon 7 Days

Sex :- Male

Patient ID :-12222381 Date :- 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company:- Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

BIOCHEMISTRY

SERUM ALBUMIN Methord:- Bromocresol Green	4.06	g/dl	2.80 - 4.50
SERUM GLOBULIN Methor¢:- CALCULATION	2.70	gm/dl	2.20 - 3.50
A/G RATIO	1.50		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR .in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.

VIKARANTJI

Technologist
Page No: 11 of 16



O B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

+91 141 4824885 maxcarediagnostics1@gmail.com

32 Yrs 9 Mon 7 Days

Male Sex :-

Methord:- ECLIA

Patient ID :-12222381

Date :- 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

TOTAL THYROID PROFILE

IMMIINOASSAV

		0.100111	
Test Name	Value	Unit	Biological Ref Interval
THYROID-TRIIODOTHYRONINE T3	1.10	ng/mL	0.70 - 2.04

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simoultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1. Primary hyperthyroidism is accompanied by †serum T3 & T4 values along with *TSH level.2. Low TSH.high FT4 and TSH receptor antibody(TRAb) +ve seen in patients with Graves disease 3.Low TSH,high FT4 and TSH receptor antibody(TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.HighTSH,Low FT4 and Thyroid microsomantibody increased seen in patients with Hashimotos thyroiditis 5.HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency 6.Low

TSH,Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism
7. Primary hypothyroidism is accompanied by 1 serum T3 and T4 values & 'serum TSH levels B. Normal T4 levels accompanied by 1 T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis9. Normal T3 & T4 along with 1 TSH indicate mild / Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild / Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .11. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .12. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroidism .13. Normal T3 & T4 along with 1 TSH indicate mild | Subclinical Hypothyroid

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester: 0.10-2.50 uIU/mL 2nd Trimester: 0.20-3.00 uIU/mL 3rd Trimester: 0.30-3.00 uIU/mL The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher THYROID ace THYROXHTEH (14) is due to a real chance with ace of Anti-Greating proceduring of the Anti-Anti-Burglized thyroid disease in the elderly. *** 5.10 - 14.10

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simoultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidsm is accompanied by resrum T3 & T4 values along with "TSH level.2.Low TSH,high FT4 and TSH receptor antibody(TRAb) +ve seen in patients with Graves disease 3.Low TSH,high FT4 and TSH receptor antibody(TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.HighTSH,Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis 5.HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency 6.Low TSH, Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertlary hypothyroidism 7.Primary, hypothyroidism is accompanied by 1 serum T3 and T4 values & "serum TSH levels.Normal T4 levels accompanied by "T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis9.Normal or "T3 & "T4 along with "TSH indicate mild / Subclinical Hypothyroidism .12.Normal T3 & T4 levels with "TSH indicate Mild / Subclinical Hypoth

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL 2nd Trimester : 0.20-3.00 uIU/mL 3rd Trimester : 0.30-3.00 uIU/mL The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy

REMARK-assay results should be interpreted in context to the clinical condition and associated results of other investigations, Previous treatment with conticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

TSH Methord:- ECLIA 2.760

μIU/mL

0.350 - 5.500

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simoultaneous measurement of TSH with free T4 is useful in evaluating differantial diagnosis

NTERPRETATION-Ultra Sensitive 4th generation assay

Pimary hypertriyroidism is accompanied by †serum T3 & T4 values along with ‡ TSH level.

Technologist Page No: 15 of 16



O B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

• +91 141 4824885 maxcarediagnostics1@gmail.com
NAME:- Mr. DINESH KUMAR SAINI

Age :-

32 Yrs 9 Mon 7 Days

Sex :-

Male



Patient ID: -12222381

Date :- 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

IMMUNOASSAY

2.Low TSH,high FT4 and TSH receptor antibody(TRAb) +ve seen in patients with Graves disease
3.Low TSH,high FT4 and TSH receptor antibody(TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter

A HighTSH,Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis

5. HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with Hashimotos thyroiditis

5. HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with lodine deficiency/Congenital T4 synthesis deficiency

6. Low TSH,Low FT4 and TRH stimulation tost -Delayed response seen in patients with Tertiary hypothyroidism

7. Primary hypothyroidism is accompanied by 1 serum T3 and T4 values & 1 serum TSH levels

8. Normal T4 levels accompanied by † T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis

S.Normal or | T3 & | T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)

10.Normal T3 & T4 along with | T5H indicate mild / Subclinical Hyperthyroidism.

11.Normal T3 & | T4 along with | T5H is seen in Hypothyroidism.

12.Normal T3 & T4 levels with | T5H is dicate Mild / Subclinical Hypothyroidism.

13.Slightly | T3 levels may be found in pregnancy and in estrogen therapy while | levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propanolol.

14.Although † TSH levels are nearly always indicative of Primary Hypothroldism , rarely they can result from TSH secreting pituitary tumours.

DURING PREGNANCY - REFERENCE RANGE for TSH IN UIU/mL (As per American Thyroid Association)

1st Trimester: 0.10-2.50 uIU/mL 2nd Trimester: 0.20-3.00 uIU/mL 3rd Trimester: 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved.TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real chance with age or an increasing proportion of unrecognized thyroid disease in the elderly.

*** End of Report *

VIKARANTJI

Technologist Page No: 16 of 16



 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

• +91 141 4824885 maxcarediagnostics1@gmail.com
NAME:- Mr. DINESH KUMAR SAINI

Age :-32 Yrs 9 Mon 7 Days

Sex :-Male

Patient ID :-12222381 Date :- 08/11/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 08/11/2022 17:44:44

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
	DALENE	I I OW	DATE VELLOW
COLOUR	PALE YEI	LLOW	PALE YELLOW
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH)	6.0		5.0 - 7.5
SPECIFIC GRAVITY	1.025		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL	and the second	NIL
BILIRUBIN	NEGATIV	Æ	NEGATIVE
UROBILINOGEN	NORMAL	. A	NORMAL
KETONES	NEGATIV	E A	NEGATIVE
NITRITE	NEGATIV	E CONTRACTOR	NEGATIVE
MICROSCOPY EXAMINATION	CHARLES AND A STATE OF THE PARTY OF THE PART		
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT	100	

VIKARANTJI

Technologist Page No: 12 of 16



 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

© +91 141 4824885 maxcarediagnostics1@gmail.com



NAME:	MR. DINESH KUMAR SAINI	AGE	32 YRS/M
REF.BY	BANK OF BARODA	DATE	08/11/2022

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected.

Shallni

DR.SHALINI GOEL
M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954



 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023
 +91 141 4824885 maxcarediagnostics1@gmail.com



MR. DINESH KUMAR SAINI	32 Y/Male
Registration Date: 08/11/2022	Ref. by: BANK OF BARODA

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (11.7 cm). Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Right kidney is measuring approx. 11.1 x 4.7 cm.

Left kidney is measuring approx. 11.1 x 4.7 cm.

Urinary bladder does not show any calculus or mass lesion.

Prostate is normal in size with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified. No significant free fluid is seen in pelvis.


IMPRESSION: Normal Study.



DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)


RMC no.: 21954




Ref.: BANK OF BARODA Test Date: 08-Nov-2022(10:26:09) Notch: 50Hz 0.05Hz - 100Hz 10mm/mV 12229451322392/Mr Dinesh Kumar Saini 32Yrs-7Mcinths/Male 3-14, Vidhyanagar Nagar, Enclave, Phase-2, Jaipur P3 HEALTH SOLUTIONS LLP http://www.rmsindia.com.jp.RMS_EGG_(VESTA_v3c0.3) FINDINGS: Normal Sinus Rhythm P-QRS-T axis: 67 • 72 • 54 • (Deg) Vent Rate: 60 bpm; PR Interval: 146 ms; QRS Duration: 92 ms; QT/QTc Int: 372/372 ms Comments: avR 2 ≾ Kgs/ Cms BP: 25mm/Sec Dr. Naresh Kumar Mohanka mmHg HR: 60 bpm 4 ప QT/QTc: 372/372ms P-QRS-T Axis: 67 - 72 - 54 (Deg) QRS Duration: 92 ms PR Interval: 146 ms Dr. Naresh Kurhar Mohanka MBBS, DIP. CARDIO (ESCORTS)
D.E.M. (RCGP-UK) RMC No.: 35703

B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur 1322203/MR DINESH KUMAR SAINI 32 Yrs/Male 0 Kg/0 Cms Date: 08-Nov-2022 10:29:45 AM Ref. By: BANK OF BARODA Protocol: BRUCE

StageTime PhaseTime Speed Grade METS H.R. B.P. R.D. PVG Gommonts 1.0 601 10/080 172 1.0 91 130/80 103 1.0 91 130/80 105 1								
StageTime PhaseTime Speed Glade METS H.R. B.P. R.P.P. VC Comments 1.0 60 170 180 190 190 190 190 190 190 190 190 190 19		O (ESCORTS	P. CARDI	ABBS, DI				
StageTime PhaseTime Speed Grade METS H.R. B.P. R.P.P. PVG Comments 1:0 60 170/80 103		35703	RMC No.:					
StageTime PhaseTime Speed Grade Mr.Ts H.R. B.P. R.P.P. PVG Comments STL L.		ar Mohanka	esh Kum	Dr. Nare				
StageTime PhaseTime Speed Grade METS H.R. B.P. R.P.P. PVG Comments STL Example METS H.R. METS H.R. METS M		7					ments:	Advice/Comr
StageTime PhaseTime Speed Grade METS H.R. B.P. R.P.P. PVC Comments STIL Assume the point 1.0 60 1.0 66 120/80 103			_					
StageTime PhaseTime based Grade METS H.R. B.P. R.P.P. PVC Comments STILL Assumble StageTime PhaseTime based Grade METS H.R. B.P. R.P.P. PVC Comments STILL Assumble StageTime PhaseTime based Grade METS H.R. B.P. R.P.P. PVC Comments STILL Assumble 1.0 60.00 1.0 60.120/30 103 . III Assumble III 3301 3.02 1.7 10.0 4.1 30/80 126 . Ask Ask 21.7 8.18 3.4 14.0 9.5 164 150/85 246 . Ask Ask 2.00 0.0 0.0 1.2 115 150/85 172 . Ask . Ask . Ask . Ask . Ask . . Ask 		7						
StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVG Comments STL Assembly StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVG Comments StageTime StageTime Speed Grade METs H.R. B.P. R.P.P. PVG Comments StageTime StageTime Speed Grade METs H.R. B.P. R.P.P. PVG Comments StageTime StageTime Speed Grade METs H.R. B.P. R.P.P. PVG Comments StageTime StageTime Speed Grade METs H.R. B.P. R.P.P. PVG Comments StageTime StageTime Speed StageTime Speed StageTime Speed StageTime Speed StageTime Speed S	6 9 12 15 18 21							
Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments STL Damidulus Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 2 PR METS MET			7	V	-			
Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments STL Abandology Stage	V6	2,	1	- -	-	4		
Stage Time Phase Time Speed Grade MrETs H.R. B.P. R.P.P. PVC Comments STL D.S. D.S. D. D.S. D. D. D				rance)	fort loter	ned :9.5(Good t	WorkLoad attai	Max
Stage Time Phase Time Speed Grade (n) METS H.R. B.P. R.P.P. PVC Comments STL ASAMADA MMISSI 100 60 120/80 72 1 2 PR MMISSI 110 60 120/80 72 1 2 PR MMISSI 110 86 120/80 103 1 1 2 PR MMISSI 110 86 120/80 103 1 1 2 PR MMISSI 120 1.0 88 120/80 105 1 1 2 PR MMISSI 120 1.0 4.7 120/80 109 1 1 4.7 120/80 109 1 4.7 4.7 120/80 126 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 4.7 <td>V5</td> <td></td> <td></td> <td></td> <td>-</td> <td>mHg)</td> <td>(BP : 160/85(m</td> <td>Max</td>	V5				-	mHg)	(BP : 160/85(m	Max
StageTime PhaseTime Speed Grade METS H.R. B.P. R.P.P. PVG Comments S 1 2 PR 100 600 1.00 86 120/80 72 11 10 86 120/80 103 11 10			le HR 188	Predictab	7% of Max		HR Attained	Max
STREETIME Speed Grade METS H.R. B.P. R.P.P. PVG Comments STR Comments STRE Comments STREETIME Speed Grade METS H.R. B.P. R.P.P. PVG Comments STREETIME Speed Strate	V4 Alaman					:08:17	rcise Time	Exer
Stage Time Phase Time Speed Grade METS H.R. B.P. R.P.P. PVC Comments STL Community Comments Stage Time Phase Time Speed Grade METS H.R. B.P. R.P.P. PVC Comments Stage Time Stage Time Phase Time Speed Stage								Findings:
StageTime PhaseTime Speed Grade METS H.R. B.P. R.P.P. PVC Comments S 1 2 PR Image: Comment Scale	Va Polantina							
Stage(Time Phase(Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Mainson) STL 05.mm/Dbm (1.0 min) (1.0	V2			1.0	0.0	0.0	4:00	Recovery
StageTime PhaseTime Speed Grade (h) METS (h) H, R. (h) B, P. (h) R, P. P. PVC Comments S 1 2 PR (h) 1.0 60 120780 72 - (h)		+		1.0	0.0	0.0	3:00	Recovery
StageTime PhaseTime Speed Grade (mph) METS (h) H.R. (b) B.P. (mmhs) R.P.P. PVC Comments S 1 2 PR Numscol (mms) (h) (V	+		1.0	0.0	0.0	2:00	Recovery
StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1 In Main Scot Minister Main Scot M	V3 avF chromotoped with			1.2	0.0	0.0	1:00	Recovery
Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments STL Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Stage Time Stage Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Stage Time Stage Time Speed Stage Time Stage Time Speed Stage Time Sta	A			9.5	14.0		2:17	PeakEx
Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments STL Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Stage Time Stage Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Stage Time Stage Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Stage Time Stage Time Speed Stage Time Stage Time Speed Stage Time		1	-	7.1	12.0		3:01	Stage 2
Stage Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments STL Stage Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Stage Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments Stage Time Speed Stage Time Stage Time Speed Stage Time				4.7	10.0		3:01	Stage 1
Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments STL				1.0				ExStart
STL Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1				1.0				¥
STL Stage Time Phase Time Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1	-			1.0				Standing
STL StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments (A) (Opin) (mmHg) k100			_	1.0				Supine
StageTime PhaseTime Speed Grade METs H.R. B.P. R.P.P. PVC Comments S 1		+	(bpm)		(%)			
STL	VC Comments s 1	-	H.R	METs	Grade			Stage
								Objective:





12222381 Mr.DINESH KUMAR SAINI 32YRS BANK OF BARODA M 08.NOV.2022 MAXCARE DIAGNOSTIC (ASSOCIATES OF P3 HEALTH SOLUTIONS LLP)

