

akriti Labs

3, Mahatma Gandhi Marg, Gandhi Nagar Mod, Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661 www.aakritilabs.com CIN No. U85195RJ2004PTC019563



: Ms. TANUJA PALIWAL Name

Age/Gender: 31 Y 2 M 1 D/Female

Patient ID : 012305070042

BarcodeNo: 10085026

Referred By: Self

Registration No: 25602

Registered

: 07/May/2023 10:53AM

Analysed

: 09/May/2023 11:01AM

Reported

: 09/May/2023 11:01AM

Panel

MEDI WHEEL (ARCOFEMI

HEALTHCARE LTD)

USG: WHOLE ABDOMEN (Female)

LIVER

: Is normal in size, shape and echogenecity.

The IHBR and hepatic radicals are not dilated. No evidence of focal echopoor/echorich lesion seen. Portal vein diameter and Common bile duct normal in size

GALL

: Is normal in size, shape and echotexture. Walls are smooth and

BLADDER

regular with normal thickness. There is no evidence of cholelithiasis.

PANCREAS: Is normal in size, shape and echotexture. Pancreatic duct is not dilated. : Is normal in size, shape and echogenecity. Spleenic hilum is not dilated. SPLEEN

KIDNEYS: Right Kidney:-Size: 104 x 43 mm, Left Kidney:-Size: 112 x 47 mm.

Bilateral Kidneys are normal in size, shape and echotexture, corticomedullary differentiation is fair and ratio appears normal.

Pelvi calyceal system is normal. No evidence of hydronephrosis/ nephrolithiasis.

22 mm size cystic lesion seen at mid pole of right kidney.

URINARY : Bladder walls are smooth, regular and normal thickness.

BLADDER: No evidence of mass or stone in bladder lumen.

UTERUS

: Uterus is anteverted with normal in size shape & echotexture

Uterine muscular shadows normal echopattern.

Endometrium is normal and centrally placed with size: 3 mm.

No evidence of mass lesion is seen. Size of uterus: 60 x 42 x 30 mm.

ADNEXA :

Both the ovaries are normal in size shape and echotexture.

No mass lesion/ polycystic ovarian cyst is seen.

SPECIFIC: No evidence of retroperitoneal mass or free fluid seen in peritoneal cavity.

NO evidence of lymphadenopathy or mass lesion in retroperitoneum. Visualized bowel loop appear normal. Great vessels appear normal.

IMPRESSION: Right renal cortical cyst

*** End Of Report ***

Page 1 of 1



Dr. Neera Mehta M.B.B.S., D.M.R.D. RMCNO.005807/14853



akriti Labs

3, Mahatma Gandhi Marg, Gandhi Nagar Mod, Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661 www.aakritilabs.com CIN No. U85195RJ2004PTC019563

NAME		MR TANUJA PALIWAL			AGE	31YRS		SEX	FERRALE
REF BY	MEDIWHEEL				DATE	09/05/	2023	REG NO	FEMALE
			ECH	OCARDIOG		DODT	2023	KEG NO	
WINDOV	V- POC	R/ADEQ	UATE/GO	OODVALVE	TO THE INL	FUNI			
MITRAL			NORMA			TRICUSPID			
AORTIC NO			NORMAI	The state of the s		PULMONARY		NORMAL	
2D/M-M					1 OLIVIO	DIVART		NORMAL	
IVSD mm		8.8		IVSS mm	12.9		AODT	w tolerand	
LVID mm		39.2	10 20	LVIS mm	22.7			A mm	20.6
LVPWD n	1.11	8.8		LVPWS mm	13.2		LA mr	n	27.4
CHAMBE	RS			Local Sections of the Control of the	13.2		EF%		60%
	LA			NORMAL					NAME OF
LV			NO	NORMAL		RA RV		NORMAL	
	PERICARDIUM			NORMAL				NORN	MAL
DOPPLER									
PEAK VELOCITY m/s E/A		1.03	3/0.71	PEAK GRADIANT MmHg					
MEAN VEL					MEAN GRADIANT MmHg				
MVA cm2	(PLAN	TMETERY	Y)		MVA cm2 (PHT)		g		
MR				I WAY	CITIZ (FIII)		4		
AORTIC	F								
PEAK VELC			1.30		PEAK	GRADIANT	MmU-		
MEAN VEL	OCITY	m/s			PEAK GRADIANT MmHg MEAN GRADIANT MmHg		21		
AR		as Fig.				SHADIAN	IVIIIH	5	
TRICUSPID					6				Ne Control of the Con
PEAK VELOCITY m/s		0.59	0.59		RADIANT	MmHa			
MEAN VELOCITY m/s			NA NA		PEAK GRADIANT MmHg MEAN GRADIANT MmHg				
TR			BIPS	PASP n	nmHa	WITHIN		100	
PULMONA		100 mm		M	1,01	ining .			
PEAK VELO	CITY m	/s	1.55		PEAK G	RADIANT	Applic		
MEAN VELOCITY m/s				V ~ ~	MEAN	GRADIANT	Manue		1.0
OD.				1417	UNADIANI	IVIMHg			

IMPRESSION

- NORMAL LV SYSTOLIC & DIASTOLIC FUNCTION
- **NO RWMA LVEF 60%**
- NORMAL RV FUNCTION
- NORMAL CHAMBER DIMENSIONS
- NORMAL VALVULAR ECHO
- INTACT IAS / IVS
- NO THROMBUS, NO VEGETATION, NORMAL PERICARDIUM.
- IVC NORMAL

CONCLUSION: FAIR LV FUNCTION.

Cardiologist



Aakriti Labs

3, Mahatma Gandhi Marg, Gandhi Nagar Mod, Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661 www.aakritilabs.com

CIN No. U85195RJ2004PTC019563

Name : Ms. TANUJA PALIWAL

Age/Gender: 31 Y 2 M 1 D/Female

Patient ID : 012305070042

BarcodeNo:10085026

Referred By: Self

Registration No: 25602

Registered

: 07/May/2023 10:53AM

Analysed

: 09/May/2023 01:54PM

Reported

: 09/May/2023 01:54PM

Panel

: MEDI WHEEL (ARCOFEMI

HEALTHCARE LTD)

DIGITAL X-RAY CHEST PA VIEW

Soft tissue shadow and bony cages are normal.

Trachea is central.

Bilateral lung field and both CP angle are clear.

Domes of diaphragm are normally placed.

Transverse diameter of heart appears with normal limits.

IMPRESSION:- NO OBVIOUS ABNORMALITY DETECTED.

wellness partner

*** End Of Report ***

Page 1 of 1



Dr. Neera Mehta M.B.B.S., D.M.R.D. RMCNO.005807/14853

Vent Rate : 90 bpm

PR Interval : 136 ms

QRS Duration: 74 ms

QT/QTc Int : 348/401 ms

P-QRS-T axis: 61.00 59.00 49.00 Allengers ECG (Pisces)(PIS216200529) Aakriti Labs
56307 / MS TANUJA PALIWAL / 31 Yrs / F/ Non Smoker
Heart Rate : 90 bpm / Tested On : 09-May-23 10.30.43 / HF 0.05 Hz - LF 100 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s
/ Refd By.: MEDIWHEEL R 59.00° T 49.00° \$ 61.00° Reported ByDANITIZ GOYAL

ECG





PATIENT NAME: TANUJA PALIWAL

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WE000796**

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00
RECEIVED :07/05/2023 11:14:04
REPORTED :11/05/2023 19:51:31

Test Report Status Final Results Biological Reference Interval Units

н	AEMATOLOGY - CBC		
MEDI WHEEL FULL BODY HEALTH CHECKUP BE	LOW 40FEMALE		
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB)	10.4 Low	12.0 - 15.0	g/dL
METHOD: CYANIDE FREE DETERMINATION RED BLOOD CELL (RBC) COUNT METHOD: ELECTRICAL IMPEDANCE	4.54	3.8 - 4.8	mi l /μL
WHITE BLOOD CELL (WBC) COUNT METHOD: ELECTRICAL IMPEDANCE	8.10	4.0 - 10.0	thou/µL
PLATELET COUNT METHOD: ELECTRONIC IMPEDANCE RBC AND PLATELET INDICES	287	150 - 410	thou/µL
HEMATOCRIT (PCV) METHOD: CALCULATED PARAMETER	32.8 Low	36 - 46	%
MEAN CORPUSCULAR VOLUME (MCV) METHOD: CALCULATED PARAMETER	72.0 Low	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH) METHOD: CALCULATED PARAMETER	22.9 Low	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC) METHOD: CALCULATED PARAMETER	31.8	31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW) METHOD: CALCULATED PARAMETER	15.8 High	11.6 - 14.0	%
MENTZER INDEX	15.9		
MEAN PLATELET VOLUME (MPV) METHOD: CALCULATED PARAMETER	9.9	6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT			
NEUTROPHILS METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	60	40 - 80	%
LYMPHOCYTES METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	34	20 - 40	%
MONOCYTES METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	05	2 - 10	%
EOSINOPHILS METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	01	1 - 6	%
BASOPHILS	00	0 - 2	%

4.86



Dr. Akansha Jain Consultant Pathologist



2.0 - 7.0



Page 1 Of 14

View Details

View Report

METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY

ABSOLUTE NEUTROPHIL COUNT



thou/µL





PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS : C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WE000796**PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00
RECEIVED :07/05/2023 11:14:04
REPORTED :11/05/2023 19:51:31

Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
METHOD: CALCULATED PARAMETER			
ABSOLUTE LYMPHOCYTE COUNT	2.75	1.0 - 3.0	thou/µL
METHOD: CALCULATED PARAMETER			
ABSOLUTE MONOCYTE COUNT	0.40	0.2 - 1.0	thou/μL
METHOD: CALCULATED PARAMETER			
ABSOLUTE EOSINOPHIL COUNT	0.08	0.02 - 0.50	thou/µL
METHOD : CALCULATED PARAMETER			
ABSOLUTE BASOPHIL COUNT	0 Low	0.02 - 0.10	thou/μL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	1.8		

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

Dr. Akansha Ja

Dr. Akansha Jain Consultant Pathologist





Page 2 Of 14

View Details

View Report





REF. DOCTOR: SELF **PATIENT NAME: TANUJA PALIWAL**

CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WE000796

PATIENT ID : TANUF070592251 CLIENT PATIENT ID:

ABHA NO

AGE/SEX :31 Years Female :07/05/2023 00:00:00 DRAWN

RECEIVED: 07/05/2023 11:14:04 REPORTED :11/05/2023 19:51:31

Biological Reference Interval Test Report Status Results <u>Final</u> Units

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD

E.S.R 0 - 20mm at 1 hr

METHOD: AUTOMATED (PHOTOMETRICAL CAPILLARY STOPPED FLOW KINETIC ANALYSIS)"

Interpretation(s)
ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION:

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition CRP is superior to ESR because it is more sensitive and reflects a more rapid change

TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging,

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia
False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, salicylates)

REFERENCE:

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition.

Dr. Akansha Jain Consultant Pathologist



Page 3 Of 14







PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WE000796

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO

AGE/SEX :31 Years Female DRAWN :07/05/2023 00:00:00 RECEIVED: 07/05/2023 11:14:04

REPORTED :11/05/2023 19:51:31

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

TYPE B **ABO GROUP**

METHOD: TUBE AGGLUTINATION

POSITIVE RH TYPE

METHOD: TUBE AGGLUTINATION

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.'

The test is performed by both forward as well as reverse grouping methods.

Dr. Akansha Jain **Consultant Pathologist** Page 4 Of 14













PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WE000796**PATIENT ID : TANUF070592251

PATIENT ID : TANUFU/059225

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00
RECEIVED :07/05/2023 11:14:04
REPORTED :11/05/2023 19:51:31

Test Report Status <u>Final</u> Results Biological Reference Interval Units

RTOCHEMISTR	v

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR) 95 74 - 99 mg/dL

METHOD : GLUCOSE OXIDASE

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE

BLOOD

HBA1C 5.7 Non-diabetic: < 5.7 %

Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 Therapeutic goals: < 7.0 Action suggested: > 8.0 (ADA Guideline 2021)

METHOD: HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC)

ESTIMATED AVERAGE GLUCOSE(EAG) 116.9 High < 116.0 mg/dL

METHOD: CALCULATED PARAMETER

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR) 101 70 - 140 mg/dL

METHOD: GLUCOSE OXIDASE LIPID PROFILE, SERUM

CHOLESTEROL, TOTAL 143 < 200 Desirable mg/dL

200 - 239 Borderline High

>/= 240 High

METHOD : CHOLESTEROL OXIDASE

TRIGLYCERIDES 77 < 150 Normal mg/dL

150 - 199 Borderline High

200 - 499 High >/=500 Very High

 ${\tt METHOD: LIPASE/GPO-PAP\ NO\ CORRECTION}$

METHOD: DIRECT CLEARANCE METHOD

HDL CHOLESTEROL 47 < 40 Low mg/dL

>/=60 High

CHOLESTEROL LDL 81 < 100 Optimal mg/dL

100 - 129

Near optimal/ above optimal

130 - 159
Borderline High
160 - 189 High

>/= 190 Very High

Dr. Akansha Jain Consultant Pathologist



Page 5 Of 14









PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: **0251WE000796**PATIENT ID: TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00
RECEIVED :07/05/2023 11:14:04
REPORTED :11/05/2023 19:51:31

	j	i	
Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
NON HDL CHOLESTEROL	96	Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	
METHOD: CALCULATED PARAMETER VERY LOW DENSITY LIPOPROTEIN CHOL/HDL RAΠΟ	15.4 3.0 Low	= 30.0 3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk 11.0	mg/dL
LDL/HDL RATIO Interpretation(s)	1.7	High Risk 0.5 - 3.0 Desirable/Low Ris 3.1 - 6.0 Borderline/Modera Risk >6.0 High Risk	
LIVER FUNCTION PROFILE, SERUM			
BILIRUBIN, TOTAL	0.29	0 - 1	mg/dL
METHOD: DIAZO WITH SULPHANILIC ACID BILIRUBIN, DIRECT METHOD: DIAZO WITH SULPHANILIC ACID	0.14	0.00 - 0.25	mg/dL
BILIRUBIN, INDIRECT METHOD: CALCULATED PARAMETER	0.15	0.1 - 1.0	mg/dL
TOTAL PROTEIN METHOD: BIURET REACTION, END POINT	6.8	6.4 - 8.2	g/dL
ALBUMIN	4.2	3.8 - 4.4	g/dL
METHOD: BROMOCRESOL GREEN GLOBULIN	2.6	2.0 - 4.1	g/dL
METHOD : CALCULATED PARAMETER ALBUMIN/GLOBULIN RATIO	1.6	1.0 - 2.1	RATIO
METHOD: CALCULATED PARAMETER ASPARTATE AMINOTRANSFERASE(AST/SGOT)	23	0 - 31	U/L
METHOD: TRIS BUFFER NO P5P IFCC / SFBC 37° C ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: TRIS BUFFER NO P5P IFCC / SFBC 37° C	16	0 - 31	U/L

Dr. Akansha Jain Consultant Pathologist





Page 6 Of 14







PATIENT NAME: TANUJA PALIWAL

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WE000796**

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00
RECEIVED :07/05/2023 11:14:04
REPORTED :11/05/2023 19:51:31

Test Report Status <u>Final</u>	Results	Biological Reference	e Interval Units
ALKALINE PHOSPHATASE METHOD: AMP OPTIMISED TO IFCC 37° C	64	39 - 117	U/L
GAMMA GLUTAMYL TRANSFER METHOD: GAMMA GLUTAMYL-3 CARBOXY-		7 - 32	U/L
LACTATE DEHYDROGENASE	385	230 - 460	U/L
BLOOD UREA NITROGEN (BUN	I), SERUM		
BLOOD UREA NITROGEN METHOD: UREASE KINETIC CREATININE, SERUM	5	5.0 - 18.0	mg/dL
CREATININE METHOD: ALKALINE PICRATE NO DEPROT BUN/CREAT RATIO	0.62 EINIZATION	0.6 - 1.2	mg/dL
BUN/CREAT RATIO METHOD: CALCULATED PARAMETER URIC ACID, SERUM	8.06		
URIC ACID METHOD: URICASE PEROXIDASE WITH AS TOTAL PROTEIN, SERUM	4.2 SCORBATE OXIDASE	2.4 - 5.7	mg/dL
TOTAL PROTEIN METHOD: BIURET REACTION, END POINT ALBUMIN, SERUM	6.8	6.4 - 8.3	g/dL
ALBUMIN METHOD: BROMOCRESOL GREEN GLOBULIN	4.2	3.8 - 4.4	g/dL
GLOBULIN	2.6	2.0 - 4.1	g/dL
ELECTROLYTES (NA/K/CL), SI	ERUM		
SODIUM, SERUM METHOD: ION-SELECTIVE ELECTRODE	142.4	137 - 145	mmo l /L
POTASSIUM, SERUM METHOD: ION-SELECTIVE ELECTRODE	4.23	3.6 - 5.0	mmo l /L
CHLORIDE, SERUM METHOD: ION-SELECTIVE ELECTRODE	103.1	98 - 107	mmol/L
Interpretation(s)			<u></u>
Sodium	Potassium	Chloride	

Dr. Akansha Jain Consultant Pathologist





Page 7 Of 14









PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

1ATPUR 302017 9314660100

ACCESSION NO: 0251WE000796 PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO

AGE/SEX :31 Years Female DRAWN :07/05/2023 00:00:00 RECEIVED: 07/05/2023 11:14:04 REPORTED :11/05/2023 19:51:31

Test Report Status Results **Biological Reference Interval** Units **Final**

Decreased in:CCF cirrhosis Decreased in: Low potassium Decreased in: Vomiting, diarrhea, vomiting, diarrhea, excessive intake, prolonged vomiting or diarrhea, renal failure combined with salt sweating, salt-losing RTA types I and II, deprivation, over-treatment with nephropathy, adrenal insufficiency, hyperaldosteronism, Cushing's diuretics, chronic respiratory acidosis, nephrotic syndrome, water syndrome, osmotic diuresis (e.g. diabetic ketoacidosis, excessive intoxication, SIADH. Drugs: hyperglycemia), alkalosis, familial sweating, SIADH, salt-losing thiazides, diuretics, ACE inhibitors, periodic paralysis,trauma nephropathy, porphyria, expansion of chlorpropamide,carbamazepine,anti (transient). Drugs: Adrenergic agents, extracellular fluid volume, depressants (SSRI), antipsychotics. adrenalinsufficiency, diuretics. hyperaldosteronism, metabolic alkalosis. Drugs: chronic laxative, corticosteroids, diuretics. Increased in: Dehydration Increased in: Massive hemolysis, Increased in: Renal failure, nephrotic (excessivesweating, severe severe tissue damage, rhabdomyolysis, syndrome, RTA, dehydration, vomiting or diarrhea), diabetes acidosis, dehydration, renal failure. overtreatment with Addison's disease, RTA type IV, saline, hyperparathyroidism, diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate hyperkalemic familial periodic insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory water intake. Drugs: steroids. paralysis. Drugs: potassium salts, licorice.oral contraceptives. potassium-sparing diuretics, NSAIDs, alkalosis.hyperadrenocorticism. beta-blockers. ACE inhibitors, high-Drugs: acetazolamide, androgens, $dose\ trime tho prim-sulfame tho xazole.$ hydrochlorothiazide, salicylates. Interferences: Severe lipemia or Interferences: Hemolysis of sample, Interferences:Test is helpful in hyperproteinemi, if sodium analysis delayed separation of serum, assessing normal and increased anion involves a dilution step can cause prolonged fist clenching during blood gap metabolic acidosis and in spurious results. The serum sodium drawing, and prolonged tourniquet distinguishing hypercalcemia due to hyperparathyroidism (high serum falls about 1.6 mEq/L for each 100 placement. Very high WBC/PLT counts mg/dL increase in blood glucose. may cause spurious. Plasma potassium chloride) from that due to malignancy levels are normal. (Normal serum chloride)

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides. Decreased in:Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease,

malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol;sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic

index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. GLYCOSYLATED HEMOGLOBIN (HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 2. Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).
The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

- eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
 eAG gives an evaluation of blood glucose levels for the last couple of months.
- 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to :

1. Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss,hemolytic anemia) will falsely lower HbA1c test results.Fructosamine is recommended in these patients which indicates diabetes control over 15 days. 2. Vitamin C & E are reported to falsely lower test results (possibly by inhibiting glycation of hemoglobin.



Consultant Pathologist





Page 8 Of 14







Female

PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

1ATPUR 302017 9314660100

ACCESSION NO: 0251WE000796 PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO

DRAWN :07/05/2023 00:00:00 RECEIVED: 07/05/2023 11:14:04 REPORTED :11/05/2023 19:51:31

:31 Years

AGE/SEX

Test Report Status Results **Biological Reference Interval Final** Units

- 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism,chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.
- 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
- b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
- c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. **Elevated levels** results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in bull bloods activity. The activity and the blood. ALT is found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen

in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease. **GGT** is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to: Myasthenia Gravis, Muscuophy

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic

syndrome Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis
TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic

syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Dr. Akansha Jain Consultant Pathologist



Page 9 Of 14









PATIENT NAME: TANUJA PALIWAL

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WE000796**

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00

RECEIVED : 07/05/2023 11:14:04 REPORTED :11/05/2023 19:51:31

Test Report Status Final Results Biological Reference Interval Units

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, URINE

COLOR

SAMPLE NOT RECEIVED

Dr. Akansha Jain Consultant Pathologist



Page 10 Of 14

Minus Dataile







PATIENT NAME: TANUJA PALIWAL

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : 0251WE000796

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female DRAWN :07/05/2023 00:00:00

RECEIVED : 07/05/2023 11:14:04 REPORTED :11/05/2023 19:51:31

Test Report Status Final Results Biological Reference Interval Units

CYTOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PAPANICOLAOU SMEAR

TEST METHOD

SAMPLE NOT RECEIVED

Dr. Akansha Jain Consultant Pathologist





Page 11 Of 14









PATIENT NAME: TANUJA PALIWAL

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

COLOUR

REF. DOCTOR: SELF
ACCESSION NO: 0251WE000796 AGE

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00

RECEIVED : 07/05/2023 11:14:04 REPORTED :11/05/2023 19:51:31

Test Report Status Final Results Biological Reference Interval Units

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, STOOL

METHOD: GROSS EXAMINATION

SAMPLE NOT RECEIVED

Jakat Jana Jana

Dr. Abhishek Sharma Consultant Microbiologist





View Details

\(\(\text{i} = \text{i} \)

Page 12 Of 14





0.550 - 4.780



PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WE000796**

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00
RECEIVED :07/05/2023 11:14:04
REPORTED :11/05/2023 19:51:31

μIU/mL

Test Report Status <u>Final</u> Results Biological Reference Interval Units

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

THYROID PANEL, SERUM

T3 111.37 60.0 - 181.0 ng/dL

METHOD : CHEMILUMINESCENCE

T4 7.00 4.5 - 10.9 μg/dL

7.00 4.5 - 10.9 μg/dL метнод : CHEMILUMINESCENCE

2.709

TSH (ULTRASENSITIVE)

METHOD: CHEMILUMINESCENCE

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. owidctlparowidctlparBelow mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism

Dr. Akansha Jain Consultant Pathologist



Page 13 Of 14









Units

PATIENT NAME: TANUJA PALIWAL REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WE000796**

PATIENT ID : TANUF070592251

CLIENT PATIENT ID: ABHA NO : AGE/SEX :31 Years Female
DRAWN :07/05/2023 00:00:00
RECEIVED :07/05/2023 11:14:04
REPORTED :11/05/2023 19:51:31

Test Report Status <u>Final</u> Results Biological Reference Interval

8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. **NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.**TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

End Of Report
Please visit www.srlworld.com for related Test Information for this accession

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- 8. Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII,

Dr. Akansha Jain Consultant Pathologist



Page 14 Of 14





