

BMI CHART

Hiranandani Fortis Hospital Mini Seashore Road, Sector 10 - A, Vashi, Navi Mumbai - 400 703.

Tel.: +91-22-3919 9222 Fax: +91-22-3919 9220/21 Email: vashi@vashihospital.com

Date: 12/1428

Signature

		Heig	tht (c		(%)									yrs									
EIGHT lbs 10				oms)					W	eighi	likas	3).					BMI		÷				
	00 1		,		-	-				9		-/:			-		J	-					_
	00 1													(2)							¥.1		
kgs 45		05 100	115	120	125	130	135	140	145	150.	155	160	165	170	175	180	185	190	195	200	205	210	2
	5.5 4	7.7 50.5	0 52.3	54.5	56.8	59.1	61.4	63.6	65.9	68.2	70.5	72.7	75.0	77,3	79.5	81.8	84.1	86.4	88.6	90.9	93.2	95.5	9
IGHT in/cm	-] L	Inderwe	ight	-0		Heal	lthy	•	-6		Ove	rweigl	nt ·			Ober	ie.			Ext	reme	y Ob	es
" - 152.4	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	4
* - 154.9	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	36	37	38	39	4
- 157.4	3 19	20	21	22	22	23	24	25	26	27	28	29	30	31	32	33	33	34	35	36	37	38	3
" - 160:0 17	7. 18	19	20	21	22	23	24	24	25	26	27	28	29	30	31	32	32	33	34 *	35	36	37	3
" - 1,62.5	7 18	3 18	19	20	21	22	23	24	24	25	26	27	28	29	30	31	31	32	33	34	35	36	3
" - 165.1	3 17	7 18	19	20	20	21	22	23	24	25	25	26	27	28	29	30	30	31	32	33	34	35	3
* - 167.6	17	7 17	18	19	20	21	21	22	23	24	25	25	26	27	28	29	29	30	31	32	33	34	3
" - 170:1 15	5 16	5 17	18	18	19	20	21	22	22	23	24	25	25	26	27	28	29	29	30	31	32	33	3
* - 172.7 1 5	5 16	6 16	17	18				21						25 ·			28-	28	29	30	31	32	3
" - 176.2 14	1 1:	5 16	17	17	18	19	20	20	21	22	22	23	24	25	25	26	27	28	28	29	30	31	3
0" - 177.8 14	1 15	5 15	16	17	18	18	19	20	20	21	22	23	23	24	25	25	26	27	28	28	29	30	3
1" - 180.3	1 14	1 15	16	16	17	18	18	19	20	21	21	22	23	23	24	25	25	26	27	28	28	29	3
- 182.8	3 14	1 14	15	16	17	17	18	19	19	20	21	21	22	23	23	24	25	25	26	27	27	28	2
- 185.4	3 13	3 14	15	15	16	17	17	18	19	19	20	21	21	22	23	23	24	25	25	26	27	27	2
" - 187.9 12	13	3 14	14	15	16	16	17	18	18	19	19	20	21	21	22	23	23	24	25	25	26	27	2
" - 190.5 12	2 13	3 13	14	15	15	16	16	17	18			-	-	21			-	The second	-	-	-	-	2
" - 193.0 12	2 12	2 13	14 .	14	15	15	16	17	17	18	18	19	20	20	21	22	22	23	23	24	25	25	2

runanamam ricanneare PVI. Ltd. Mini Sea Shore Road, Sector 10 -A, Vashi, Navi Mumbai - 400703

Board Line: 022 - 39199222 | Fax: 022 - 39199220

Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199222 | Health Checkup: 022 - 39199300

www.fortishealthcare.com

CIN: U85100MH2005PTC154823

GST IN: 27AABCH5894D1ZG | PAN NO: AABCH5894D





(A 12 Fortis Network Hospital)

OPD Opthal 14		Health Check-up		
	Mr.Anuj Kumar Tripathi	Sex	Male	Age 38
UHID	8064047	Date	12/12/2	023

Her Pre-HTW.

Drug allergy: -> Not Know

Mini Sea Shore Road, Sector 10 -A, Vashi, Navi Mumbai - 400703

Board Line: 022 - 39199222 | Fax: 022 - 39199220 Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199222 | Health Checkup: 022 - 39199300

www.fortishealthcare.com |

CIN: U85100MH2005PTC154823

GST IN: 27AABCH5894D1ZG | PAN NO: AABCH5894D





Date	12/12/2	023	
			38
7.50 50			
	Sex	Sex Male	Date 12/12/2023 Sex Male Age Health Check-up

OF

- generalized attrition

Stains calculus 444
Almited
RC = 61 (Toptve)

Drug allergy: Sys illness:

Freatment

Ald-OScaling Grade III

PC C - 6/6

Dr. Tugeti





REF. DOCTOR:



PATIENT NAME: MR.ANUJ KUMAR TRIPATHI

CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001 ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00
RECEIVED :12/12/2023 09:02:41
REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status <u>Final</u> Results Biological Reference Interval Units

	HA	EMATOLOGY - CBC		
	CBC-5, EDTA WHOLE BLOOD			
5	BLOOD COUNTS, EDTA WHOLE BLOOD			
	HEMOGLOBIN (HB) METHOD: SLS METHOD	14.5	13.0 - 17.0	g/dL
	RED BLOOD CELL (RBC) COUNT METHOD: HYDRODYNAMIC FOCUSING	5.01	4.5 - 5.5	mil/μL
	WHITE BLOOD CELL (WBC) COUNT METHOD: FLUORESCENCE FLOW CYTOMETRY	7.29	4.0 - 10.0	thou/µL
	PLATELET COUNT METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION	226	150 - 410	thou/µL
	RBC AND PLATELET INDICES			
	HEMATOCRIT (PCV)	44.0	40.0 50.0	
	METHOD : CUMULATIVE PULSE HEIGHT DETECTION METHOD	44.0	40.0 - 50.0	%
	MEAN CORPUSCULAR VOLUME (MCV) METHOD: CALCULATED PARAMETER	87.8	83.0 - 101.0	fL
	MEAN CORPUSCULAR HEMOGLOBIN (MCH) METHOD: CALCULATED PARAMETER	28.9	27.0 - 32.0	pg
	MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION(MCHC) METHOD: CALCULATED PARAMETER	33.0	31.5 - 34.5	g/dL
	RED CELL DISTRIBUTION WIDTH (RDW) METHOD: CALCULATED PARAMETER	13.0	11.6 - 14.0	%
	MENTZER INDEX	17.5		
	METHOD: CALCULATED PARAMETER			
	MEAN PLATELET VOLUME (MPV)	12.5 High	6.8 - 10.9	fL
	METHOD: CALCULATED PARAMETER			

WBC DIFFERENTIAL COUNT



Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) Consultant Pathologist Page 1 Of 17





View Details

View Report











CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

REF. DOCTOR:

PATIENT ID : FH.8064047

CLIENT PATIENT ID: UID:8064047

ACCESSION NO: 0022WL001984

ABHA NO

AGE/SEX :38 Years Male DRAWN :12/12/2023 08:59:00

RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units				
NEUTROPHILS METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING	65	40.0 - 80.0	%			
LYMPHOCYTES METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING	29	20.0 - 40.0	%			
MONOCYTES METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING	4	2.0 - 10.0	%			
EOSINOPHILS METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING	2	1 - 6	%			
BASOPHILS METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING	0	0 - 2	%			
ABSOLUTE NEUTROPHIL COUNT METHOD: CALCULATED PARAMETER	4.74	2.0 - 7.0	thou/μL			
ABSOLUTE LYMPHOCYTE COUNT METHOD : CALCULATED PARAMETER	2.11	1.0 - 3.0	thou/μL			
ABSOLUTE MONOCYTE COUNT METHOD : CALCULATED PARAMETER	0.29	0.2 - 1.0	thou/μL			
ABSOLUTE EOSINOPHIL COUNT METHOD: CALCULATED PARAMETER	0.15	0.02 - 0.50	thou/μL			
ABSOLUTE BASOPHIL COUNT METHOD: CALCULATED PARAMETER	0.00 Low	0.02 - 0.10	thou/μL			
NEUTROPHIL LYMPHOCYTE RATIO (NLR) METHOD: CALCULATED	2.2					

MORPHOLOGY

RBC

METHOD: MICROSCOPIC EXAMINATION

WBC

METHOD: MICROSCOPIC EXAMINATION

PLATELETS

METHOD: MICROSCOPIC EXAMINATION

PREDOMINANTLY NORMOCYTIC NORMOCHROMIC

NORMAL MORPHOLOGY

ADEQUATE



Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**





Page 2 Of 17















CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

REF. DOCTOR:

ACCESSION NO: 0022WL001984 PATIENT ID

: FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male :12/12/2023 08:59:00 DRAWN

RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval

Units

Interpretation(s)

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) In patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.4, 46.1% COVID-19 patients with mild disease might become severe.

3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504. This ratio element is a calculated parameter and out of NABL scope.

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**





Page 3 Of 17

View Details

View Report



PERFORMED AT :

Email: -







CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

REF. DOCTOR :

ACCESSION NO: 0022WL001984 PATIENT ID : FH 8064047

CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

METHOD: WESTERGREN METHOD

Test Report Status

Final

Results

Biological Reference Interval

HAEMATOLOGY

ERYTHROCYTE SEDIMENTATION RATE (ESR), EDTA BLOOD

E.S.R

0 - 14

mm at 1 hr

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD

HBA1C

8.2 High

Non-diabetic: < 5.7

0/

Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5Therapeutic goals: < 7.0 Action suggested: > 8.0

(ADA Guideline 2021)

METHOD: HB VARIANT (HPLC)

METHOD: CALCULATED PARAMETER

ESTIMATED AVERAGE GLUCOSE(EAG)

188.6 High

< 116.0

mg/dL

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-**TEST DESCRIPTION**:
ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-**TEST DESCRIPTION**:
Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy,

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm/hr(95 if anemic). ESR returns to normal 4th week post partum,

Decreased in: Polycythermia vera, Sickle cell anemia

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc.), Hypercholesterolemia
False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, salicylates)




Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377)

Consultant Pathologist



Page 4 Of 17

View Report



PERFORMED AT :









CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI. MUMBAI 440001

REF. DOCTOR :

ACCESSION NO: 0022WL001984 PATIENT ID : FH.8064047

CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval

Units

REFERENCE :

- 1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:
- Evaluating the long-te
 Diagnosing diabetes. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 3. Identifying patients at increased risk for diabetes (prediabetes).

3. Identifying patients at increased risk for diabetes (preciabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

1. eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.

2. eAG gives an evaluation of blood glucose levels for the last couple of months.

3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c - 46.7

HbA1c Estimation can get affected due to :

- 1. Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

 2. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperblirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.

 4. Interference of hemoglobinopathies in HbA1c estimation is seen in

- Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
- b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
 c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) Consultant Pathologist



Page 5 Of 17

View Details

View Report



PERFORMED AT:









Male

PATIENT NAME: MR.ANUJ KUMAR TRIPATHI

CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval Units

IMMUNOHAEMATOLOGY

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP

TYPE B

METHOD: TUBE AGGLUTINATION

RH TYPE

POSITIVE

METHOD: TUBE AGGLUTINATION

Interpretation(s)
ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) Consultant Pathologist

Page 6 Of 17







PERFORMED AT:









CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

REF. DOCTOR: ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047

CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male DRAWN :12/12/2023 08:59:00

RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval Units

		BIOCHEMISTRY		
	LIVER FUNCTION PROFILE, SERUM			
1	BILIRUBIN, TOTAL METHOD: JENDRASSIK AND GROFF	0.92	0.2 - 1.0	mg/dL
	BILIRUBIN, DIRECT METHOD: JENDRASSIK AND GROFF	0.19	0.0 - 0.2	mg/dL
	BILIRUBIN, INDIRECT METHOD: CALCULATED PARAMETER	0.73	0.1 - 1.0	mg/dL
	TOTAL PROTEIN METHOD: BIURET	7.4	6.4 - 8.2	g/dL
	ALBUMIN METHOD: BCP DYE BINDING	3.9	3.4 - 5.0	g/dL
	GLOBULIN METHOD: CALCULATED PARAMETER	3.5	2.0 - 4.1	g/dL
	ALBUMIN/GLOBULIN RATIO METHOD: CALCULATED PARAMETER	1.1	1.0 - 2.1	RATIO
	ASPARTATE AMINOTRANSFERASE(AST/SGOT) METHOD: UV WITH P5P	18	15 - 37	U/L
	ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: UV WITH P5P	35	< 45.0	U/L
	ALKALINE PHOSPHATASE METHOD: PNPP-ANP	75	30 - 120	U/L
	GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: GAMMA GLUTAMYLCARBOXY 4NITROANILIDE	35	15 - 85	U/L
	LACTATE DEHYDROGENASE METHOD: LACTATE -PYRUVATE	111	85 - 227	U/L

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR)

164 High

Normal: < 100

mg/dL

Pre-diabetes: 100-125 Diabetes: >/=126

METHOD: HEXOKINASE

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**

Page 7 Of 17















CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years

Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421

CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval Units

KIDNEY PANEL - 1

BLOOD UREA NITROGEN (BUN), SERUM

GLOMERULAR FILTRATION RATE (MALE)

BLOOD UREA NITROGEN

METHOD: UREASE - UV

6 - 20

mg/dL

CREATININE EGFR- EPI

CREATININE

0.94

0.90 - 1.30

mg/dL

vears

METHOD: ALKALINE PICRATE KINETIC JAFFES AGE

38

106.41

Refer Interpretation Below

mL/min/1.73m2

METHOD: CALCULATED PARAMETER

METHOD: CALCULATED PARAMETER

BUN/CREAT RATIO

BUN/CREAT RATIO

7.45

5.00 - 15.00

URIC ACID, SERUM

URIC ACID

METHOD: URICASE UV

4.8

3.5 - 7.2

mg/dL

TOTAL PROTEIN, SERUM

TOTAL PROTEIN METHOD : BIURET

7.4

6.4 - 8.2

g/dL


Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**

Page 8 Of 17





View Report















Male

PATIENT NAME: MR.ANUJ KUMAR TRIPATHI

CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

ACCESSION NO: 0022WL001984

: FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

PATIENT ID

AGE/SEX :38 Years

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status Final Results Biological Reference Interval Units

ALBUMIN, SERUM

ALBUMIN 3.9 3.4 - 5.0g/dL METHOD: BCP DYE BINDING

GLOBULIN

GLOBULTN 3.5 2.0 - 4.1g/dL METHOD: CALCULATED PARAMETER

ELECTROLYTES (NA/K/CL), SERUM

136	SODIUM, SERUM 136 METHOD: ISE INDIRECT	66 - 145 mmol/l	L
3.50	POTASSIUM, SERUM METHOD: ISE INDIRECT 4.52	50 - 5.10 mmol/l	L
98 -	CHLORIDE, SERUM 101	3 - 107 mmol/l	L

Interpretation(s)

Interpretation(s)

Interpretation(s)
LIVER FUNCTION PROFILE, SERUMBilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.



Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377)

Consultant Pathologist





Page 9 Of 17

View Report



PERFORMED AT:

Agilus Diagnostics Ltd. Hiranandani Hospital-Vashi, Mini Seashore Road, Sector 10, Navi Mumbai, 400703 Maharashtra, India Tel: 022-39199222,022-49723322,

CIN - U74899PB1995PLC045956

Email: -







Male

PATIENT NAME: MR.ANUJ KUMAR TRIPATHI

Final

CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

REF. DOCTOR :

ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

Test Report Status

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Results

Biological Reference Interval

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver,liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity.ALT test measures the amount of this enzyme in the blood.ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatitis, increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic

hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver, Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

urine.

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids, phenytoin, estrogen, thiazides.

Decreased in:Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical, stomach, fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g. galactosemia), Drugs-insulin, ethanol, propranolol; sulfonylureas, tolbutamide, and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin/(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

BLOOD URRA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet; Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE EGFR- EPI-- Kidney disease outcomes quality initiative (KDOQI) guidelines state that estimation of GFR is the best overall indices of the Kidney function.

- It gives a rough measure of number of functioning nephrons. Reduction in GFR implies progression of underlying disease.

- The GFR is a calculation based on serum creatinine test.

- Creatinine is mainly derived from the metabolism of creatine in muscle, and its generation is proportional to the total muscle mass. As a result, mean creatinine general.

- The GFR is a calculation based on serum creatinine test.

- Creatinine is mainly derived from the metabolism of creatinine in muscle, and its generation is proportional to the total muscle mass. As a result, mean creatinine generation is higher in men than in women, in younger than in older individuals, and in blacks than in whites.

- Creatinine is filtered from the blood by the kidneys and excreted into urine at a relatively steady rate.

- When kidney function is compromised, excretion of creatinine decreases with a consequent increase in blood creatinine levels. With the creatinine test, a reasonable estimate of the actual GFR can be determined.

- This equation takes into account several factors that impact creatinine production, including age, gender, and race.

- CKD EPI (Chronic kidney disease epidemiology collaboration) equation performed better than MDRD equation especially when GFR is high(>60 ml/min per 1.73m2).. This formula has less bias and greater accuracy which helps in early diagnosis and also reduces the rate of false positive diagnosis of CKD.

National Kidney Foundation (NKF) and the American Society of Nephrology (ASN).
Estimated GFR Calculated Using the CKD-EPI equation-https://testguide.labmed.uw.edu/guideline/egfr
Ghuman JK, et al. Impact of Removing Race Variable on CKD Classification Using the Creatinine-Based 2021 CKD-EPI Equation. Kidney Med 2022, 4:100471. 35756325
Harrison's Principle of Internal Medicine, 21st ed. pg 62 and 334
URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome Causes of decreased levels-Low Zinc intake, OCP, Multiple Sclerosis
TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.
Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.

Athati

Maharashtra, India

Page 10 Of 17

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**





View Details

View Report



Tel: 022-39199222,022-49723322, CIN - U74899PB1995PLC045956 Email: -





REF. DOCTOR:



PATIENT NAME: MR.ANUJ KUMAR TRIPATHI

CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

· UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval Units

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.



Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**



Page 11 Of 17



PERFORMED AT :







CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

PATIENT ID : FH.8064047

CLIENT PATIENT ID: UID:8064047

ACCESSION NO: 0022WL001984

ABHA NO

AGE/SEX :38 Years Male DRAWN

:12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421

CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval

Units

BIOCHEMISTRY - LIPID

LIPID PROFILE, SE	RUM
-------------------	-----

CHOLESTEROL, TOTAL

168

< 200 Desirable

mg/dL

200 - 239 Borderline High

>/= 240 High

METHOD: ENZYMATIC/COLORIMETRIC, CHOLESTEROL OXIDASE, ESTERASE, PEROXIDASE

TRIGLYCERIDES

228 High

< 150 Normal

mg/dL

150 - 199 Borderline High

200 - 499 High

>/=500 Very High

< 40 Low

mg/dL

METHOD : DIRECT MEASURE - PEG

METHOD : ENZYMATIC ASSAY HDL CHOLESTEROL

LDL CHOLESTEROL, DIRECT

83

47

< 100 Optimal

>/=60 High

mg/dL

100 - 129 Near or above optimal

130 - 159 Borderline High

160 - 189 High >/= 190 Very High

METHOD: DIRECT MEASURE WITHOUT SAMPLE PRETREATMENT

NON HDL CHOLESTEROL

121

Desirable: Less than 130

mg/dL Above Desirable: 130 - 159 Borderline High: 160 - 189

High: 190 - 219

Very high: > or = 220

METHOD: CALCULATED PARAMETER

VERY LOW DENSITY LIPOPROTEIN

45.6 High

</=30.0

mg/dL

METHOD: CALCULATED PARAMETER

CHOL/HDL RATIO

3.6

4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk

> 11.0 High Risk

3.3 - 4.4 Low Risk

METHOD: CALCULATED PARAMETER



Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**

Page 12 Of 17





View Report















CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
LDL/HDL RATIO	1.8	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk
METHOD CALCULATED BARRANETTE		>6.0 High Risk

METHOD: CALCULATED PARAMETER

Interpretation(s)

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**



Page 13 Of 17

View Details



PERFORMED AT :

Email: -







CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAT 440001

REF. DOCTOR:

ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421

CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval

Units

CLINICAL PATH - URINALYSIS

URINALYSIS

PHYSICAL EXAMINATION, URINE

COLOR

PALE YELLOW

METHOD: PHYSICAL

APPEARANCE

CLEAR

METHOD: VISUAL

CHEMICAL EXAMINATION, URINE

PH

6.0

4.7 - 7.5

METHOD: REFLECTANCE SPECTROPHOTOMETRY- DOUBLE INDICATOR METHOD

SPECIFIC GRAVITY

<=1.005

1.003 - 1.035

METHOD: REFLECTANCE SPECTROPHOTOMETRY (APPARENT PKA CHANGE OF PRETREATED POLYELECTROLYTES IN RELATION TO IONIC CONCENTRATION)

PROTEIN

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY - PROTEIN-ERROR-OF-INDICATOR PRINCIPLE

GLUCOSE

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, DOUBLE SEQUENTIAL ENZYME REACTION-GOD/POD

KETONES

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, ROTHERA'S PRINCIPLE

BLOOD

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, PEROXIDASE LIKE ACTIVITY OF HAEMOGLOBIN

BILIRUBIN

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, DIAZOTIZATION- COUPLING OF BILIRUBIN WITH DIAZOTIZED SALT UROBILINOGEN

NORMAL

NORMAL

METHOD: REFLECTANCE SPECTROPHOTOMETRY (MODIFIED EHRLICH REACTION)

LEUKOCYTE ESTERASE

NITRITE NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, CONVERSION OF NITRATE TO NITRITE

NOT DETECTED NOT DETECTED

NOT DETECTED METHOD: REFLECTANCE SPECTROPHOTOMETRY, ESTERASE HYDROLYSIS ACTIVITY

Dr. Rekha Nair, MD (Reg No. MMC 2001/06/2354) Microbiologist

Page 14 Of 17







Dr. Akshay Dhotre, MD

Consultant Pathologist

(Reg,no. MMC 2019/09/6377)









CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41 REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421

CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Results **Final**

Biological Reference Interval Units

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS

METHOD: MICROSCOPIC EXAMINATION

PUS CELL (WBC'S)

METHOD: MICROSCOPIC EXAMINATION

EPITHELIAL CELLS

METHOD: MICROSCOPIC EXAMINATION

CASTS

METHOD: MICROSCOPIC EXAMINATION CRYSTALS

METHOD: MICROSCOPIC EXAMINATION **BACTERIA**

METHOD: MICROSCOPIC EXAMINATION

YEAST

METHOD: MICROSCOPIC EXAMINATION

REMARKS

1-2

0 - 1

NOT DETECTED

NOT DETECTED

/HPF

0-5

0-5

/HPF /HPF

NOT DETECTED

NOT DETECTED

NOT DETECTED

NOT DETECTED

NOT DETECTED

NOT DETECTED

URINARY MICROSCOPIC EXAMINATION DONE ON URINARY CENTRIFUGED SEDIMENT.

Interpretation(s)

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**

Rucha. N

Dr. Rekha Nair, MD (Reg No. MMC 2001/06/2354) Microbiologist





Page 15 Of 17



PERFORMED AT:





CODE/NAME & ADDRESS: C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

REF. DOCTOR: ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047

CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX : 38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

METHOD: ELECTROCHEMILUMINESCENCE, SANDWICH IMMUNOASSAY

Results

Biological Reference Interval Units

SPECIALISED CHEMISTRY - HORMONE

THYROID PANEL, SERUM

109.6 80.0 - 200.0 ng/dL METHOD: ELECTROCHEMILUMINESCENCE IMMUNOASSAY, COMPETITIVE PRINCIPLE 7.71 5.10 - 14.10 µg/dL METHOD: ELECTROCHEMILUMINESCENCE IMMUNOASSAY, COMPETITIVE PRINCIPLE TSH (ULTRASENSITIVE) 1.270 0.270 - 4.200µIU/mL

Interpretation(s)



Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**



Page 16 Of 17



PERFORMED AT:

Agilus Diagnostics Ltd. Hiranandani Hospital-Vashi, Mini Seashore Road, Sector 10, Navi Mumbai, 400703 Maharashtra, India

Tel: 022-39199222,022-49723322, CIN - U74899PB1995PLC045956

Email: -





CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001

REF. DOCTOR: ACCESSION NO: 0022WL001984

PATIENT ID : FH.8064047

CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male

DRAWN :12/12/2023 08:59:00 RECEIVED: 12/12/2023 09:02:41

REPORTED :12/12/2023 14:48:42

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval

Units

SPECIALISED CHEMISTRY - TUMOR MARKER

PROSTATE SPECIFIC ANTIGEN, SERUM

PROSTATE SPECIFIC ANTIGEN

0.361

0.0 - 1.4

ng/mL

METHOD: ELECTROCHEMILUMINESCENCE, SANDWICH IMMUNOASSAY

Interpretation(s)

PROSTATE SPECIFIC ANTIGEN, SERUM-- PSA is detected in the male patients with normal, benign hyperplastic and malignant prostate tissue and in patients with prostatitis.

- PSA is not detected (or detected at very low levels) in the patients without prostate tissue (because of radical prostatectomy or cystoprostatectomy) and also in the female

patients.

- It a suitable marker for monitoring of patients with Prostate Cancer and it is better to be used in conjunction with other diagnostic procedures.

- Serial PSA levels can help determine the success of prostatectomy and the need for further treatment, such as radiation, endocrine or chemotherapy and useful in detecting residual disease and early recurrence of tumor.

- Elevated levels of PSA can be also observed in the patients with non-malignant diseases like Prostatitis and Benign Prostatic Hyperplasia.

- Specimens for total PSA assay should be obtained before biopsy, prostatectomy or prostatic massage, since manipulation of the prostate gland may lead to elevated PSA (false positive) levels persisting up to 3 weeks. (false positive) levels persisting up to 3 weeks.

- As per American urological guidelines, PSA screening is recommended for early detection of Prostate cancer above the age of 40 years. Following Age specific reference

- Age can be used as a guide lines.

- Measurement of total PSA alone may not clearly distinguish between benign prostatic hyperplasia (BPH) from cancer, this is especially true for the total PSA values

between 4-10 ng/mL.

- Total PSA values determined on patient samples by different testing procedures cannot be directly compared with one another and could be the cause of erroneous medical interpretations. Recommended follow up on same platform as patient result can vary due to differences in assay method and reagent specificity.

Burtis CA, Ashwood ER, Bruns DE, Teitz textbook of clinical chemistry and Molecular Diagnostics. 4th edition.
 Williamson MA, Snyder LM. Wallach's interpretation of diagnostic tests. 9th edition.

End Of Report Please visit www.agilusdiagnostics.com for related Test Information for this accession




Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) **Consultant Pathologist**





Page 17 Of 17

View Report







REF. DOCTOR:



PATIENT NAME: MR.ANUJ KUMAR TRIPATHI

CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI, MUMBAI 440001 ACCESSION NO: 0022WL002024

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO :

AGE/SEX :38 Years Male
DRAWN :12/12/2023 11:31:00

RECEIVED : 12/12/2023 11:32:30 REPORTED :12/12/2023 12:18:57

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

METHOD : HEXOKINASE

Final

Results

Biological Reference Interval

Units

BIOCHEMISTRY

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR)

226 High

70 - 140

mg/dL

Interpretation(s)
GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c

End Of Report
Please visit www.agilusdiagnostics.com for related Test Information for this accession

(HOURS

Dr. Akshay Dhotre, MD (Reg,no. MMC 2019/09/6377) Consultant Pathologist



Page 1 Of 1

View Details

View Report



PERFORMED AT :

Agilus Diagnostics Ltd. Hiranandani Hospital-Vashi, Mini Seashore Road, Sector 10, Navi Mumbai, 400703 Maharashtra, India Tel: 022-39199222,022-49723322,

CIN - U74899PB1995PLC045956

Email:-







CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

ACCESSION NO : 0022WL001989

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ABHA NO

AGE/SEX :38 Years Male DRAWN :12/12/2023 09:13:00

RECEIVED: 12/12/2023 09:15:45

REPORTED :12/12/2023 10:24:29

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421

CORP-OPD

BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval

Units

CLINICAL PATH - STOOL ANALYSIS

STOOL: OVA & PARASITE

PHYSICAL EXAMINATION, STOOL

COLOUR

METHOD: VISUAL CONSISTENCY

METHOD: VISUAL

MUCUS

METHOD: VISUAL

VISIBLE BLOOD

METHOD: VISUAL

BROWN

WELL FORMED

NOT DETECTED

NOT DETECTED

ABSENT ABSENT

CHEMICAL EXAMINATION, STOOL

OCCULT BLOOD

NOT DETECTED

NOT DETECTED

MICROSCOPIC EXAMINATION, STOOL

PUS CELLS

METHOD: MICROSCOPIC EXAMINATION

METHOD: GUAIAC ACID METHOD

RED BLOOD CELLS

METHOD: MICROSCOPIC EXAMINATION

CYSTS

METHOD: MICROSCOPIC EXAMINATION

OVA

METHOD: MICROSCOPIC EXAMINATION

LARVAE

METHOD: MICROSCOPIC EXAMINATION

TROPHOZOITES

METHOD: MICROSCOPIC EXAMINATION

1-2

NOT DETECTED

NOT DETECTED

NOT DETECTED

/HPF

/hpf

NOT DETECTED

NOT DETECTED

NOT DETECTED

NOT DETECTED

NOT DETECTED

NOT DETECTED

Dr. Rekha Nair, MD (Reg No. MMC 2001/06/2354) Microbiologist



Page 1 Of 2

View Report



Agilus Diagnostics Ltd. Hiranandani Hospital-Vashi, Mini Seashore Road, Sector 10, Navi Mumbai, 400703 Maharashtra, India Tel: 022-39199222,022-49723322,

CIN - U74899PB1995PLC045956

Email: -











CODE/NAME & ADDRESS : C000045507

FORTIS VASHI-CHC -SPLZD FORTIS HOSPITAL # VASHI,

MUMBAI 440001

REF. DOCTOR:

PATIENT ID : FH.8064047 CLIENT PATIENT ID: UID:8064047

ACCESSION NO: 0022WL001989

ABHA NO

Male AGE/SEX :38 Years

:12/12/2023 09:13:00 DRAWN RECEIVED: 12/12/2023 09:15:45 REPORTED: 12/12/2023 10:24:29

CLINICAL INFORMATION:

UID:8064047 REQNO-1636421 CORP-OPD BILLNO-1501230PCR069824 BILLNO-1501230PCR069824

Test Report Status

Final

Results

Biological Reference Interval Units

Interpretation(s)

End Of Report Please visit www.agilusdiagnostics.com for related Test Information for this accession

Rucha. N

Dr. Rekha Nair, MD (Reg No. MMC 2001/06/2354) Microbiologist



Page 2 Of 2

View Report



Agilus Diagnostics Ltd. Hiranandani Hospital-Vashi, Mini Seashore Road, Sector 10, Navi Mumbai, 400703

Maharashtra, India Tel: 022-39199222,022-49723322, CIN - U74899PB1995PLC045956

Email: -



IC	(James)						E
MA 96:55:54 AM	axis, V-rate 50- 99 QRS area>0 in V2			92	94		F 50~ 0.50-100 Hz W
	normal P	- OTHERWISE NORMAL ECG - Unconfirmed Diagnosis	\$	Z -	-5		mm/mV Chest: 10.0 mm/mV
мале	. Abnormal R-wave progression, early transition.	52 53 8 Standard Placement	W. T.	TAR	ave.		Speed: 25 mm/sec Limb: 10
	Rate 66 PR 129 QRSD 89 QT 370 QTC 388	AXIS P 52 QRS 53 T 8 12 Lead; Stand	Н		I		Device:

Mini Sea Shore Road, Sector 10-A, Vashi, Navi Mumbai - 400703.

Board Line: 022 - 39199222 | Fax: 022 - 39133220 Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199200 | Health Checkup: 022 - 39199300

www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN: 27AABCH5894D1ZG PAN NO: AABCH5894D





DEPARTMENT OF NIC

Date: 12/Dec/2023

Name: Mr. Anuj Kumar Tripathi Age | Sex: 38 YEAR(S) | Male

Order Station: FO-OPD

Bed Name:

UHID | Episode No: 8064047 | 71058/23/1501 Order No | Order Date: 1501/PN/OP/2312/147512 | 12-Dec-2023 Admitted On | Reporting Date: 12-Dec-2023 16:41:35

Order Doctor Name: Dr.SELF.

ECHOCARDIOGRAPHY TRANSTHORACIC

FINDINGS:

- No left ventricle regional wall motion abnormality at rest.
- Normal left ventricle systolic function. LVEF = 60%.
- · No left ventricle diastolic dysfunction.
- · No left ventricle hypertrophy. No left ventricle dilatation.
- · Structurally normal valves.
- No mitral regurgitation.
- No aortic regurgitation. No aortic stenosis.
- No tricuspid regurgitation. No pulmonary hypertension.
- · Intact IAS and IVS.
- No left ventricle clot/vegetation/pericardial effusion.
 - Normal right atrium and right ventricle dimensions.
 - Normal left atrium and left ventricle dimension.
 - Normal right ventricle systolic function. No hepatic congestion.

M-MODE MEASUREMENTS:

LA	22	
AO Root	32	mm
	20	mm
AO CUSP SEP	16	min
LVID (s)	27	
LVID (d)		mm
IVS (d)	42	mm
LVPW (d)	10	mm
	10	mm
RVID (d)	29	mm
RA	28	=====
LVEF		mm
	60	%

Page 2 of 2

Mini Sea Shore Road, Sector 10-A, Vashi, Navi Mumbai - 400703.

Board Line: 022 - 39199222 | Fax: 022 - 39133220

Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199200 | Health Checkup: 022 - 39199300

www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN : 27AABCH5894D1ZG PAN NO : AABCH5894D





DEPARTMENT OF NIC

Date: 12/Dec/2023

Name: Mr. Anuj Kumar Tripathi

Age | Sex: 38 YEAR(S) | Male Order Station : FO-OPD

Bed Name:

UHID | Episode No : 8064047 | 71058/23/1501 Order No | Order Date: 1501/PN/OP/2312/147512 | 12-Dec-2023

Admitted On | Reporting Date: 12-Dec-2023 16:41:35

Order Doctor Name: Dr.SELF.

DOPPLER STUDY:

E WAVE VELOCITY: 0.9 m/sec. A WAVE VELOCITY: 0.5 m/sec

E/A RATIO:1.4

		MEAN (mmHg)	GRADE OF REGURGITATION
MITRAL VALVE	N		Nil
AORTIC VALVE	05		Nil
TRICUSPID VALVE	N		Nil
PULMONARY VALVE	2.0		Nil

Final Impression:

· Normal 2 Dimensional and colour doppler echogardiography study.

DR. PRASHANT PAWAR DNB(MED), DNB (CARD)

DR.AMIT SINGH, MD(MED),DM(CARD) Hiranandani Healthcare Pvt. Ltd.

Mini Sea Shore Road, Sector 10-A, Vashi, Navi Mumbai - 400703.

Board Line: 022 - 39199222 | Fax: 022 - 39133220 Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199200 | Health Checkup: 022 - 39199300

www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN: 27AABCH5894D1ZG PAN NO: AABCH5894D





DEPARTMENT OF RADIOLOGY

Date: 12/Dec/2023

Name: Mr. Anuj Kumar Tripathi UHID | Episode No : 8064047 | 71058/23/1501

 Age | Sex: 38 YEAR(S) | Male
 Order No | Order Date: 1501/PN/OP/2312/147512 | 12-Dec-2023

 Order Station : FO-OPD
 Admitted On | Reporting Date : 12-Dec-2023 13:52:26

Bed Name: Order Doctor Name: Dr.SELF.

X-RAY-CHEST- PA

Findings:

Both lung fields are clear.

The cardiac shadow appears within normal limits.

Trachea and major bronchi appears normal.

Both costophrenic angles are well maintained.

Bony thorax is unremarkable.


DR. YOGINI SHAH

DMRD., DNB. (Radiologist)

miranangani Healthcare Pvt. Ltd.

Mini Sea Shore Road, Sector 10-A, Vashi, Navi Mumbai - 400703.

Board Line: 022 - 39199222 | Fax: 022 - 39133220 Emergency: 022 - 39199100 | Ambulance: 1255



www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN: 27AABCH5894D1ZG PAN NO: AABCH5894D





Patient Name		Anuj Kumar Tripathi	Patient ID		8064047
Sex / Age	:	M / 38Y 5M 7D	Accession No.		PHC.7092127
Modality		US	Scan DateTime		12-12-2023 10:51:40
IPID No		71058/23/1501	ReportDatetime	+	12-12-2023 13:02:20

USG - WHOLE ABDOMEN

LIVER is normal in size and shows mildly raised echogenicity. No IHBR dilatation. No focal lesion is seen in liver. Portal vein appears normal in caliber.

GALL BLADDER is physiologically distended. Gall bladder reveals normal wall thickness. No evidence of calculi in gall bladder. No evidence of pericholecystic collection.

CBD appears normal in caliber.

SPLEEN is normal in size and echogenicity.

BOTH KIDNEYS are normal in size and echogenicity. The central sinus complex is normal. No evidence of calculi/hydronephrosis. Right kidney measures 11.0×4.3 cm. Left kidney measures 10.4×6.1 cm.

PANCREAS: Head and body of pancreas is visualised and appears normal. Rest of the pancreas is obscured.

URINARY BLADDER is normal in capacity and contour. Few internal echoes are seen within. Bladder wall is normal in thickness. No evidence of intravesical calculi.

PROSTATE is normal in size & echogenicity. It measures ~ 18.2 cc in volume.

No evidence of ascites.

Impression:

- Grade I fatty infiltration of liver.
- Few internal echoes within urinary bladder, s/o cystitis. Recommended urine anlysis correlation.

DR. KUNAL NIGAM M.D. (Radiologist)