PID No. : MED121173651 Register On : 22/07/2023 2:27 PM : 522311814 Collection On : 22/07/2023 3:22 PM SID No. Age / Sex : 57 Year(s) / Male Report On : 24/07/2023 2:58 PM

Type : OP

: 10/10/2023 2:39 PM Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
BLOOD GROUPING AND Rh	'O' 'Positive'		

TYPING

 $({\rm EDTA~Blood} Agglutination)$

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.

Printed On

Complete Blood Count With - ESR

Haemoglobin (EDTA Blood/Spectrophotometry)	15.1	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	43.1	%	42 - 52
RBC Count (EDTA Blood)	4.92	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	87.7	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	30.7	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	35.0	g/dL	32 - 36
RDW-CV	13.1	%	11.5 - 16.0
RDW-SD	40.21	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	8400	cells/cu.m m	4000 - 11000
Neutrophils (Blood)	50.3	%	40 - 75
Lymphocytes (Blood)	36.1	%	20 - 45
Eosinophils (Blood)	5.3	%	01 - 06





APPROVED BY

The results pertain to sample tested.

Page 1 of 7

 PID No.
 : MED121173651
 Register On
 : 22/07/2023 2:27 PM

 SID No.
 : 522311814
 Collection On
 : 22/07/2023 3:22 PM

 Age / Sex
 : 57 Year(s) / Male
 Report On
 : 24/07/2023 2:58 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood)	7.4	%	01 - 10
Basophils (Blood)	0.9	%	00 - 02
INTERPRETATION: Tests done on Automated Fi	ive Part cell counter. Al	l abnormal results are r	eviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	4.23	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	3.03	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.45	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.62	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.08	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	260	10^3 / μl	150 - 450
MPV (Blood)	6.9	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.18	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	3	mm/hr	< 20
BUN / Creatinine Ratio	13.7		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	200.43	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 10/10/2023 2:39 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.





The results pertain to sample tested.

Page 2 of 7

Lab Address: BANGALORE REFERENCE LABORATORY-CLUMAX DIAGNOSTIC AND RESEARCH CENTRE PVT LTD. (A UNIT OF MEDALL HEALTHCARE PVT. LTD.),Old No 66 & New No 1, 2nd Main Road, Bashyam Circle, Sankey Cross Road Malleswaram, Gayathri Devi Park Extension, Kodandarampura, Sadashiva Nagar, Bengaluru, Karnataka-560003,.

 PID No.
 : MED121173651
 Register On
 : 22/07/2023 2:27 PM

 SID No.
 : 522311814
 Collection On
 : 22/07/2023 3:22 PM

 Age / Sex
 : 57 Year(s) / Male
 Report On
 : 24/07/2023 2:58 PM

Type : OP Printed On : 10/10/2023 2:39 PM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
Glucose Postprandial (PPBS)	189.79	mg/dL	70 - 140
(Plasma - PP/GOD-PAP)			

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	10.4	mg/dL	7.0 - 21
(Serum/Urease UV / derived)			
Creatinine	0.76	mg/dL	0.9 - 1.3
(Serum/Modified Jaffe)			

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc

Uric Acid (Serum/Enzymatic) Liver Function Test	3.34	mg/dL	3.5 - 7.2
Bilirubin(Total) (Serum/DCA with ATCS)	0.46	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.16	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.30	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	20.45	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	21.04	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	26.79	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	105.1	U/L	56 - 119





APPROVED BY

The results pertain to sample tested.

Page 3 of 7

Lab Address: BANGALORE REFERENCE LABORATORY-CLUMAX DIAGNOSTIC AND RESEARCH CENTRE PVT LTD. (A UNIT OF MEDALL HEALTHCARE PVT. LTD.),Old No 66 & New No 1, 2nd Main Road, Bashyam Circle, Sankey Cross Road Malleswaram, Gayathri Devi Park Extension, Kodandarampura, Sadashiva Nagar, Bengaluru, Karnataka-560003,.

 PID No.
 : MED121173651
 Register On
 : 22/07/2023 2:27 PM

 SID No.
 : 522311814
 Collection On
 : 22/07/2023 3:22 PM

 Age / Sex
 : 57 Year(s) / Male
 Report On
 : 24/07/2023 2:58 PM

 Age / Sex
 : 57 Year(s) / Male
 Report On
 : 24/07/2023 2:58 PM

 Type
 : OP
 Printed On
 : 10/10/2023 2:39 PM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
Total Protein (Serum/Biuret)	6.83	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.33	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.50	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.73		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	119.28	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	253.01	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	34.02	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	34.7	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189

Very High: >= 190





APPROVED BY

PID No. : MED121173651 Register On : 22/07/2023 2:27 PM : 522311814 SID No. Collection On : 22/07/2023 3:22 PM Age / Sex : 57 Year(s) / Male Report On : 24/07/2023 2:58 PM

Type : OP **Printed On** : 10/10/2023 2:39 PM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
VLDL Cholesterol (Serum/Calculated)	50.6	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	85.3	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	3.5	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	7.4	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	1	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

Glycosylated Haemoglobin (HbA1c)

% HbA1C 9.0 Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 (Whole Blood/HPLC) Diabetic: \geq 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 211.6 mg/dL

(Whole Blood)





APPROVED BY

SID No. : 522311814 Collection On : 22/07/2023 3:22 PM

Age / Sex: 57 Year(s) / Male **Report On**: 24/07/2023 2:58 PM

Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

: 10/10/2023 2:39 PM

ng/ml

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values. Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

Prostate specific antigen - Total(PSA)

0.562

Normal: 0.0 - 4.0

Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0

Suspicious of Malignant disease of

Prostate: > 10.0

THYROID PROFILE / TFT

(Serum/Manometric method)

T3 (Triiodothyronine) - Total 1.15 ng/ml 0.4 - 1.81

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 8.77 µg/dl 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.70 μ IU/mL 0.35 - 5.50

(Serum/ECLIA)





APPROVED BY

Printed On

Age / Sex: 57 Year(s) / Male **Report On**: 24/07/2023 2:58 PM

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u>
Value Reference Interval

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

: 10/10/2023 2:39 PM

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.





APPROVED BY

-- End of Report --

Name	SIDDALINGAIAH	ID	MED121173651
Age & Gender	57-Male		7/24/2023 2:58:57 PM
Ref Doctor Name	MediWheel		



ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (12.8 cm) and shows increased echogenicity. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER - Postprandial status.

CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

BOTH KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

•	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	11.1	1.6
Left Kidney	10.8	1.7

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern. It measures 3.1 x 2.7 x 3.7 cms and vol: 15.9cc.

No evidence of ascites.

IMPRESSION:

- Grade I fatty infiltration of liver.
- No other significant abnormality detected.

- 1.This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any, with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

Name	SIDDALINGAIAH	ID	MED121173651
Age & Gender	57-Male	Visit Date	7/24/2023 2:58:57 PM
Ref Doctor Name	MediWheel		



DR. HEMANANDINI V. N. CONSULTANT RADIOLOGIST Hn/mj

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3.Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- O.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false oninion
- $11. Disputes, if any\ , with regard\ to\ the\ report\ findings\ are\ subject\ to\ the\ exclusive\ jurisdiction\ of\ the\ competent\ courts\ chennai\ only.$

Name	SIDDALINGAIAH	ID	MED121173651
Age & Gender	57-57-Male	Visit Date	7/24/2023 2:58:57 PM
Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHIC STUDY

M-mode measurement:

AORTA 2.62 cms. LEFT ATRIUM 2.84 cms. **AVS** 1.47 cms. LEFT VENTRICLE (DIASTOLE) 4.29 cms. (SYSTOLE) 2.43 cms. **VENTRICULAR SEPTUM** 0.96 (DIASTOLE) cms. (SYSTOLE) 1.35 cms. **POSTERIOR WALL** (DIASTOLE) 1.08 cms. (SYSTOLE) 1.40 cms. **EDV** 62 ml. **ESV** 20 ml. FRACTIONAL SHORTENING 43 % **EJECTION FRACTION** 60 %

DOPPLER MEASUREMENTS:

EPSS

RVID

MITRAL VALVE: E - 0.4 m/s A - 0.6 m/s NO MR.

AORTIC VALVE: 1.1 m/s NO AR.

TRICUSPID VALVE: E - 0.2 m/s A - 0.3 m/s NO TR.

PULMONARY VALVE: 0.8 m/s NO PR.

REPORT DISCLAIMER

cms.

cms.

1.80

- 1.This is only a radiological imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any, with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chemai only.

Name	SIDDALINGAIAH	ID	MED121173651
Age & Gender	57-57-Male	Visit Date	7/24/2023 2:58:57 PM
Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHY FINDINGS:

Left Ventricle : Normal size, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal.Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION: FAIR ECHO WINDOW

- NORMAL SIZED CARDIAC CHAMBERS.
- GRADE I LV DIASTOLIC DYSFUNCTION.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. YASHODA RAVI

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3.Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any, with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

Name	SIDDALINGAIAH	ID	MED121173651
Age & Gender	57-57-Male		7/24/2023 2:58:57 PM
Ref Doctor Name	MediWheel		



CONSULTANT CARDIOLOGIST

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11. Disputes, if any, with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr. SIDDALINGAIAH	ID	MED121173651
Age & Gender	57Y/M	Visit Date	Jul 22 2023 2:27PM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Aortic unfolding is noted.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

• No significant abnormality detected.

DR.HEMANANDHINI CONSULTANT RADIOLOGIST