PID No.
 : MED111391697
 Register On
 : 24/11/2022 9:16 AM

 SID No.
 : 222020132
 Collection On
 : 24/11/2022 10:20 AI

 SID No.
 : 222020132
 Collection On : 24/11/2022 10:20 AM

 Age / Sex : 31 Year(s) / Male
 Report On : 24/11/2022 5:43 PM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING	'O' 'Positive'		
(EDTA Blood/Agglutination)			
INTERPRETATION: Reconfirm the Blood group	p and Typing before	blood transfusion	
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	14.9	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	45.4	%	42 - 52
RBC Count (EDTA Blood/Impedance Variation)	4.92	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	92.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	30.3	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	32.8	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	13.6	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	44.7	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	6600	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	56.7	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	32.5	%	20 - 45
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	2.3	%	01 - 06



 PID No.
 : MED111391697
 Register On
 : 24/11/2022 9:16 AM

 SID No.
 : 222020132
 Collection On
 : 24/11/2022 10:20 AM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	7.6	%	01 - 10
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.9	%	00 - 02
INTERPRETATION: Tests done on Automated	l Five Part cell count	ter. All abnormal results a	are reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.74	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.15	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.15	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.50	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.06	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	292	$10^3 / \mu l$	150 - 450
MPV (EDTA Blood/Derived from Impedance)	7.5	fL	7.9 - 13.7
PCT (EDTA Blood/Automated Blood cell Counter)	0.218	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	14	mm/hr	< 15
BUN / Creatinine Ratio	10.3		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	89.5	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126



Age / Sex : 31 Year(s) / Male

 PID No.
 : MED111391697
 Register On
 : 24/11/2022 9:16 AM

 SID No.
 : 222020132
 Collection On
 : 24/11/2022 10:20 AM

Report On



Type : OP Printed On : 05/12/2022 5:28 PM

Ref. Dr : MediWheel

Investigation	Observed Unit	<u>Biological</u>
-	Value	Reference Interval

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

: 24/11/2022 5:43 PM

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	124.2	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	9.4	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.91	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine etc.

Cic.			
Uric Acid (Serum/Enzymatic)	6.0	mg/dL	3.5 - 7.2
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.94	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.29	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.65	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	28.2	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase)	35.5	U/L	5 - 41



APPROVED BY

The results pertain to sample tested.

(Serum/Modified IFCC)

Page 3 of 7

PID No. : MED111391697 Register On : 24/11/2022 9:16 AM

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	34.5	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	127.7	U/L	53 - 128
Total Protein (Serum/Biuret)	7.79	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.39	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	3.40	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.29		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	202.7	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	138.8	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol 43.6 mg/dL Optimal(Negative Risk Factor): >= (Serum/Immunoinhibition) 60

Borderline: 40 - 59 High Risk: < 40



Age / Sex : 31 Year(s) / Male **Report On** : 24/11/2022 5:43 PM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
LDL Cholesterol (Serum/Calculated)	131.3	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	27.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	159.1	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	4.6	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	3.2	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	3	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
Glycosylated Haemoglobin (HbA1c)		
HbA1C (Whole Blood/HPLC)	5.4 %	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %



Age / Sex : 31 Year(s) / Male **Report On** : 24/11/2022 5:43 PM

Type : OP Printed On : 05/12/2022 5:28 PM

Ref. Dr : MediWheel

Investigation

Observed Value

Unit
Value

Biological
Reference Interval

Estimated Average Glucose

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.08 ng/ml 0.7 - 2.04

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 9.70 $\mu g/dl$ 4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.42 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



APPROVED BY

The results pertain to sample tested.

Page 6 of 7

 SID No.
 : 222020132
 Collection On : 24/11/2022 10:20 AM

 Age / Sex
 : 31 Year(s) / Male
 Report On : 24/11/2022 5:43 PM

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u> Urine Analysis - Routine</u>			
COLOUR (Urine)	Pale yellow		Yellow to Amber
APPEARANCE (Urine)	Clear		Clear
Protein (Urine/Protein error of indicator)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Pus Cells (Urine/Automated 6"Flow cytometry)	Occasional	/hpf	NIL
Epithelial Cells (Urine/Automated 6"Flow cytometry)	Occasional	/hpf	NIL
RBCs (Urine/Automated 6"Flow cytometry)	NIL	/hpf	NIL
Casts (Urine/Automated 6"Flow cytometry)	NIL	/hpf	NIL
Crystals (Urine/Automated 6"Flow cytometry)	NIL	/hpf	NIL
Others (Urine)	NIL		

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.



APPROVED BY

-- End of Report --