

Units

#### PATIENT NAME: NITHYA K N REF. DOCTOR : DR. BANK OF BARODA CODE/NAME & ADDRESS : C000138396 ACCESSION NO : 0183WD000644 AGE/SEX :29 Years Female ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) :07/04/2023 00:00:00 PATIENT ID : NITHF020793183 DRAWN F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED :07/04/2023 17:33:08 DELHI ABHA NO REPORTED :24/04/2023 10:50:37 : NEW DELHI 110030 8800465156 **Test Report Status** Biological Reference Interval

Results

# MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

<u>Final</u>

XRAY-CHEST				
»»	BOTH THE LUNG FIELDS AF	RE CLEAR		
»»	BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGELS ARE CLEAR			
»»	BOTH THE HILA ARE NORMAL			
»»	CARDIAC AND AORTIC SHADOWS APPEAR NORMAL			
»»	BOTH THE DOMES OF THE	DIAPHRAM ARE NORMAL		
»»	VISUALIZED BONY THORAX	X IS NORMAL		
IMPRESSION	NO ABNORMALITY DETECT	ED		
ECG				
ECG	WITHIN NORMAL LIMITS			
MEDICAL HISTORY				
RELEVANT PRESENT HISTORY	NOT SIGNIFICANT			
RELEVANT PAST HISTORY	NOT SIGNIFICANT			
RELEVANT PERSONAL HISTORY	UNMARRIED			
RELEVANT FAMILY HISTORY	FATHER H/O HTN			
OCCUPATIONAL HISTORY	NOT SIGNIFICANT			
HISTORY OF MEDICATIONS	NOT SIGNIFICANT			
ANTHROPOMETRIC DATA & BMI				
HEIGHT IN METERS	1.52		mts	
WEIGHT IN KGS.	53		Kgs	
BMI	23	BMI & Weight Status as foll Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight	o <b>∖kg</b> ∕sqmts	

### **GENERAL EXAMINATION**

MENTAL / EMOTIONAL STATE	NORMAL
PHYSICAL ATTITUDE	NORMAL
GENERAL APPEARANCE / NUTRITIONAL STATUS	HEALTHY
BUILT / SKELETAL FRAMEWORK	AVERAGE
FACIAL APPEARANCE	NORMAL
SKIN	NORMAL

**Dr.Karthick Prabhu R Consultant Pathologist** 







Vie<u>w Report</u>

30.0 and Above: Obese



Test Report Status

<u>Final</u>



**Biological Reference Interval** Units

PATIENT NAME : NITHYA K N	<b>REF. DOCTOR</b> : D	DR. BANK OF BARODA
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	ACCESSION NO: <b>0183WD000644</b> PATIENT ID : NITHF020793183 CLIENT PATIENT ID: ABHA NO :	AGE/SEX: 29 YearsFemaleDRAWN: 07/04/202300:00:00RECEIVED: 07/04/202317:33:08REPORTED: 24/04/202310:50:37

Results

	-	
UPPER LIMB	NORMAL	
LOWER LIMB	NORMAL	
NECK	NORMAL	
NECK LYMPHATICS / SALIVARY GLANDS	NOT ENLARGED OR TENDER	
THYROID GLAND	NOT ENLARGED	
CAROTID PULSATION	NORMAL	
BREAST (FOR FEMALES)	NORMAL	
TEMPERATURE	NORMAL	
PULSE	74/MINS, REGULAR, ALL PERIPHERAL PULSES WELL F BRUIT	ELT, NO CAROTID
RESPIRATORY RATE	NORMAL	
CARDIOVASCULAR SYSTEM		
ВР	120/80 MM HG (SITTING)	mm/Hg
PERICARDIUM	NORMAL	
APEX BEAT	NORMAL	
HEART SOUNDS	NORMAL	
MURMURS	ABSENT	
RESPIRATORY SYSTEM		
SIZE AND SHAPE OF CHEST	NORMAL	
MOVEMENTS OF CHEST	SYMMETRICAL	
BREATH SOUNDS INTENSITY	NORMAL	
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)	
ADDED SOUNDS	ABSENT	
PER ABDOMEN		
APPEARANCE	NORMAL	
VENOUS PROMINENCE	ABSENT	
LIVER	NOT PALPABLE	
SPLEEN	NOT PALPABLE	
HERNIA	ABSENT	
CENTRAL NERVOUS SYSTEM		
HIGHER FUNCTIONS	NORMAL	
CRANIAL NERVES	NORMAL	



Dr.Karthick Prabhu R Consultant Pathologist

Page 2 Of 22









PATIENT NAME: NITHYA K N		<b>REF. DOCTOR :</b> D	R. BANK O	BARODA	
CODE/NAME & ADDRESS : C000138396	ACCESSION NO	: 0183WD000644	AGE/SEX	:29 Years	Female
	PATIENT ID	: NITHF020793183	DRAWN	:07/04/2023	00:00:00
F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	CLIENT PATIENT	TID:	RECEIVED	:07/04/2023	17:33:08
NEW DELHI 110030	ABHA NO	:	REPORTED	:24/04/2023	10:50:37
8800465156					
	<u> </u>				
Test Report Status <u>Final</u>	Results	Biological	Reference	e Interval 🛛 🛛	Jnits

CEREBELLAR FUNCTIONS	NORMAL
SENSORY SYSTEM	NORMAL
MOTOR SYSTEM	NORMAL
REFLEXES	NORMAL
MUSCULOSKELETAL SYSTEM	
SPINE	NORMAL
JOINTS	NORMAL
BASIC EYE EXAMINATION	
CONJUNCTIVA	NORMAL
EYELIDS	NORMAL
EYE MOVEMENTS	NORMAL
CORNEA	NORMAL
DISTANT VISION RIGHT EYE WITH GLASSES	REFRACTIVE ERROR NEED CORRECTION
DISTANT VISION LEFT EYE WITH GLASSES	REFRACTIVE ERROR NEED CORRECTION
NEAR VISION RIGHT EYE WITH GLASSES	REFRACTIVE ERROR NEED CORRECTION
NEAR VISION LEFT EYE WITH GLASSES	REFRACTIVE ERROR NEED CORRECTION
COLOUR VISION	NORMAL
BASIC ENT EXAMINATION	
EXTERNAL EAR CANAL	NORMAL
TYMPANIC MEMBRANE	NORMAL
NOSE	NO ABNORMALITY DETECTED
SINUSES	NORMAL
THROAT	NO ABNORMALITY DETECTED
TONSILS	NOT ENLARGED
BASIC DENTAL EXAMINATION	
TEETH	NORMAL
GUMS	HEALTHY
SUMMARY	
RELEVANT HISTORY	NOT SIGNIFICANT
RELEVANT GP EXAMINATION FINDINGS	NOT SIGNIFICANT
RELEVANT LAB INVESTIGATIONS	WITHIN NORMAL LIMITS
RELEVANT NON PATHOLOGY DIAGNOSTICS	NO ABNORMALITIES DETECTED

Dr.Karthick Prabhu R Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in







View Report





**Biological Reference Interval** Units

PATIENT NAME: NITHYA K N	<b>REF. DOCTOR</b> : D	R. BANK OF BARODA
	ACCESSION NO : 0183WD000644	AGE/SEX : 29 Years Female
	PATIENT ID : NITHF020793183	DRAWN :07/04/2023 00:00:00
F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	CLIENT PATIENT ID:	RECEIVED :07/04/2023 17:33:08
NEW DELHI 110030	ABHA NO :	REPORTED :24/04/2023 10:50:37
8800465156		
· · · · · · · · · · · · · · · · · · ·		

**REMARKS / RECOMMENDATIONS** 

**Final** 

NONE

Results

### FITNESS STATUS

**Test Report Status** 

FITNESS STATUS

FIT (AS PER REQUESTED PANEL OF TESTS)

#### Comments

OUR PANEL OF DOCTORS :

GENERAL PHYSICIANS - DR.S.B.PRAVEEN., M.B.B.S., M.Sc(Psy)., F.Diab., AFIH. RADIOLOGIST - DR.DEBABRATA NITYARANJAN DAS, MD(RAD)., M.R.FELLOW(USA)., GYNECOLOGIST - DR.PREMALATHA KRISHNAKUMAR.MD.,MRCOG.,Dip.in Colposcopy(UK). CARDIOLOGIST - DR. A.PREM KRISHNA, MD., MRCP(UK)., DNB., DM. THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY HEAD THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

#### Interpretation(s)

MEDICAL

HISTORY-\*\*\*\*\* THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

FITNESS STATUS-Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history as well as the comprehensiveness of the diagnostic panel which has been requested for . These are then further correlated with details of the job under consideration to eventually fit the right man to the right job.

Basis the above, SRL classifies a candidate's Fitness Status into one of the following categories:

• Fit (As per requested panel of tests) - SRL Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.

• Fit (with medical advice) (As per requested panel of tests) - This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician<sup>111</sup>'s consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job. • Fitness on Hold (Temporary Unfit) (As per requested panel of tests) - Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit

(With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.

Unfit (As per requested panel of tests) - An unfit report by SRL Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.



**Dr.Karthick Prabhu R** Consultant Pathologist







Details





PATIENT NAME : NITHYA K N	<b>REF. DOCTOR :</b>	DR. BANK OF BARODA
	ACCESSION NO : 0183WD000644	AGE/SEX : 29 Years Female
F-703, F-703, LADO SARAT, MEHRAULISOUTH WEST	PATIENT ID : NITHF020793183	DRAWN :07/04/2023 00:00:00
DELHI	CLIENT PATIENT ID:	RECEIVED :07/04/2023 17:33:08
NEW DELHI 110030 8800465156	ABHA NO :	REPORTED :24/04/2023 10:50:37
0000405150		
Test Report Status <u>Final</u>	Results Biological	Reference Interval Units

HAEMATOLOGY - CBC			
MEDI WHEEL FULL BODY HEALTH CHECKUP BEI	OW 40FEMALE		
BLOOD COUNTS, EDTA WHOLE BLOOD			
HEMOGLOBIN (HB)	12.6	12.0 - 15.0	g/dL
RED BLOOD CELL (RBC) COUNT	4.68	3.8 - 4.8	mil/µL
WHITE BLOOD CELL (WBC) COUNT	7.70	4.0 - 10.0	thou/µL
PLATELET COUNT	266	150 - 410	thou/µL
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV)	37.3	36 - 46	%
MEAN CORPUSCULAR VOLUME (MCV)	80.0 Low	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	27.0	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	33.9	31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW)	13.4	11.6 - 14.0	%
MENTZER INDEX	17.1		
MEAN PLATELET VOLUME (MPV)	7.6	6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT			
NEUTROPHILS	59	40 - 80	%
LYMPHOCYTES	30	20 - 40	%
MONOCYTES	5	2 - 10	%
EOSINOPHILS	5	1 - 6	%
BASOPHILS	1	< 1 - 2	%
ABSOLUTE NEUTROPHIL COUNT	4.54	2.0 - 7.0	thou/µL
ABSOLUTE LYMPHOCYTE COUNT	2.31	1.0 - 3.0	thou/µL
ABSOLUTE MONOCYTE COUNT	0.38	0.2 - 1.0	thou/µL
ABSOLUTE EOSINOPHIL COUNT	0.38	0.02 - 0.50	thou/µL
ABSOLUTE BASOPHIL COUNT	0.08	0.02 - 0.10	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	2.0		

#### Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

Dr.Karthick Prabhu R Consultant Pathologist





View Report

Page 5 Of 22

View Details





PATIENT NAME: NITHYA K N		REF. DOCTOR : D	R. BANK OF	BARODA	
CODE/NAME & ADDRESS : C000138396	ACCESSION NO	: 0183WD000644	AGE/SEX	:29 Years	Female
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	PATIENT ID	: NITHF020793183	DRAWN	:07/04/2023	00:00:00
DELAI	CLIENT PATIEN ABHA NO	TID:		: 07/04/2023 :24/04/2023	
8800465156					
Test Report Status Final	Results	Biological	Reference	e Interval L	Inits

diagnosing a case of beta thalassaemia trait. WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.3, COVID-19 patients tend to show mild disease. (Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.



**Dr.Karthick Prabhu R Consultant Pathologist** 

**PERFORMED AT :** Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in

Page 6 Of 22





V<u>iew Details</u>





PATIENT NAME: NITHYA K N		<b>REF. DOCTOR :</b> D	R. BANK O	F BARODA	
CODE/NAME & ADDRESS : C000138396	ACCESSION NO	: 0183WD000644	AGE/SEX	:29 Years	Female
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	PATIENT ID	: NITHF020793183	DRAWN	:07/04/2023	00:00:00
DELAI	CLIENT PATIEN	ΓID:		:07/04/2023	
NEW DELHI 110030	ABHA NO	:	REPORTED	:24/04/2023	10:50:37
8800465156					
Test Report Status <u>Final</u>	Results	Biological	Reference	e Interval U	Inits

	HAEMATOLOGY				
MEDI WHEEL FULL BODY HEALTH	MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE				
ERYTHROCYTE SEDIMENTATION BLOOD	RATE (ESR),WHOLE				
E.S.R	24 High	0 - 20	mm at 1 hr		

#### Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-**TEST DESCRIPTION** :-Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy,

Estrogen medications, vasculities, inhammatory artificis, kenal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy fissue injury, Pregnancy, Estrogen medication, Aging. Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythermia vera, Sickle cell anemia

#### LIMITATIONS

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia False Decreased : Poikilocytosis, (Sickle Cells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine,

salicylates)

**REFERENCE** :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.



**Dr.Karthick Prabhu R Consultant Pathologist** 



View Report

Page 7 Of 22

**PERFORMED AT :** Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in

Details





PATIENT NAME: NITHYA K N	REF. DOCTOR	DR. BANK OF BARODA
	ACCESSION NO : <b>0183WD000644</b> PATIENT ID : NITHF020793183	AGE/SEX :29 Years Female DRAWN :07/04/2023 00:00:00
DELHI	CLIENT PATIENT ID: ABHA NO :	RECEIVED :07/04/2023 17:33:08 REPORTED :24/04/2023 10:50:37
8800465156	Results Biologic	al Reference Interval Units

	IMMUNOHAEMATOLOGY	
MEDI WHEEL FULL BODY HEAL	1 CHECKUP BELOW 40FEMALE	
ABO GROUP & RH TYPE, EDTA \	HOLE BLOOD	
ABO GROUP	TYPE B	
RH TYPE	POSITIVE	

Interpretation(s) ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.



**Dr.Karthick Prabhu R Consultant Pathologist** 













PATIENT NAME : NITHYA K N	REF. DOCTOR :	DR. BANK OF BARODA
CODE/NAME & ADDRESS : C000138396 ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	ACCESSION NO : <b>0183WD000644</b> PATIENT ID : NITHF020793183	AGE/SEX : 29 Years Female DRAWN :07/04/2023 00:00:00
F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	CLIENT PATIENT ID:	RECEIVED :07/04/2023 17:33:08 REPORTED :24/04/2023 10:50:37
NEW DELHI 110030 8800465156		10.30.37
Test Report Status <u>Final</u>	Results Biological	Reference Interval Units

	BIOCHEMISTRY		
MEDI WHEEL FULL BODY HEALTH CHECKUP BEI	OW 40FEMALE		
GLUCOSE FASTING, FLUORIDE PLASMA			
FBS (FASTING BLOOD SUGAR) METHOD : HEXOKINASE / SPECTROPHOTOMETRY	80	74 - 99	mg/dL
GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA V BLOOD	VHOLE		
HBA1C	5.2	Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 ADA Target: 7.0 Action suggested: > 8.0	%
ESTIMATED AVERAGE GLUCOSE(EAG) GLUCOSE, POST-PRANDIAL, PLASMA	102.5	< 116.0	mg/dL
PPBS(POST PRANDIAL BLOOD SUGAR) METHOD : HEXOKINASE / SPECTROPHOTOMETRY	109	70 - 139	mg/dL
LIPID PROFILE, SERUM			
CHOLESTEROL, TOTAL	163	< 200 Desirable 200 - 239 Borderline High >/= 240 High	mg/dL
METHOD : CHOLESTEROL OXIDASE / SPECTROPHOTOMETRY		, 5	
TRIGLYCERIDES	49	< 150 Normal 150 - 199 Borderline High 200 - 499 High >/=500 Very High	mg/dL
HDL CHOLESTEROL	48	< 40 Low >/=60 High	mg/dL
CHOLESTEROL LDL	105 High	< 100 Optimal 100 - 129 Near optimal/ above optima 130 - 159 Borderline High 160 - 189 High >/= 190 Very High	mg/dL I



**Dr.Karthick Prabhu R Consultant Pathologist** 

**PERFORMED AT :** Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in









PATIENT NAME : NITHYA K N		REF. DOCTOR : D	R. BANK OF	BARODA	
CODE/NAME & ADDRESS : C000138396 ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : <b>018</b> PATIENT ID : NITH CLIENT PATIENT ID: ABHA NO :	IF020793183	RECEIVED :	29 Years 07/04/2023 07/04/2023 24/04/2023	17:33:08
Test Report Status <u>Final</u>	Results	Biological	Reference	Interval l	Jnits
NON HDL CHOLESTEROL	115	Desirable: Above Des Borderline High: 190 Very high:	irable: 130 High: 160 - 219	0 - 159 ) - 189	/dL
VERY LOW DENSITY LIPOPROTEIN	9.8	= 30.0</td <td></td> <td></td> <td>/dL</td>			/dL
CHOL/HDL RATIO	3.4	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Ri 7.1 - 11.0 Moderate F > 11.0 High Risk			
LDL/HDL RATIO	2.2	0.5 - 3.0 E 3.1 - 6.0 E Risk >6.0 High	orderline/		

### Interpretation(s)

1) Cholesterol levels help assess the patient risk status and to follow the progress of patient under treatment to lower serum cholesterol concentrations.

2) Serum Triglyceride (TG) are a type of fat and a major source of energy for the body. Both quantity and composition of the diet impact on plasma triglyceride concentrations. Elevations in TG levels are the result of overproduction and impaired clearance. High TG are associated with increased risk for CAD (Coronary artery disease) in patients with other risk factors, such as low HDL-C, some patient groups with elevated apolipoprotein B concentrations, and patients with forms of LDL that may be particularly atherogenic.

3)HDL-C plays a crucial role in the initial step of reverse cholesterol transport, this considered to be the primary atheroprotective function of HDL

4) LDL -C plays a key role in causing and influencing the progression of atherosclerosis and, in particular, coronary sclerosis. The majority of cholesterol stored in atherosclerotic plaques originates from LDL, thus LDL-C value is the most powerful clinical predictor.

5)Non HDL cholesterol: Non-HDL-C measures the cholesterol content of all atherogenic lipoproteins, including LDL hence it is a better marker of risk in both primary and secondary prevention studies. Non-HDL-C also covers, to some extent, the excess ASCVD risk imparted by the sdLDL, which is significantly more atherogenic than the normal large buoyant particles, an elevated non-HDL-C indirectly suggests greater proportion of the small, dense variety of LDL particles

Serum lipid profile is measured for cardiovascular risk prediction.Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

### Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Dr.Karthick Prabhu R Consultant Pathologist

Page 10 Of 22





View Report

View Details





#### PATIENT NAME: NITHYA K N REF. DOCTOR : DR. BANK OF BARODA CODE/NAME & ADDRESS : C000138396 ACCESSION NO : 0183WD000644 AGE/SEX :29 Years Female ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) PATIENT ID DRAWN :07/04/2023 00:00:00 : NITHF020793183 F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED :07/04/2023 17:33:08 DELHI ABHA NO REPORTED :24/04/2023 10:50:37 : NEW DELHI 110030 8800465156

## Test Report Status Final

Results

**Biological Reference Interval** Units

Risk Category			
Extreme risk group	A.CAD with $> 1$ feature of high risk group		
	B. CAD with $> 1$ feature of Very high risk §	B. CAD with > 1 feature of Very high risk group or recurrent ACS (within 1 year) despite LDL-C	
	< or $=$ 50 mg/dl or polyvascular disease		
Very High Risk		major risk factors or evidence of end organ damage 3.	
	Familial Homozygous Hypercholesterolemi	a	
High Risk		abetes with 1 major risk factor or no evidence of end	
		organ damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6.	
	Coronary Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid		
	plaque		
Moderate Risk	2 major ASCVD risk factors		
Low Risk	0-1 major ASCVD risk factors		
Major ASCVD (Ath	Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors		
1. Age $>$ or $=$ 45 year	ears in males and $>$ or $= 55$ years in females 3. Current Cigarette smoking or tobacco use		
2. Family history of p	remature ASCVD 4. High blood pressure		
5. Low HDL			

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
Category A	< OR = 30)	< OR = 60)		
Extreme Risk Group	<or 30<="" =="" td=""><td><or 60<="" =="" td=""><td>&gt; 30</td><td>&gt;60</td></or></td></or>	<or 60<="" =="" td=""><td>&gt; 30</td><td>&gt;60</td></or>	> 30	>60
Category B				
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR=70	>OR=100
Moderate Risk	<100	<130	>OR=100	>OR=130
Low Risk	<100	<130	>OR=130*	>OR=160

\*After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

## LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	0.80	0.2 - 1.0	mg/dL
METHOD : DIAZOTIZED SULFANILIC ACID / SPECTROPHOTOMETRY			
BILIRUBIN, DIRECT	0.10	0.0 - 0.2	mg/dL
METHOD : DIAZOTIZED SULFANILIC ACID / SPECTROPHOTOMETRY			
BILIRUBIN, INDIRECT	0.70	0.1 - 1.0	mg/dL
TOTAL PROTEIN	6.8	6.4 - 8.2	g/dL
ALBUMIN	4.1	3.4 - 5.0	g/dL

Dr.Karthick Prabhu R Consultant Pathologist



Page 11 Of 22



PATIENT NAME : NITHYA K N	REF. DOCTOR : D	DR. BANK OF BARODA
CODE/NAME & ADDRESS : C000138396 ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO: <b>0183WD000644</b> PATIENT ID : NITHF020793183 CLIENT PATIENT ID: ABHA NO :	AGE/SEX       :29 Years       Female         DRAWN       :07/04/2023       00:00:00         RECEIVED       :07/04/2023       17:33:08         REPORTED       :24/04/2023       10:50:37
Test Report Status <u>Final</u>	Results Biological	Reference Interval Units

METHOD : BCP DYE BINDING / SPECTOPHOTOMETER			
GLOBULIN	2.7	2.0 - 4.1	g/dL
ALBUMIN/GLOBULIN RATIO	1.5	1.0 - 2.1	RATIO
ASPARTATE AMINOTRANSFERASE(AST/SGOT) METHOD : UV WITH PYRIDOXAL 5 PHOSPHATE / SPECTROPHOTOMET	12 Low ER	15 - 37	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD : UV WITH PYRIDOXAL 5 PHOSPHATE / SPECTROPHOTOMET	<b>29</b> ER	< 34.0	U/L
ALKALINE PHOSPHATASE	29 Low	30 - 120	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD : GCNA / SPECTROPHOTOMETRY	27	5 - 55	U/L
LACTATE DEHYDROGENASE	98 Low	100 - 190	U/L
METHOD : LACTATE PYRUVATE UV/ L.LACTATE / SPECTOPHOTOMETER	R		
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	7	6 - 20	mg/dL
METHOD : UREASE / GLDH / SPECTROPHOTOMETRY			
CREATININE, SERUM			
CREATININE	0.49 Low	0.60 - 1.10	mg/dL
METHOD : PICRATE/ JAFFE / SPECTOPHOTOMETER			
BUN/CREAT RATIO			
BUN/CREAT RATIO	14.29	5.00 - 15.00	
URIC ACID, SERUM			
URIC ACID	2.7	2.6 - 6.0	mg/dL
METHOD : URICASE / CATALASE UV / SPECTROPHOTOMETRY			
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	6.8	6.4 - 8.2	g/dL
ALBUMIN, SERUM			
ALBUMIN	4.1	3.4 - 5.0	g/dL
METHOD : BCP DYE BINDING / SPECTOPHOTOMETER			
GLOBULIN			
GLOBULIN	2.7	2.0 - 4.1	g/dL
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM, SERUM	138.4	136 - 145	mmol/L
POTASSIUM, SERUM	4.47	3.50 - 5.10	mmol/L
CHLORIDE, SERUM	107.2 High	98 - 107	mmol/L
	-		



Dr.Karthick Prabhu R Consultant Pathologist



3

View Details View Report





PATIENT NAME: NITHYA K N	REF. DOCTOR : D	R. BANK OF BARODA
F-703, F-703, LADO SARAL MEHRAULISOUTH WEST	ACCESSION NO : <b>0183WD000644</b> PATIENT ID : NITHF020793183 CLIENT PATIENT ID: ABHA NO :	AGE/SEX :29 Years Female DRAWN :07/04/2023 00:00:00 RECEIVED :07/04/2023 17:33:08 REPORTED :24/04/2023 10:50:37
Test Report Status Final	Results Biological	Reference Interval Units

#### Comments

NOTE : SERUM ELECTROLYTE VALUES RECHECKED AND CONFIRMED Interpretation(s)

Sodium	Potassium	Chloride
Decreased in:CCF, cirrhosis,	Decreased in: Low potassium	Decreased in: Vomiting, diarrhea,
vomiting, diarrhea, excessive	intake,prolonged vomiting or diarrhea,	renal failure combined with salt
sweating, salt-losing	RTA types I and II,	deprivation, over-treatment with
nephropathy,adrenal insufficiency,	hyperaldosteronism, Cushing's	diuretics, chronic respiratory acidosis,
nephrotic syndrome, water	syndrome,osmotic diuresis (e.g.,	diabetic ketoacidosis, excessive
intoxication, SIADH. Drugs:	hyperglycemia),alkalosis, familial	sweating, SIADH, salt-losing
thiazides, diuretics, ACE inhibitors,	periodic paralysis,trauma	nephropathy, porphyria, expansion of
chlorpropamide,carbamazepine,anti	(transient).Drugs: Adrenergic agents,	extracellular fluid volume,
depressants (SSRI), antipsychotics.	diuretics.	adrenalinsufficiency,
		hyperaldosteronism, metabolic
		alkalosis. Drugs: chronic
		laxative,corticosteroids, diuretics.
Increased in: Dehydration	Increased in: Massive hemolysis,	Increased in: Renal failure, nephrotic
(excessivesweating, severe	severe tissue damage, rhabdomyolysis,	syndrome, RTA, dehydration,
vomiting or diarrhea),diabetes	acidosis, dehydration,renal failure,	overtreatment with
mellitus, diabetesinsipidus,	Addison's disease, RTA type IV,	saline, hyperparathyroidism, diabetes
hyperaldosteronism, inadequate	hyperkalemic familial periodic	insipidus, metabolic acidosis from
water intake. Drugs: steroids,	paralysis. Drugs: potassium salts,	diarrhea (Loss of HCO3-), respiratory
licorice, oral contraceptives.	potassium- sparing diuretics,NSAIDs,	alkalosis, hyperadrenocorticism.
	beta-blockers, ACE inhibitors, high-	Drugs: acetazolamide, and rogens,
	dose trimethoprim-sulfamethoxazole.	hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or	Interferences: Hemolysis of sample,	Interferences: Test is helpful in
hyperproteinemi, if sodium analysis	delayed separation of serum,	assessing normal and increased anion
involves a dilution step can cause	prolonged fist clenching during blood	gap metabolic acidosis and in
spurious results. The serum sodium	drawing, and prolonged tourniquet	distinguishing hypercalcemia due to
falls about 1.6 mEq/L for each 100	placement. Very high WBC/PLT counts	hyperparathyroidism (high serum
mg/dL increase in blood glucose.	may cause spurious. Plasma potassium	chloride) from that due to malignancy
	levels are normal.	(Normal serum chloride)

Interpretation(s) GLUCOSE FASTING,FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine.

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides. Decreased in :Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease,

malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol

sulfonylureas, tolbutamide, and other oral hypoglycemic agents. NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic

index & response to food consumed,Alimentary Hypoglycemia,Increased insulin response & sensitivity etc. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.

Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for



**Dr.Karthick Prabhu R Consultant Pathologist** 



Page 13 Of 22

View Report

view Details





PATIENT NAME: NITHYA K N		REF. DOCTOR : D	R. BANK OI	BARODA	
CODE/NAME & ADDRESS : C000138396	ACCESSION NC	: 0183WD000644	AGE/SEX	:29 Years	Female
	PATIENT ID	: NITHF020793183	DRAWN	:07/04/2023	00:00:00
DELHI	CLIENT PATIEN		i	:07/04/2023	
NEW DELHI 110030	ABHA NO	:	REPORTED	:24/04/2023	10:50:37
8800465156					
Test Report Status Final	Results	Biological	Reference	Interval U	Inits

well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

1. eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.

2. eAG gives an evaluation of blood glucose levels for the last couple of months. 3. eAG is calculated as eAG (mg/dl) = 28.7 \* HbA1c - 46.7

#### HbA1c Estimation can get affected due to :

1. Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

2.Vitamin C & E are reported to falsely lower test results.(possibly by inhibiting glycation of hemoglobin.

3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.

4. Interference of hemoglobinopathies in HbA1c estimation is seen in

a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
 b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c) HDF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

**Bilirubin** is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. **Elevated levels** results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin. **AST** is no enzyme found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured

**AST** is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver,liver cancer,kidney failure,hemolytic anemia,pancreatitis,hemochromatosis. AST levels may also increase after a heart attack or strenuous activity.ALT test measures the amount of this enzyme in the blood.ALT is found mainly in the liver, but also in smaller amounts in the kidneys,heart,muscles, and pancreas.It is commonly measured as a part of a diagnostic evaluation of hepatotcellular injury, to determine liver health.AST levels increase during acute hepatitis,sometimes due to a viral infection, ischemia to the liver, chronic hepatotcitis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

**GGT** is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

**Total Protein** also known as total protein, is a biochemical test for measuring the total amount of protein in serum.Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to:Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular percentility or decreased lymphatic clearance majurition and wasting etc.

permeability or decreased lymphatic clearance,malnutrition and wasting etc BLOOD UREA NITROGEN (BUN), SERUM-**Causes of Increased** levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

 Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:• Myasthenia Gravis, Muscuophy URIC ACID, SERUM-Causes of Increased levels-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic syndrome Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome. Protein-losing enteropathy etc.

ALBUMIN, SERUM-

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns,



Dr.Karthick Prabhu R Consultant Pathologist



Page 14 Of 22

View Details View Report





PATIENT NAME: NITHYA K N	<b>REF. DOCTOR :</b>	DR. BANK OF BARODA
CODE/NAME & ADDRESS : C000138396	ACCESSION NO : 0183WD000644	AGE/SEX : 29 Years Female
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	PATIENT ID : NITHF020793183	DRAWN :07/04/2023 00:00:00
F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	CLIENT PATIENT ID:	RECEIVED :07/04/2023 17:33:08
NEW DELHI 110030	ABHA NO :	REPORTED :24/04/2023 10:50:37
8800465156		
		<u> </u>
Test Report Status Final	Results Biological	Reference Interval Units

hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.



Dr.Karthick Prabhu R Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in Page 15 Of 22





View Report





PATIENT NAME: NITHYA K N	REF	DOCTOR : D	R. BANK OF	F BARODA	
	ACCESSION NO : 0183WDC	00644	AGE/SEX	:29 Years	Female
	PATIENT ID : NITHF0207	793183	DRAWN	:07/04/2023	00:00:00
F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	CLIENT PATIENT ID:		RECEIVED	:07/04/2023	17:33:08
NEW DELHI 110030	ABHA NO :		REPORTED	:24/04/2023	10:50:37
8800465156					

Test Report Status Final

Results

**Biological Reference Interval** Units

CLINI	CAL PATH - URINALYSIS		)
MEDI WHEEL FULL BODY HEALTH CHECKUP BE	LOW 40FEMALE		
PHYSICAL EXAMINATION, URINE			
COLOR	PALE YELLOW		
APPEARANCE	CLOUDY		
CHEMICAL EXAMINATION, URINE			
PH	5.5	4.7 - 7.5	
SPECIFIC GRAVITY	1.025	1.003 - 1.035	
PROTEIN	NOT DETECTED	NOT DETECTED	
GLUCOSE	NOT DETECTED	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
BLOOD	NOT DETECTED	NOT DETECTED	
BILIRUBIN	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
NITRITE	NOT DETECTED	NOT DETECTED	
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED	
MICROSCOPIC EXAMINATION, URINE			
RED BLOOD CELLS	1 - 2	NOT DETECTED	/HPF
PUS CELL (WBC'S)	10-15	0-5	/HPF
EPITHELIAL CELLS	8-10	0-5	/HPF
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED		
BACTERIA	DETECTED	NOT DETECTED	
YEAST	NOT DETECTED	NOT DETECTED	

### Comments

URINALYSIS :- MICROSCOPIC EXAMINATION OF URINE IS CARRIED OUT ON CENTRIFUGED URINARY SEDIMENT. Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of

Conditions

Dr.Karthick Prabhu R Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in Page 16 Of 22





View Report





PATIENT NAME: NITHYA K N		<b>REF. DOCTOR</b> : D	R. BANK O	F BARODA	
CODE/NAME & ADDRESS : C000138396	ACCESSION NO	) : <b>0183WD000644</b>	AGE/SEX	:29 Years	Female
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	PATIENT ID	: NITHF020793183	DRAWN	:07/04/2023	00:00:00
DELHI	CLIENT PATIEN		1	:07/04/2023	
NEW DELHI 110030	ABHA NO	:	REPORTED	:24/04/2023	10:50:37
8800465156					

# Test Report Status Final

Results

**Biological Reference Interval** Units

Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind
	of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

R

Dr.Karthick Prabhu R Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in Page 17 Of 22



View Report





PATIENT NAME: NITHYA K N	REF. DOCTOR : D	R. BANK OF BARODA	
CODE/NAME & ADDRESS : C000138396	ACCESSION NO : 0183WD000644	AGE/SEX : 29 Years Fem	ale
	PATIENT ID : NITHF020793183	DRAWN :07/04/2023 00:00	0:00
DELHI		RECEIVED :07/04/2023 17:33	
NEW DELHI 110030	ABHA NO :	REPORTED :24/04/2023 10:50	):37
8800465156			
(	i		

Test	Report	Status	<u>Final</u>
------	--------	--------	--------------

Results

**Biological Reference Interval** Units

CL	INICAL PATH - STOOL ANAL	ſSIS	
MEDI WHEEL FULL BODY HEALTH CHECK	(UP BELOW 40FEMALE		
PHYSICAL EXAMINATION, STOOL			
COLOUR	BROWN		
CONSISTENCY	WELL FORMED		
MUCUS	NOT DETECTED	NOT DETECTED	
VISIBLE BLOOD	ABSENT	ABSENT	
ADULT PARASITE	NOT DETECTED		
CHEMICAL EXAMINATION, STOOL			
STOOL PH	NEGATIVE		
OCCULT BLOOD	NOT DETECTED	NOT DETECTED	
MICROSCOPIC EXAMINATION, STOOL			
PUS CELLS	1-2		/hpf
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
CYSTS	NOT DETECTED	NOT DETECTED	
OVA	NOT DETECTED		
LARVAE	NOT DETECTED	NOT DETECTED	
TROPHOZOITES	NOT DETECTED	NOT DETECTED	
FAT	ABSENT		
VEGETABLE CELLS	ABSENT		
Interpretation(s)			

## Interpretation(s)


Stool routine analysis is only a screening test for disorders of gastrointentestinal tract like infection, malabsorption, etc. The following table describes the probable conditions, in which the analytes are present in stool.

PRESENCE OF	CONDITION
Pus cells	Pus in the stool is an indication of infection
Red Blood cells	Parasitic or bacterial infection or an inflammatory bowel condition such as ulcerative colitis

Dr.Karthick Prabhu R Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in Page 18 Of 22







**Test Report Status** 

<u>Final</u>



**Biological Reference Interval** Units

PATIENT NAME: NITHYA K N	<b>REF. DOCTOR</b> : D	R. BANK OF BARODA
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST	PATIENT ID : NITHF020793183 CLIENT PATIENT ID:	AGE/SEX :29 Years Female DRAWN :07/04/2023 00:00:00 RECEIVED :07/04/2023 17:33:08 REPORTED :24/04/2023 10:50:37
(		•

Results

Parasites	Infection of the digestive system. Stool examination for ova and parasite detects presence of parasitic infestation of gastrointestinal tract. Various forms of parasite that can be detected include cyst, trophozoite and larvae. One negative result does not rule out the possibility of parasitic infestation. Intermittent
	shedding of parasites warrants examinations of multiple specimens tested on consecutive days.Stool specimens for parasitic examination should be collected before initiation of antidiarrheal therapy or antiparasitic therapy. This test does not detect presence of opportunistic parasites like Cyclospora, Cryptosporidia and Isospora species. Examination of Ova and Parasite has been carried out by
	direct and concentration techniques.
Mucus	Mucus is a protective layer that lubricates, protects& reduces damage due to
	bacteria or viruses.
Charcot-Leyden crystal	Parasitic diseases.
Ova & cyst	Ova & cyst indicate parasitic infestation of intestine.
Frank blood	Bleeding in the rectum or colon.
Occult blood	Occult blood indicates upper GI bleeding.
Macrophages	Macrophages in stool are an indication of infection as they are protective cells.
Epithelial cells	Epithelial cells that normally line the body surface and internal organs show up in stool when there is inflammation or infection.
Fat	Increased fat in stool maybe seen in conditions like diarrhoea or malabsorption.
рН	Normal stool pH is slightly acidic to neutral. Breast-fed babies generally have an acidic stool.

# **ADDITIONAL STOOL TESTS :**

- Stool Culture:- This test is done to find cause of GI infection, make decision about best treatment for GI infection & to find out if 1. treatment for GI infection worked.
- Fecal Calprotectin: It is a marker of intestinal inflammation. This test is done to differentiate Inflammatory Bowel Disease (IBD) 2. from Irritable Bowel Syndrome (IBS).
- 3. Fecal Occult Blood Test(FOBT): This test is done to screen for colon cancer & to evaluate possible cause of unexplained anaemia.
- 4. Clostridium Difficile Toxin Assay: This test is strongly recommended in healthcare associated bloody or waterydiarrhoea, due to overuse of broad spectrum antibiotics which alter the normal GI flora.
- 5. Biofire (Film Array) GI PANEL: In patients of Diarrhoea, Dysentry, Rice watery Stool, FDA approved, Biofire Film Array Test, (Real Time Multiplex PCR) is strongly recommended as it identifies organisms, bacteria, fungi, virus, parasite and other opportunistic pathogens, Vibrio cholera infections only in 3 hours. Sensitivity 96% & Specificity 99%.
- 6. Rota Virus Immunoassay: This test is recommended in severe gastroenteritis in infants & children associated with watery diarrhoea, vomitting& abdominal cramps. Adults are also affected. It is highly contagious in nature.

**Dr.Karthick Prabhu R Consultant Pathologist** 

Page 19 Of 22



Details

Patient Ref. No. 775000002841801



PATIENT NAME: NITHYA K N	<b>REF. DOCTOR :</b> D	R. BANK OF BARODA
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	CLIENT PATIENT ID:	AGE/SEX :29 Years Female DRAWN :07/04/2023 00:00:00 RECEIVED :07/04/2023 17:33:08 REPORTED :24/04/2023 10:50:37
Test Report Status <u>Final</u>	Results Biological	Reference Interval Units

### SPECIALISED CHEMISTRY - HORMONE

SPECIALISED CHEMISTRY - HORMONE				
MEDI WHEEL FULL BODY HEALTH CHE	CKUP BELOW 40FEMALE			
THYROID PANEL, SERUM				
ТЗ	105.40	Non-Pregnant Women 80.0 - 200.0 Pregnant Women 1st Trimester:105.0 - 230.0 2nd Trimester:129.0 - 262.0 3rd Trimester:135.0 - 262.0	0	
Τ4	7.63	Non-Pregnant Women 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70	)	
TSH (ULTRASENSITIVE)	1.820	Non Pregnant Women 0.27 - 4.20 Pregnant Women 1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15	µIU/mL	

## Interpretation(s)

**Triiodothyronine T3**, **Thyroxine T4**, and **Thyroid Stimulating Hormone TSH** are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3.Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism.Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment

Dr.Karthick Prabhu R Consultant Pathologist



Page 20 Of 22

View Report

View Details





#### PATIENT NAME: NITHYA K N REF. DOCTOR : DR. BANK OF BARODA CODE/NAME & ADDRESS : C000138396 ACCESSION NO : 0183WD000644 AGE/SEX :29 Years Female ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) :07/04/2023 00:00:00 PATIENT ID DRAWN : NITHF020793183 F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED :07/04/2023 17:33:08 DELHI ABHA NO REPORTED :24/04/2023 10:50:37 : NEW DELHI 110030 8800465156

## Test Report Status <u>Final</u>

Results

**Biological Reference Interval** Units

_	1	1				
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid	
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto	
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical	
					inflammation, drugs like amphetamines, Iodine containing drug and	
					dopamine antagonist e.g. domperidone and other physiological reasons.	
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism	
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre	
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid	
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4	
					replacement therapy (7) First trimester of Pregnancy	
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism	
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor	
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent	
					treatment for Hyperthyroidism	
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness	
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies	

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. **NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.**TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

> \*\*End Of Report\*\* Please visit www.srlworld.com for related Test Information for this accession



Dr.Karthick Prabhu R Consultant Pathologist









Vie<u>w Report</u>





PATIENT NAME: NITHYA K N	REF. DOCTOR : D	R. BANK OF BARODA
ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI	PATIENT ID : NITHF020793183 CLIENT PATIENT ID:	AGE/SEX:29 YearsFemaleDRAWN:07/04/202300:00:00RECEIVED:07/04/202317:33:08REPORTED:24/04/202310:50:37
Test Report Status <u>Final</u>	Results Biological	Reference Interval Units

## **CONDITIONS OF LABORATORY TESTING & REPORTING**

 It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
 All tests are performed and reported as per the turnaround time stated in the AGILUS Directory of Services.

3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.

4. A requested test might not be performed if:

i. Specimen received is insufficient or inappropriate

- ii. Specimen quality is unsatisfactory
- iii. Incorrect specimen type

iv. Discrepancy between identification on specimen container label and test requisition form

5. AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.

6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.

7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.

8. Test results cannot be used for Medico legal purposes.

9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

(91115 91115) within 48 hours of the report.

#### Agilus Diagnostics Ltd

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062



Dr.Karthick Prabhu R Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd (Formerly SRL Ltd) 57, Cowley Brown Road, R S Puram Coimbatore, 641002 Tamilnadu, India Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956 Email : customercare.coimbatore@srl.in Page 22 Of 22





View Repor

