



Age/Gender : 28 Y 0 M 12 D/M UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM

Received : 09/Dec/2023 11:00AM Reported : 09/Dec/2023 03:00PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF HAEMATOLOGY

ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324

PERIPHERAL SMEAR , WHOLE BLOOD EDTA	

Page 1 of 12







Age/Gender : 28 Y 0 M 12 D/M UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM
Received : 09/Dec/2023 11:00AM

Reported : 09/Dec/2023 03:00PM

Status : Final Report
Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF HAEMATOLOGY ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324 Test Name Result Unit Bio. Ref. Range Method

15.1	g/dL	13-17	CYANIDE FREE COLOUROMETER
46.40	%	40-50	PULSE HEIGHT AVERAGE
4.8	Million/cu.mm	4.5-5.5	Electrical Impedence
96.8	fL	83-101	Calculated
31.5	pg	27-32	Calculated
32.5	g/dL	31.5-34.5	Calculated
13	%	11.6-14	Calculated
6,050	cells/cu.mm	4000-10000	Electrical Impedance
(DLC)			
54.9	%	40-80	Electrical Impedance
35.1	%	20-40	Electrical Impedance
1.6	%	1-6	Electrical Impedance
7.2	%	2-10	Electrical Impedance
1.2	%	<1-2	Electrical Impedance
			•
3321.45	Cells/cu.mm	2000-7000	Calculated
2123.55	Cells/cu.mm	1000-3000	Calculated
96.8	Cells/cu.mm	20-500	Calculated
435.6	Cells/cu.mm	200-1000	Calculated
72.6	Cells/cu.mm	0-100	Calculated
167000	cells/cu.mm	150000-410000	IMPEDENCE/MICROSCOPY
02	mm at the end of 1 hour	0-15	Modified Westergren
	46.40 4.8 96.8 31.5 32.5 13 6,050 (DLC) 54.9 35.1 1.6 7.2 1.2 3321.45 2123.55 96.8 435.6 72.6 167000	46.40 % 4.8 Million/cu.mm 96.8 fL 31.5 pg 32.5 g/dL 13 % 6,050 cells/cu.mm (DLC) 54.9 % 35.1 % 1.6 % 7.2 % 1.2 % 3321.45 Cells/cu.mm 2123.55 Cells/cu.mm 96.8 Cells/cu.mm 435.6 Cells/cu.mm 72.6 Cells/cu.mm 02 mm at the end	46.40 % 40-50 4.8 Million/cu.mm 4.5-5.5 96.8 fL 83-101 31.5 pg 27-32 32.5 g/dL 31.5-34.5 13 % 11.6-14 6,050 cells/cu.mm 4000-10000 (DLC) 54.9 % 40-80 35.1 % 20-40 1.6 % 1-6 7.2 % 2-10 1.2 % <1-2

RBCs ARE NORMOCYTIC NORMOCHROMIC.

TLC, DLC WITHIN NORMAL LIMIT. NO IMMATURE CELLS ARE SEEN.

PLATELETS ARE ADEQUATE.

NO HEMOPARASITES SEEN

Page 2 of 12





: szfgsdzfxh

Age/Gender : 28 Y 0 M 12 D/M

UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777
Ref Doctor : Dr.SELF

Emp/Auth/TPA ID

Collected : 09/Dec/2023 10:15AM
Received : 09/Dec/2023 11:00AM
Reported : 09/Dec/2023 03:00PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF HAEMATOLOGY						
ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324						
Test Name	Test Name Result Unit Bio. Ref. Range Method					

BLOOD GROUP ABO AND RH FACTOR , WHOLE BLOOD EDTA				
BLOOD GROUP TYPE	0		Forward & Reverse Grouping with Slide/Tube Aggluti	
Rh TYPE	POSITIVE		Forward & Reverse Grouping with Slide/Tube Agglutination	

Page 3 of 12







Age/Gender : 28 Y 0 M 12 D/M

UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected

: 09/Dec/2023 02:23PM

Received : 09/Dec/2023 05:56PM

Reported : 09/Dec/2023 06:54PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF BIOCHEMISTRY

ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324					
Test Name Result Unit Bio. Ref. Range Method					

				•
GLUCOSE, FASTING , NAF PLASMA	89	mg/dL	70-100	GOD - POD

Comment:

As per American Diabetes Guidelines, 2023

Fasting Glucose Values in mg/dL	Interpretation
70-100 mg/dL	Normal
100-125 mg/dL	Prediabetes
≥126 mg/dL	Diabetes
<70 mg/dL	Hypoglycemia

Note:

- 1. The diagnosis of Diabetes requires a fasting plasma glucose of > or = 126 mg/dL and/or a random / 2 hr post glucose value of > or = 200 mg/dL on at least 2 occasions.
- 2. Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis & is considered critical.

GLUCOSE, POST PRANDIAL (PP), 2	106	mg/dL	70-140	GOD - POD
HOURS , SODIUM FLUORIDE PLASMA (2				
HR)				

Comment:

It is recommended that FBS and PPBS should be interpreted with respect to their Biological reference ranges and not with each other.

Conditions which may lead to lower postprandial glucose levels as compared to fasting glucose levels may be due to reactive hypoglycemia, dietary meal content, duration or timing of sampling after food digestion and absorption, medications such as insulin preparations, sulfonylureas, amylin analogues, or conditions such as overproduction of insulin.

Page 4 of 12







Age/Gender : 28 Y 0 M 12 D/M UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM

Received : 09/Dec/2023 01:27PM Reported : 09/Dec/2023 02:35PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF BIOCHEMISTRY

ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324					
Test Name	Result	Unit	Bio. Ref. Range	Method	

HBA1C, GLYCATED HEMOGLOBIN , WHOLE BLOOD EDTA	5.1	%	HPLC
ESTIMATED AVERAGE GLUCOSE (eAG),	100	mg/dL	Calculated
WHOLE BLOOD EDTA			

Comment:

Reference Range as per American Diabetes Association (ADA) 2023 Guidelines:

REFERENCE GROUP	нва1С %	
NON DIABETIC	<5.7	
PREDIABETES	5.7 – 6.4	
DIABETES	≥ 6.5	
DIABETICS		
EXCELLENT CONTROL	6 – 7	
FAIR TO GOOD CONTROL	7 – 8	
UNSATISFACTORY CONTROL	8 – 10	
POOR CONTROL	>10	

Note: Dietary preparation or fasting is not required.

- 1. HbA1C is recommended by American Diabetes Association for Diagnosing Diabetes and monitoring Glycemic Control by American Diabetes Association guidelines 2023.
- 2. Trends in HbA1C values is a better indicator of Glycemic control than a single test.
- 3. Low HbA1C in Non-Diabetic patients are associated with Anemia (Iron Deficiency/Hemolytic), Liver Disorders, Chronic Kidney Disease. Clinical Correlation is advised in interpretation of low Values.
- 4. Falsely low HbA1c (below 4%) may be observed in patients with clinical conditions that shorten erythrocyte life span or decrease mean erythrocyte age. HbA1c may not accurately reflect glycemic control when clinical conditions that affect erythrocyte survival are present.
- $5.\ In\ cases\ of\ Interference\ of\ Hemoglobin\ variants\ in\ HbA1C, alternative\ methods\ (Fructosamine)\ estimation\ is\ recommended\ for\ Glycemic\ Control$
 - A: HbF >25%
 - B: Homozygous Hemoglobinopathy.
 - (Hb Electrophoresis is recommended method for detection of Hemoglobinopathy)







Age/Gender : 28 Y 0 M 12 D/M

UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777
Ref Doctor : Dr.SELF

Emp/Auth/TPA ID : szfgsdzfxh

Collected : 09/Dec/2023 10:15AM

Received : 09/Dec/2023 04:09PM Reported : 09/Dec/2023 05:21PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF BIOCHEMISTRY

ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324						
ARCOT LIMI - MILDIWITELE - I GLE BODT ARROAL I LOS MALE - 2D LOTIO - I AR INDIA - I 12324						
Test Name	Result	Unit	Bio. Ref. Range	Method		
)			

LIPID PROFILE, SERUM							
TOTAL CHOLESTEROL	128	mg/dL	<200	CHE/CHO/POD			
TRIGLYCERIDES	67	mg/dL	<150	Enzymatic			
HDL CHOLESTEROL	48	mg/dL	>40	CHE/CHO/POD			
NON-HDL CHOLESTEROL	80	mg/dL	<130	Calculated			
LDL CHOLESTEROL	66.6	mg/dL	<100	Calculated			
VLDL CHOLESTEROL	13.4	mg/dL	<30	Calculated			
CHOL / HDL RATIO	2.67		0-4.97	Calculated			

Comment:

Reference Interval as per National Cholesterol Education Program (NCEP) Adult Treatment Panel III Report.

	Desirable	Borderline High	High	Very High
TOTAL CHOLESTEROL	< 200	200 - 239	≥ 240	
TRIGLYCERIDES	<150	150 - 199	200 - 499	≥ 500
III .I D1 .	Optimal < 100 Near Optimal 100-129	130 - 159	160 - 189	≥ 190
HDL	≥ 60			
INON-HDL CHOLESTEROL	Optimal <130; Above Optimal 130-159	160-189	190-219	>220

- 1. Measurements in the same patient on different days can show physiological and analytical variations.
- 2. NCEP ATP III identifies non-HDL cholesterol as a secondary target of therapy in persons with high triglycerides.
- 3. Primary prevention algorithm now includes absolute risk estimation and lower LDL Cholesterol target levels to determine eligibility of drug therapy.
- **4.** Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.
- **5.** As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.
- **6.** VLDL, LDL Cholesterol Non HDL Cholesterol, CHOL/HDL RATIO, LDL/HDL RATIO are calculated parameters when Triglycerides are below 350mg/dl. When Triglycerides are more than 350 mg/dl LDL cholesterol is a direct measurement.

Page 6 of 12







Age/Gender : 28 Y 0 M 12 D/M UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh

: 09/Dec/2023 10:15AM Collected Received : 09/Dec/2023 04:09PM

Reported : 09/Dec/2023 05:10PM

: Final Report Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF BIOCHEMISTRY

ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324						
Test Name Result Unit Bio. Ref. Range Me						

Status

LIVER FUNCTION TEST (LFT), SERUM								
0.30	mg/dL	0.20-1.20	DIAZO METHOD					
0.20	mg/dL	0.0-0.3	Calculated					
0.10	mg/dL	0.0-1.1	Dual Wavelength					
35	U/L	21-72	UV with P-5-P					
36.0	U/L	17-59	UV with P-5-P					
93.00	U/L	38-126	p-nitrophenyl phosphate					
7.50	g/dL	6.3-8.2	BIURET METHOD					
4.40	g/dL	3.5 - 5	Bromocresol Green					
3.10	g/dL	2.0-3.5	Calculated					
1.42		0.9-2.0	Calculated					
	0.20 0.10 35 36.0 93.00 7.50 4.40 3.10	0.20 mg/dL 0.10 mg/dL 35 U/L 36.0 U/L 93.00 U/L 7.50 g/dL 4.40 g/dL 3.10 g/dL	0.20 mg/dL 0.0-0.3 0.10 mg/dL 0.0-1.1 35 U/L 21-72 36.0 U/L 17-59 93.00 U/L 38-126 7.50 g/dL 6.3-8.2 4.40 g/dL 3.5 - 5 3.10 g/dL 2.0-3.5					

Comment:

LFT results reflect different aspects of the health of the liver, i.e., hepatocyte integrity (AST & ALT), synthesis and secretion of bile (Bilirubin, ALP), cholestasis (ALP, GGT), protein synthesis (Albumin)

Common patterns seen:

1. Hepatocellular Injury:

- AST Elevated levels can be seen. However, it is not specific to liver and can be raised in cardiac and skeletal injuries.
- ALT Elevated levels indicate hepatocellular damage. It is considered to be most specific lab test for hepatocellular injury. Values also correlate well with increasing BMI.
- Disproportionate increase in AST, ALT compared with ALP.
- · Bilirubin may be elevated.
- AST: ALT (ratio) In case of hepatocellular injury AST: ALT > 1In Alcoholic Liver Disease AST: ALT usually >2. This ratio is also seen to be increased in NAFLD, Wilsons's diseases, Cirrhosis, but the increase is usually not >2.

2. Cholestatic Pattern:

- ALP Disproportionate increase in ALP compared with AST, ALT.
- · Bilirubin may be elevated.
- ALP elevation also seen in pregnancy, impacted by age and sex.
- To establish the hepatic origin correlation with GGT helps. If GGT elevated indicates hepatic cause of increased ALP.

3. Synthetic function impairment:

- Albumin- Liver disease reduces albumin levels.
- Correlation with PT (Prothrombin Time) helps.

Page 7 of 12







Age/Gender : 28 Y 0 M 12 D/M
UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM Received : 09/Dec/2023 04:09PM

Reported : 09/Dec/2023 05:21PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF BIOCHEMISTRY ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324 Test Name Result Unit Bio. Ref. Range Method

CREATININE	0.80	mg/dL	0.66-1.25	Creatinine amidohydrolase
UREA	24.00	mg/dL	19-43	Urease
BLOOD UREA NITROGEN	11.2	mg/dL	8.0 - 23.0	Calculated
URIC ACID	5.50	mg/dL	3.5-8.5	Uricase
CALCIUM	9.60	mg/dL	8.4 - 10.2	Arsenazo-III
PHOSPHORUS, INORGANIC	3.80	mg/dL	2.5-4.5	PMA Phenol
SODIUM	133	mmol/L	135-145	Direct ISE
POTASSIUM	4.6	mmol/L	3.5-5.1	Direct ISE
CHLORIDE	101	mmol/L	98 - 107	Direct ISE





Age/Gender : 28 Y 0 M 12 D/M UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM Received : 09/Dec/2023 04:09PM

Reported : 09/Dec/2023 05:21PM

: Final Report Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF BIOCHEMISTRY							
ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324							
Test Name Result Unit Bio. Ref. Range Method							

Status

GAMMA GLUTAMYL TRANSPEPTIDASE	14.00	U/L	15-73	Glyclyclycine
(GGT) , SERUM				Nitoranalide

Page 9 of 12







Age/Gender : 28 Y 0 M 12 D/M UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM

Received : 09/Dec/2023 11:00AM Reported : 09/Dec/2023 03:07PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF IMMUNOLOGY

ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324							
AROOI LIIII - IIILDIIVIILLE - I	ARCOT LIM - MEDIWITELE - FOLE BODT ARROAL FEOS MALE - 2D EGITO - FAR INDIA - F12324						
Test Name Result Unit Bio. Ref. Range Method							
1 COL HAIRE	Result	Oilit	Dio. Non Range	method			

THYROID PROFILE TOTAL (T3, T4, TSH) , SERUM							
TRI-IODOTHYRONINE (T3, TOTAL)	1.45	ng/mL	0.67-1.81	ELFA			
THYROXINE (T4, TOTAL)	7.43	μg/dL	4.66-9.32	ELFA			
THYROID STIMULATING HORMONE (TSH)	1.960	μIU/mL	0.25-5.0	ELFA			

Comment:

lkor nregnant females	Bio Ref Range for TSH in uIU/ml (As per American Thyroid Association)
First trimester	0.1 - 2.5
Second trimester	0.2 - 3.0
Third trimester	0.3 - 3.0

- 1. TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH activates production of T3 (Triiodothyronine) and its prohormone T4 (Thyroxine). Increased blood level of T3 and T4 inhibit production of TSH.
- **2.** TSH is elevated in primary hypothyroidism and will be low in primary hyperthyroidism. Elevated or low TSH in the context of normal free thyroxine is often referred to as sub-clinical hypo- or hyperthyroidism respectively.
- 3. Both T4 & T3 provides limited clinical information as both are highly bound to proteins in circulation and reflects mostly inactive hormone. Only a very small fraction of circulating hormone is free and biologically active.
- 4. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, medication & circulating antibodies.

TSH	Т3	Т4	FT4	Conditions
High	Low	Low	Low	Primary Hypothyroidism, Post Thyroidectomy, Chronic Autoimmune Thyroiditis
High	N	N	N	Subclinical Hypothyroidism, Autoimmune Thyroiditis, Insufficient Hormone Replacement Therapy.
N/Low	Low	Low	Low	Secondary and Tertiary Hypothyroidism
Low	High	High	High	Primary Hyperthyroidism, Goitre, Thyroiditis, Drug effects, Early Pregnancy
Low	N	N	N	Subclinical Hyperthyroidism
Low	Low	Low	Low	Central Hypothyroidism, Treatment with Hyperthyroidism
Low	N	High	High	Thyroiditis, Interfering Antibodies
N/Low	High	N	N	T3 Thyrotoxicosis, Non thyroidal causes
High	High	High	High	Pituitary Adenoma; TSHoma/Thyrotropinoma

Page 10 of 12







Age/Gender : 28 Y 0 M 12 D/M
UHID/MR No : SCHI.0000016657

Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM
Received : 09/Dec/2023 12:21PM

Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

: 09/Dec/2023 02:32PM

DEPARTMENT OF CLINICAL PATHOLOGY							
ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324							
Test Name Result Unit Bio. Ref. Range Method							

Reported

COMPLETE URINE EXAMINATION (CUE)	, URINE			
PHYSICAL EXAMINATION				
COLOUR	PALE YELLOW		PALE YELLOW	Visual
TRANSPARENCY	CLEAR		CLEAR	Visual
рН	6.5		5-7.5	Bromothymol Blue
SP. GRAVITY	1.015		1.002-1.030	Dipstick
BIOCHEMICAL EXAMINATION	•		•	
URINE PROTEIN	NEGATIVE	NEGATIVE		PROTEIN ERROR OF INDICATOR
GLUCOSE	NEGATIVE		NEGATIVE	GOD-POD
URINE BILIRUBIN	NEGATIVE		NEGATIVE	AZO COUPLING
URINE KETONES (RANDOM)	NEGATIVE		NEGATIVE	NITROPRUSSIDE
UROBILINOGEN	NORMAL		NORMAL	EHRLICH
BLOOD	NEGATIVE		NEGATIVE	Dipstick
NITRITE	NEGATIVE		NEGATIVE	Dipstick
LEUCOCYTE ESTERASE	NEGATIVE		NEGATIVE	PYRROLE HYDROLYSIS
CENTRIFUGED SEDIMENT WET MOUNT	AND MICROSCOPY	(
PUS CELLS	0-2	/hpf	0-5	Microscopy
EPITHELIAL CELLS	0-2	/hpf	<10	MICROSCOPY
RBC	ABSENT	/hpf	0-2	MICROSCOPY
CASTS	ABSENT		0-2 Hyaline Cast	MICROSCOPY
CRYSTALS	ABSENT		ABSENT	MICROSCOPY

Page 11 of 12







Age/Gender : 28 Y 0 M 12 D/M

UHID/MR No : SCHI.0000016657 Visit ID : SCHIOPV23777

Ref Doctor : Dr.SELF Emp/Auth/TPA ID : szfgsdzfxh Collected : 09/Dec/2023 10:15AM Received : 09/Dec/2023 12:21PM

Reported : 09/Dec/2023 02:32PM Status : Final Report

Sponsor Name : ARCOFEMI HEALTHCARE LIMITED

DEPARTMENT OF CLINICAL PATHOLOGY

ARCOFEMI - MEDIWHEEL - F	ARCOFEMI - MEDIWHEEL - FULL BODY ANNUAL PLUS MALE - 2D ECHO - PAN INDIA - FY2324						
Test Name	Result	Unit	Bio. Ref. Range	Method			

URINE GLUCOSE(POST PRANDIAL)	NEGATIVE	NEGATIVE	Dipstick	
URINE GLUCOSE(FASTING)	NEGATIVE	NEGATIVE	Dinstick	

*** End Of Report ***

Dr. Tanish Mandal
Dr. SHWETA GUPTA
MBBS,MD (Pathology)
Consultant Pathology

Consultant Pathology

Dr Nidhi Sachdev M.B.B.S,MD(Pathology) Consultant Pathologist

Page 12 of 12

