

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

PATIENT NAME : BHAWANI SHANKAR

PATIENT ID : BHAWM050992251

ACCESSION NO : 0251VI000546 AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

**MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE**

**BLOOD COUNTS, EDTA WHOLE BLOOD**

HEMOGLOBIN	16.4	13.0 - 17.0	g/dL
METHOD : CYANIDE FREE DETERMINATION			
RED BLOOD CELL COUNT	<b>5.63</b>	<b>High</b> 4.5 - 5.5	mil/ $\mu$ L
METHOD : ELECTRICAL IMPEDANCE			
WHITE BLOOD CELL COUNT	7.20	4.0 - 10.0	thou/ $\mu$ L
METHOD : ELECTRICAL IMPEDANCE			
PLATELET COUNT	178	150 - 410	thou/ $\mu$ L
METHOD : ELECTRONIC IMPEDANCE			

**RBC AND PLATELET INDICES**

HEMATOCRIT	47.9	40 - 50	%
METHOD : CALCULATED PARAMETER			
MEAN CORPUSCULAR VOL	85.0	83 - 101	fL
METHOD : CALCULATED PARAMETER			
MEAN CORPUSCULAR HGB,	29.1	27.0 - 32.0	pg
METHOD : CALCULATED PARAMETER			
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION	34.3	31.5 - 34.5	g/dL
METHOD : CALCULATED PARAMETER			
MENTZER INDEX	15.1		
RED CELL DISTRIBUTION WIDTH	13.2	11.6 - 14.0	%
METHOD : CALCULATED PARAMETER			
MEAN PLATELET VOLUME	10.8	6.8 - 10.9	fL
METHOD : CALCULATED PARAMETER			

**WBC DIFFERENTIAL COUNT - NLR**

SEGMENTED NEUTROPHILS	62	40 - 80	%
METHOD : IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY			
ABSOLUTE NEUTROPHIL COUNT	4.46	2.0 - 7.0	thou/ $\mu$ L
METHOD : CALCULATED PARAMETER			
LYMPHOCYTES	30	20 - 40	%
METHOD : IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY			
ABSOLUTE LYMPHOCYTE COUNT	2.16	1.0 - 3.0	thou/ $\mu$ L
METHOD : CALCULATED PARAMETER			
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	2.0		
EOSINOPHILS	<b>07</b>	<b>High</b> 1 - 6	%
METHOD : IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY			
ABSOLUTE EOSINOPHIL COUNT	0.50	0.02 - 0.50	thou/ $\mu$ L



Scan to View Details



Scan to View Report

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

PATIENT NAME : BHAWANI SHANKAR

PATIENT ID : BHAWM050992251

ACCESSION NO : 0251VI000546 AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

METHOD : CALCULATED PARAMETER				
MONOCYTES	01	Low	2 - 10	%
METHOD : IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY				
ABSOLUTE MONOCYTE COUNT	0.07	Low	0.2 - 1.0	thou/ $\mu$ L
METHOD : CALCULATED PARAMETER				
BASOPHILS	00		0 - 2	%
METHOD : IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY				
ABSOLUTE BASOPHIL COUNT	0	Low	0.02 - 0.10	thou/ $\mu$ L
DIFFERENTIAL COUNT PERFORMED ON:	EDTA SMEAR			
<b>* ERYTHRO SEDIMENTATION RATE, BLOOD</b>				
SEDIMENTATION RATE (ESR)	06		0 - 14	mm at 1 hr
METHOD : WESTERGREN METHOD				
<b>GLUCOSE, FASTING, PLASMA</b>				
GLUCOSE, FASTING, PLASMA	93		74 - 99	mg/dL
METHOD : GLUCOSE OXIDASE				
<b>GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD</b>				
GLYCOSYLATED HEMOGLOBIN (HBA1C)	5.8	High	Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 ADA Target: 7.0 Action suggested: > 8.0	%
METHOD : HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC)				
MEAN PLASMA GLUCOSE	119.8	High	< 116.0	mg/dL
METHOD : CALCULATED PARAMETER				
<b>GLUCOSE, POST-PRANDIAL, PLASMA</b>				
GLUCOSE, POST-PRANDIAL, PLASMA	116		70 - 140	mg/dL
METHOD : GLUCOSE OXIDASE				
<b>CORONARY RISK PROFILE, SERUM</b>				
CHOLESTEROL	195		< 200 Desirable 200 - 239 Borderline High >/= 240 High	mg/dL
METHOD : CHOLESTEROL OXIDASE				
TRIGLYCERIDES	225	High	< 150 Normal 150 - 199 Borderline High 200 - 499 High >/=500 Very High	mg/dL
METHOD : LIPASE/GPO-PAP NO CORRECTION				
HDL CHOLESTEROL	44		< 40 Low >/=60 High	mg/dL
METHOD : DIRECT CLEARANCE METHOD				



Scan to View Details



Scan to View Report

**Patient Ref. No. 25100000154496**

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

**PATIENT NAME : BHAWANI SHANKAR**

PATIENT ID : **BHAWM050992251**

ACCESSION NO : **0251VI000546** AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Test Report Status	Final	Results	Biological Reference Interval	Units
CHOLESTEROL LDL		<b>106</b>	<b>High</b> < 100 Optimal 100 - 129 Near optimal/ above optimal 130 - 159 Borderline High 160 - 189 High >= 190 Very High	mg/dL
NON HDL CHOLESTEROL		<b>151</b>	<b>High</b> Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
METHOD : CALCULATED PARAMETER				
CHOL/HDL RATIO		4.4	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk	
LDL/HDL RATIO		2.4	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	
VERY LOW DENSITY LIPOPROTEIN		<b>45.0</b>	<b>High</b> </= 30.0	mg/dL
<b>LIVER FUNCTION PROFILE, SERUM</b>				
BILIRUBIN, TOTAL		<b>1.67</b>	<b>High</b> 0 - 1	mg/dL
METHOD : DIAZO WITH SULPHANILIC ACID				
BILIRUBIN, DIRECT		<b>0.36</b>	<b>High</b> 0.00 - 0.25	mg/dL
METHOD : DIAZO WITH SULPHANILIC ACID				
BILIRUBIN, INDIRECT		<b>1.31</b>	<b>High</b> 0.1 - 1.0	mg/dL
METHOD : CALCULATED PARAMETER				
TOTAL PROTEIN		<b>8.4</b>	<b>High</b> 6.4 - 8.2	g/dL
METHOD : BIURET REACTION, END POINT				
ALBUMIN		<b>4.7</b>	<b>High</b> 3.8 - 4.4	g/dL
METHOD : BROMOCRESOL GREEN				
GLOBULIN		3.7	2.0 - 4.1	g/dL
METHOD : CALCULATED PARAMETER				
ALBUMIN/GLOBULIN RATIO		1.3	1.0 - 2.1	RATIO
METHOD : CALCULATED PARAMETER				
ASPARTATE AMINOTRANSFERASE (AST/SGOT)		<b>39</b>	<b>High</b> 0 - 37	U/L
METHOD : TRIS BUFFER NO P5P IFCC / SFBC 37° C				
ALANINE AMINOTRANSFERASE (ALT/SGPT)		<b>85</b>	<b>High</b> 0 - 40	U/L
METHOD : TRIS BUFFER NO P5P IFCC / SFBC 37° C				



Scan to View Details



Scan to View Report

**Patient Ref. No. 251000000154496**

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

PATIENT NAME : BHAWANI SHANKAR

PATIENT ID : BHAWM050992251

ACCESSION NO : 0251VI000546 AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Test Report Status	Final	Results	Biological Reference Interval	Units
ALKALINE PHOSPHATASE		94	39 - 117	U/L
METHOD : AMP OPTIMISED TO IFCC 37° C				
GAMMA GLUTAMYL TRANSFERASE (GGT)		40	11 - 50	U/L
METHOD : GAMMA GLUTAMYL-3 CARBOXY-4 NITROANILIDE (IFCC) 37° C				
LACTATE DEHYDROGENASE		423	230 - 460	U/L
METHOD : GERMAN METHODS 37° C				
<b>SERUM BLOOD UREA NITROGEN</b>				
BLOOD UREA NITROGEN		12	5.0 - 18.0	mg/dL
METHOD : UREASE KINETIC				
<b>CREATININE, SERUM</b>				
CREATININE		1.03	0.8 - 1.3	mg/dL
METHOD : ALKALINE PICRATE NO DEPROTEINIZATION				
<b>BUN/CREAT RATIO</b>				
BUN/CREAT RATIO		11.65		
METHOD : CALCULATED PARAMETER				
<b>URIC ACID, SERUM</b>				
URIC ACID		5.8	3.4 - 7.0	mg/dL
METHOD : URICASE PEROXIDASE WITH ASCORBATE OXIDASE				
<b>TOTAL PROTEIN, SERUM</b>				
TOTAL PROTEIN		8.4	High 6.4 - 8.3	g/dL
METHOD : BIURET REACTION, END POINT				
<b>ALBUMIN, SERUM</b>				
ALBUMIN		4.7	High 3.8 - 4.4	g/dL
METHOD : BROMOCRESOL GREEN				
<b>GLOBULIN</b>				
GLOBULIN		3.7	2.0 - 4.1	g/dL
METHOD : CALCULATED PARAMETER				
<b>ELECTROLYTES (NA/K/CL), SERUM</b>				
SODIUM		143.4	137 - 145	mmol/L
METHOD : ION-SELECTIVE ELECTRODE				
POTASSIUM		4.26	3.6 - 5.0	mmol/L
METHOD : ION-SELECTIVE ELECTRODE				
CHLORIDE		104.6	98 - 107	mmol/L
METHOD : ION-SELECTIVE ELECTRODE				
<b>PHYSICAL EXAMINATION, URINE</b>				
COLOR		PALE YELLOW		
METHOD : GROSS EXAMINATION				
APPEARANCE		CLEAR		



Scan to View Details



Scan to View Report

**Patient Ref. No. 251000000154496**

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

**PATIENT NAME : BHAWANI SHANKAR**

PATIENT ID : **BHAWM050992251**

ACCESSION NO : **0251VI000546** AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

METHOD : GROSS EXAMINATION

SPECIFIC GRAVITY

1.020

1.003 - 1.035

METHOD : IONIC CONCENTRATION METHOD

**CHEMICAL EXAMINATION, URINE**

PH

6.0

4.7 - 7.5

METHOD : DOUBLE INDICATOR PRINCIPLE

PROTEIN

NOT DETECTED

NOT DETECTED

METHOD : PROTEIN ERROR OF INDICATORS WITH REFLECTANCE

GLUCOSE

NOT DETECTED

NOT DETECTED

METHOD : GLUCOSE OXIDASE PEROXIDASE / BENEDICTS

KETONES

NOT DETECTED

NOT DETECTED

METHOD : SODIUM NITROPRUSSIDE REACTION

BLOOD

NOT DETECTED

NOT DETECTED

METHOD : PEROXIDASE ANTI PEROXIDASE

BILIRUBIN

NOT DETECTED

NOT DETECTED

METHOD : DIPSTICK

UROBILINOGEN

NORMAL

NORMAL

METHOD : EHRLICH REACTION REFLECTANCE

NITRITE

NOT DETECTED

NOT DETECTED

METHOD : NITRATE TO NITRITE CONVERSION METHOD

LEUKOCYTE ESTERASE

NOT DETECTED

NOT DETECTED

**MICROSCOPIC EXAMINATION, URINE**

PUS CELL (WBC'S)

1-2

0-5

/HPF

METHOD : DIPSTICK, MICROSCOPY

EPITHELIAL CELLS

0-1

0-5

/HPF

METHOD : MICROSCOPIC EXAMINATION

ERYTHROCYTES (RBC'S)

NOT DETECTED

NOT DETECTED

/HPF

METHOD : MICROSCOPIC EXAMINATION

CASTS

NOT DETECTED

METHOD : MICROSCOPIC EXAMINATION

CRYSTALS

NOT DETECTED

METHOD : MICROSCOPIC EXAMINATION

BACTERIA

NOT DETECTED

NOT DETECTED

METHOD : MICROSCOPIC EXAMINATION

YEAST

NOT DETECTED

NOT DETECTED

**THYROID PANEL, SERUM**

T3

123,6

60,0 - 181,0

ng/dL

METHOD : CHEMILUMINESCENCE



Scan to View Details



Scan to View Report

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,
Tonk Road
JAIPUR, 302015
Rajasthan, INDIA

PATIENT NAME : BHAWANI SHANKAR

PATIENT ID : BHAWM050992251

ACCESSION NO : 0251VI000546 AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Table with 4 columns: Test Report Status, Results, Biological Reference Interval, Units. Status is Final.

T4 10.10 4.5 - 10.9 µg/dL
METHOD : CHEMILUMINESCENCE

TSH 3RD GENERATION 2.675 0.550 - 4.780 µIU/mL
METHOD : CHEMILUMINESCENCE

STOOL: OVA & PARASITE

COLOUR SAMPLE NOT RECEIVED
METHOD : GROSS EXAMINATION

\* ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE A
METHOD : TUBE AGGLUTINATION

RH TYPE POSITIVE
METHOD : TUBE AGGLUTINATION

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-

Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT - NLR-

The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients ; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

ERYTHRO SEDIMENTATION RATE, BLOOD-

Erythrocyte sedimentation rate (ESR) is a non-specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0-1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

Reference :

- 1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition
2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin
3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition"

GLUCOSE, FASTING, PLASMA-

ADA 2021 guidelines for adults, after 8 hrs fasting is as follows:

Pre-diabetics: 100 - 125 mg/dL

Diabetic: > or = 126 mg/dL

GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-

Glycosylated hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood, the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells.

Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia, increased red cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of testing such as glycated serum protein (fructosamine) should be considered.

"Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of



Scan to View Details



Scan to View Report

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,
Tonk Road
JAIPUR, 302015
Rajasthan, INDIA

PATIENT NAME : BHAWANI SHANKAR

PATIENT ID : BHAWM050992251

ACCESSION NO : 0251VI000546 AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Table with 4 columns: Test Report Status (Final), Results, Biological Reference Interval, Units

diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations."

References

- 1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006, 879-884.
2. Forsham PH. Diabetes Mellitus:A rational plan for management. Postgrad Med 1982, 71,139-154.
3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184.
GLUCOSE, POST-PRANDIAL, PLASMA-ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75grams of glucose in 300 ml water,over a period of 5 minutes.

LIVER FUNCTION PROFILE, SERUM- LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

SERUM BLOOD UREA NITROGEN-

Causes of Increased levels

Pre renal

- High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal

• Renal Failure

Post Renal

- Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

- Liver disease

• STADH.

CREATININE, SERUM-

Higher than normal level may be due to:

- Blockage in the urinary tract
• Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
• Loss of body fluid (dehydration)
• Muscle problems, such as breakdown of muscle fibers
• Problems during pregnancy, such as seizures (eclampsia), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis

• Muscular dystrophy

URIC ACID, SERUM-

Causes of Increased levels

Dietary

- High Protein Intake.

• Prolonged Fasting,

• Rapid weight loss.

Gout

Lesch nyhan syndrome.

Type 2 DM.

Metabolic syndrome.



Scan to View Details



Scan to View Report

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,
Tonk Road
JAIPUR, 302015
Rajasthan, INDIA

PATIENT NAME : BHAWANI SHANKAR

PATIENT ID : BHAWM050992251

ACCESSION NO : 0251VI000546 AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Table with 4 columns: Test Report Status (Final), Results, Biological Reference Interval, Units

Causes of decreased levels

- Low Zinc Intake
OCP's
Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels

- Drink plenty of fluids
Limit animal proteins
High Fibre foods
Vit C Intake
Antioxidant rich foods

TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum..Protein in the plasma is made up of albumin and globulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

ELECTROLYTES (NA/K/CL), SERUM-

Sodium levels are increased in dehydration, cushing's syndrome, aldosteronism & decreased in Addison's disease, hypopituitarism, liver disease. Hypokalemia (low K) is common in vomiting, diarrhea, alcoholism, folic acid deficiency and primary aldosteronism. Hyperkalemia may be seen in end-stage renal failure, hemolysis, trauma, Addison's disease, metabolic acidosis, acute starvation, dehydration, and with rapid K infusion. Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bicarbonate, diabetes insipidus, adrenocortical hyperfunction, salicylate intoxication and with excessive infusion of isotonic saline or extremely high dietary intake of salt. Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and prolonged vomiting,

MICROSCOPIC EXAMINATION, URINE-

Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.

Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia

THYROID PANEL, SERUM-

Triiodothyronine T3, is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.

Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3

Table with 4 columns: Levels in Pregnancy, TOTAL T4 (µg/dL), TSH3G (µIU/mL), TOTAL T3 (ng/dL). Rows for 1st, 2nd, and 3rd Trimester.

Below mentioned are the guidelines for age related reference ranges for T3 and T4.

Table with 2 columns: T3 (ng/dL), T4 (µg/dL). Rows for New Born, 1-3 day, and 1 Week.



Scan to View Details



Scan to View Report



**Patient Ref. No. 25100000154496**

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

**PATIENT NAME : BHAWANI SHANKAR**

PATIENT ID : **BHAWM050992251**

ACCESSION NO : **0251VI000546** AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

**Test Report Status** Final

**Results**

**Biological Reference Interval Units**

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group.

Kindly note: Method specific reference ranges are appearing on the report under biological reference range.

Reference:

1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.

2. Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.

3. Behrman R.E. Kliegman R.M., Jenson H. B. Nelson Text Book of Pediatrics, 17th Edition

STOOL: OVA & PARASITE-

Acute infective diarrhoea and gastroenteritis (diarrhoea with vomiting) are major causes of ill health and premature death in developing countries. Loss of water and electrolytes from the body can lead to severe dehydration which if untreated, can be rapidly fatal in young children, especially that are malnourished, hypoglycaemic, and generally in poor health.

Laboratory diagnosis of parasitic infection is mainly based on microscopic examination and the gross examination of the stool specimen. Depending on the nature of the parasite, the microscopic observations include the identification of cysts, ova, trophozoites, larvae or portions of adult structure. The two classes of parasites that cause human infection are the Protozoa and Helminths. The protozoan infections include amoebiasis mainly caused by Entamoeba histolytica and giardiasis caused by Giardia lamblia. The common helminthic parasites are Trichuris trichiura, Ascaris lumbricoides, Strongyloides stercoralis, Taenia sp. etc

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-

Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.



Scan to View Details



Scan to View Report

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

PATIENT NAME : BHAWANI SHANKAR

PATIENT ID : BHAWM050992251

ACCESSION NO : 0251VI000546 AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

**OUT OF RANGE REPORT**

**MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40**

**MALE BLOOD COUNTS, EDTA WHOLE BLOOD**

RED BLOOD CELL COUNT 5.63 High 4.5 - 5.5 mil/ $\mu$ L

**LIVER FUNCTION PROFILE, SERUM**

BILIRUBIN, DIRECT 0.36 High 0.00 - 0.25 mg/dL

TOTAL PROTEIN 8.4 High 6.4 - 8.2 g/dL

ALANINE AMINOTRANSFERASE (ALT/SGPT) 85 High 0 - 40 U/L

ASPARTATE AMINOTRANSFERASE (AST/SGOT) 39 High 0 - 37 U/L

BILIRUBIN, TOTAL 1.67 High 0 - 1 mg/dL

BILIRUBIN, INDIRECT 1.31 High 0.1 - 1.0 mg/dL

ALBUMIN 4.7 High 3.8 - 4.4 g/dL

**TOTAL PROTEIN, SERUM**

TOTAL PROTEIN 8.4 High 6.4 - 8.3 g/dL

**ALBUMIN, SERUM**

ALBUMIN 4.7 High 3.8 - 4.4 g/dL

**WBC DIFFERENTIAL COUNT - NLR**

ABSOLUTE BASOPHIL COUNT 0 Low 0.02 - 0.10 thou/ $\mu$ L

EOSINOPHILS 07 High 1 - 6 %

ABSOLUTE MONOCYTE COUNT 0.07 Low 0.2 - 1.0 thou/ $\mu$ L

MONOCYTES 01 Low 2 - 10 %

**GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD**

MEAN PLASMA GLUCOSE 119.8 High < 116.0 mg/dL

GLYCOSYLATED HEMOGLOBIN (HBA1C) 5.8 High  
Non-diabetic: < 5.7 %  
Pre-diabetics: 5.7 - 6.4  
Diabetics: > or = 6.5  
ADA Target: 7.0  
Action suggested: > 8.0

**CORONARY RISK PROFILE, SERUM**

TRIGLYCERIDES 225 High  
< 150 Normal mg/dL  
150 - 199 Borderline  
High  
200 - 499 High  
> / = 500 Very High



Scan to View Details



Scan to View Report

**Patient Ref. No. 251000000154496**

CLIENT CODE : C000138404

CLIENT'S NAME AND ADDRESS :

PROVISIONAL REPORT

SRL Ltd  
C/o Aakriti Labs Pvt Ltd, 3, Mahatma Gandhi Marg, Gandhi Nagar Mod,  
Tonk Road  
JAIPUR, 302015  
Rajasthan, INDIA

**PATIENT NAME : BHAWANI SHANKAR**

PATIENT ID : **BHAWM050992251**

ACCESSION NO : **0251VI000546** AGE : 30 Years SEX : Male

ABHA NO :

DRAWN : 05/09/2022 11:20

RECEIVED : 05/09/2022 12:54

REPORTED : 05/09/2022 20:01

REFERRING DOCTOR : SELF

CLIENT PATIENT ID : 012209050041

Test Report Status	Final	Results	Biological Reference Interval	Units
CHOLESTEROL LDL		106	<b>High</b> < 100 Optimal 100 - 129 Near optimal/ above optimal 130 - 159 Borderline High 160 - 189 High >/= 190 Very High	mg/dL
VERY LOW DENSITY LIPOPROTEIN		45.0	<b>High</b> </= 30.0	mg/dL
NON HDL CHOLESTEROL		151	<b>High</b> Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL

INVESTIGATOR : \_\_\_\_\_ MD  
DATE:

**\*\*End Of Report\*\***

**Please visit [www.srlworld.com](http://www.srlworld.com) for related Test Information for this accession**



Scan to View Details



Scan to View Report



Name : **Mr. BHAWANI SHANKAR**  
Age/Gender: 30 Y/Male  
Patient ID : 012209050041  
BarcodeNo : 10059652  
Referred By : Self

Registration No: 40938  
Registered : 05/Sep/2022 11:20AM  
Analysed : 06/Sep/2022 12:06PM  
Reported : 06/Sep/2022 12:06PM  
Panel : Medi Wheel (ArcoFemi  
Healthcare Ltd)


## DIGITAL X-RAY CHEST PA VIEW

Soft tissue shadow and bony cages are normal.  
Trachea is central.  
Bilateral lung field and both CP angle are clear.  
Domes of diaphragm are normally placed.  
Transverse diameter of heart appears with normal limits.


**IMPRESSION:- NO OBVIOUS ABNORMALITY DETECTED.**

\*\*\* End Of Report \*\*\*

Page 1 of 1

  
Dr. Neera Mehta  
M.B.B.S., D.M.R.D.  
RMCNO.005807/14853





NORMAL

DR. ARIF HUSSAIN KHAN  
MBBS PGDCC

RMC NUMBER 023361

(ADX\_GEM217220330)(A)Allengers



SHAWANI SHANKAR / 30 Yrs / M / 0 Cms / 0 Kg  
 / 04 / 2022  
 Angina / Hypercholesterolemia / Diabetic / Positive Estrogen / Non-Athlete

Time	Duration	Speed (mpn)	Elevation	MEIS	Rate	% THR	BP	RPP	PVC	Comments
00:05	0:05	00.0	00.0	01.0	054	28 %	114/65	061	00	
00:16	0:11	00.0	00.0	01.0	056	29 %	114/65	063	00	
00:32	0:16	00.0	00.0	01.0	078	41 %	114/65	088	00	
00:41	0:09	00.0	00.0	01.0	078	41 %	114/65	088	00	
00:46	0:05	01.7	10.0	01.1	064	34 %	114/65	072	00	
03:46	3:00	01.7	10.0	04.7	086	45 %	114/65	098	00	
06:46	3:00	02.5	12.0	07.1	099	52 %	114/65	112	00	
09:46	3:00	03.4	14.0	10.2	124	65 %	114/65	141	00	
11:18	1:32	04.2	16.0	11.9	156	82 %	114/65	177	00	
12:18	1:00	00.0	00.0	04.3	118	62 %	114/65	134	00	
13:18	2:00	00.0	00.0	01.0	102	54 %	114/65	116	00	
14:18	3:00	00.0	00.0	01.0	097	51 %	144/80	139	00	
15:18	4:00	00.0	00.0	01.0	086	45 %	144/80	123	00	
15:49	4:31	00.0	00.0	01.0	092	48 %	135/75	124	00	

**FINDINGS :**

Exercise Time : 10:32  
 Max HR Attained : 156 bpm 82% of Target 190  
 Max BP Attained : 144/80 (mm/Hg)  
 Max WorkLoad Attained : 11.9 Good response to induced stress  
 Max ST Dep Lead & Avg ST Value : avL & -0.4 mm in Recovery  
 Duke Treadmill Score : 09.5  
 Test End Reasons : Test Complete, Heart Rate Achieved

**REPORT :**

**FINAL IMPRESSION - TEST IS NEGATIVE FOR INDUCIBLE ISCHAEMIA**

**DR. ARIF HUSSAIN KHAN**  
 MBBS PGDCC  
 RMC NUMBER 023361  
 NORMAL  
 Doctor : DR. ARIF HUSSAIN