

Name	V. SATYANARAYANA	ID	MED111551707
Age & Gender	38Y/M	Visit Date	Mar 25 2023 9:20AM
Ref Doctor	MediWheel		

ULTRASOUND WHOLE ABDOMEN

Liver : Mildly enlarged in size (16.2 cm) with diffuse increase in

echotexture.

There is no evidence of IHBR / EHBR dilatation seen.

No focal space occupying lesions seen.

CBD is normal. PV normal.

Gall Bladder : Normal in volume and wall thickness.

No e/o intraluminal calculi seen.

Pancreas : Head, body and tail are identified with normal echopattern and

smooth outlines.

Spleen : Measured 10.0 cm, in size with normal echotexture.

Right kidney: Measured $10.7 \times 5.4 \text{ cm}$ in size.

Left kidney : Measured 9.7 x 5.5 cm in size.

Both kidneys are normal in size, position, with well preserved cortico medullary differentiation and normal

pelvicalyceal anatomy.

No e/o space occupying lesion seen.

No e/o suprarenal / retroperitoneal masses noted.

- 4 mm calculus is noted in lower pole of right kidney &

inter pole of left kidney.

Urinary : Normal in volume and wall thickness.
bladder No e/o intraluminal calculi / masses seen.

100 C/O mitratummar carcum/ masses seen.

Prostate : Measured 3.1 x 3.3 x 3.0 cm in size (Vol : 16.2 cc) with normal

echotexture.

No e/o ascites / pleural effusion seen. No e/o detectable bowel pathology seen.



Name	V. SATYANARAYANA	ID	MED111551707
Age & Gender	38Y/M	Visit Date	Mar 25 2023 9:20AM
Ref Doctor	MediWheel		

IMPRESSION:

- Mild hepatomegaly with Grade II steatosis To correlate with LFT.
- Bilateral non obstructive nephrolithiasis.
- For clinical correlation.

Dr.Jahn av i Barla ,MD (RD)

Consultant Radiologist

PID No. : MED111551707 **Register On** : 25/03/2023 9:21 AM : 80026874 SID No. Collection On : 25/03/2023 9:55 AM Age / Sex : 38 Year(s) / Male Report On : 25/03/2023 2:08 PM Type : OP

Printed On



Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
BLOOD GROUPING AND Rh TYPING (Blood/Agglutination)	'O' 'Positive'		
Complete Blood Count With - ESR			
Haemoglobin (Blood/Spectrophotometry)	16.0	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (Blood/Numeric Integration of MCV)	47.7	%	42 - 52
RBC Count (Blood/Electrical Impedance)	5.58	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (Blood/Calculated)	85.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (Blood/Calculated)	28.8	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (Blood/Calculated)	33.7	g/dL	32 - 36
RDW-CV (Calculated)	14.4	%	11.5 - 16.0
RDW-SD (Calculated)	43.04	fL	39 - 46
Total Leukocyte Count (TC) (Blood/Electrical Impedance)	7650	cells/cu.mm	4000 - 11000
Neutrophils (Blood/ <i>Impedance and absorbance</i>)	65.42	%	40 - 75
Lymphocytes (Blood/ <i>Impedance and absorbance</i>)	24.43	%	20 - 45
Eosinophils (Blood/Impedance and absorbance)	3.14	%	01 - 06

: 01/04/2023 12:27 PM







PID No. : MED111551707 **Register On** : 25/03/2023 9:21 AM : 80026874 SID No. Collection On : 25/03/2023 9:55 AM Age / Sex : 38 Year(s) / Male Report On : 25/03/2023 2:08 PM Type : OP

Printed On



Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes	6.70	%	01 - 10
(Blood/Impedance and absorbance)			
Basophils (Blood/Impedance and absorbance)	0.32	%	00 - 02
INTERPRETATION: Tests done on Automated F	ive Part cell counte	er. All abnormal results are re-	viewed and confirmed microscopically.
Absolute Neutrophil count (Blood/Impedance and absorbance)	5.00	10^3 / μ1	1.5 - 6.6
Absolute Lymphocyte Count (Blood/Impedance)	1.87	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (Blood/Impedance)	0.24	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (Blood/Impedance)	0.51	10^3 / μ1	< 1.0
Absolute Basophil count (Blood/Impedance)	0.02	10^3 / μ1	< 0.2
Platelet Count (Blood/Impedance)	2.15	lakh/cu.mm	1.4 - 4.5
INTERPRETATION: Platelet count less than 1.5	lakhs will be confir	med microscopically.	
MPV (Blood/Derived from Impedance)	7.20	fL	7.9 - 13.7
PCT (Calculated)	0.15	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated ESR analyser)	04	mm/hr	< 15
BUN / Creatinine Ratio	11.2		
Glucose Fasting (FBS) (Plasma - F/Glucose oxidase/Peroxidase)	95	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 01/04/2023 12:27 PM









 PID No.
 : MED111551707
 Register On : 25/03/2023 9:21 AM

 SID No.
 : 80026874
 Collection On : 25/03/2023 9:55 AM

 Age / Sex : 38 Year(s) / Male
 Report On : 25/03/2023 2:08 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

: 01/04/2023 12:27 PM

8		
Glucose, Fasting (Urine)	Negative	Negative
(Urine - F)		

Glucose Postprandial (PPBS) 180 mg/dL 70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Calculated)	11.2	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe - Alkaline Picrate)	1	mg/dL	0.9 - 1.3
Uric Acid (Serum/Uricase/Peroxidase)	5.8	mg/dL	3.5 - 7.2
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulphanilic acid)	1.0	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulphanilic acid)	0.3	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Calculated)	0.70	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC without P-5-P)	24	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC without P-5-P)	26	U/L	5 - 41







APPROVED BY

The results pertain to sample tested.

Page 3 of 8

Lab Address: MEDALL HEALTH CARE PVT LTD,#17-11-3/4,DR.GKS MANSION,OFFICIAL COLONY,MAHARANI PETA, VIZAG 530002,.

PID No. : MED111551707 Register On : 25/03/2023 9:21 AM : 80026874 SID No. Collection On : 25/03/2023 9:55 AM Age / Sex : 38 Year(s) / Male Report On 25/03/2023 2:08 PM

: 01/04/2023 12:27 PM

Type : OP **Printed On**

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Alkaline Phosphatase (SAP) (Serum/IFCC AMP Buffer)	98	U/L	53 - 128
Total Protein (Serum/Biuret)	7.4	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.8	gm/dl	3.5 - 5.2
Globulin (Serum/Calculated)	3.60	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Calculated)	1.06		1.1 - 2.2
INTERPRETATION: Enclosure: Graph GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic) Lipid Profile	19	U/L	< 55
Cholesterol Total	205	mg/dL	Optimal: < 200
(Serum/Cholesterol oxidase/Peroxidase)	200	mg/dL	Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol-phosphate oxidase/Peroxidase)	129	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_ circulating level of triglycerides during most part of the day.







 PID No.
 : MED111551707
 Register On
 : 25/03/2023 9:21 AM

 SID No.
 : 80026874
 Collection On
 : 25/03/2023 9:55 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 25/03/2023 2:08 PM

Printed On : 01/04/2023 12:27 PM

Ref. Dr : MediWheel

: OP

Type



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
HDL Cholesterol (Serum/Immunoinhibition)	52	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	127.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	25.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	153.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol	3.9	Optimal: < 3.3
Ratio		Low Risk: 3.4 - 4.4
(Serum/Calculated)		Average Risk: 4.5 - 7.1
		Moderate Risk: 7.2 - 11.0
		High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 2.5 Optimal: < 2.5 (TG/HDL) Mild to moderate risk: 2.5 - 5.0 (Serum/Calculated) High Risk: > 5.0







 PID No.
 : MED111551707
 Register On
 : 25/03/2023 9:21 AM

 SID No.
 : 80026874
 Collection On
 : 25/03/2023 9:55 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 25/03/2023 2:08 PM

Type : OP Printed On : 01/04/2023 12:27 PM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.4		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC-Ion exchange)	5.4	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Mean Blood Glucose 108.28 mg/dl

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.46 ng/ml 0.7 - 2.04

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 7.83 μg/dl 4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay (CLIA))







APPROVED BY

The results pertain to sample tested.

Page 6 of 8

Lab Address: MEDALL HEALTH CARE PVT LTD,#17-11-3/4,DR.GKS MANSION,OFFICIAL COLONY,MAHARANI PETA, VIZAG 530002,.

 PID No.
 : MED111551707
 Register On
 : 25/03/2023 9:21 AM

 SID No.
 : 80026874
 Collection On
 : 25/03/2023 9:55 AM

Age / Sex : 38 Year(s) / Male **Report On** : 25/03/2023 2:08 PM

Ref. Dr : MediWheel

: OP

Printed On : 01/04/2023 12:27 PM



<u>Investigation</u>	<u>Observed</u> <u>U</u>	<u>nit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

INTERPRETATION:

Comment:

Type

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.16 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescence)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

Urine Analysis - Routine

Others

(Urine/Microscopy)

INTERPRETATION: Note: Done with Automated Urine Analyser & microscopy

Physical Examination(Urine Routine)

Colour PALE YELLOW Yellow to Amber

(Urine/Physical examination)

Appearance Clear Clear

(Urine/Physical examination)

Chemical Examination(Urine Routine)

Protein Negative Negative

(Urine/Dipstick-Error of indicator/ Sulphosalicylic acid method)









APPROVED BY

The results pertain to sample tested.

Page 7 of 8

Lab Address: MEDALL HEALTH CARE PVT LTD,#17-11-3/4,DR.GKS MANSION,OFFICIAL COLONY,MAHARANI PETA, VIZAG 530002,.

 PID No.
 : MED111551707
 Register On
 : 25/03/2023 9:21 AM

 SID No.
 : 80026874
 Collection On
 : 25/03/2023 9:55 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 25/03/2023 2:08 PM

Printed On : 01/04/2023 12:27 PM

Ref. Dr : MediWheel

: OP

Type



<u>Investigation</u>	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine/Dip Stick Method / Glucose Oxidase - Peroxidase / Benedict s semi quantitative method.)	Negative		Negative
Microscopic Examination(Urine Routine)			
Pus Cells (Urine/Microscopy exam of urine sediment)	3-4	/hpf	0 - 5
Epithelial Cells (Urine/Microscopy exam of urine sediment)	1-2	/hpf	NIL
RBCs (Urine/Microscopy exam of urine sediment)	NIL	/hpf	0 - 5







APPROVED BY

-- End of Report --