PID No.
 : MED120924617
 Register On
 : 26/03/2022 9:02 AM

 SID No.
 : 522212695
 Collection On
 : 26/03/2022 10:50 AM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 27/03/2022 3:12 PM

Printed On

: 30/03/2022 5:38 PM

MEDALL

Type : OP

Ref. Dr : MediWheel

Investigation HAEMATOLOGY	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	15.2	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	45.6	%	42 - 52
RBC Count (EDTA Blood/Impedance Variation)	5.15	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	89.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	29.6	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	33.4	g/dL	32 - 36
RDW-CV (Derived from Impedance)	12.7	%	11.5 - 16.0
RDW-SD (Derived from Impedance)	39.56	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	7100	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	56.5	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	23.6	%	20 - 45
Eosinophils	10.9	%	01 - 06



(Blood/Impedance Variation & Flow Cytometry)



: MediWheel

 PID No.
 : MED120924617
 Register On
 : 26/03/2022 9:02 AM

 SID No.
 : 522212695
 Collection On
 : 26/03/2022 10:50 AM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 27/03/2022 3:12 PM

Printed On

: 30/03/2022 5:38 PM

MEDALI

Type : OP

Ref. Dr

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood/Impedance Variation & Flow Cytometry)	8.4	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.6	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.01	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.68	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.77	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.60	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.04	10^3 / µl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	235	10^3 / μl	150 - 450
MPV (Blood/Derived from Impedance)	9.4	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.22	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	26	mm/hr	< 15





: MediWheel

 PID No.
 : MED120924617
 Register On
 : 26/03/2022 9:02 AM

 SID No.
 : 522212695
 Collection On
 : 26/03/2022 10:50 AM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 27/03/2022 3:12 PM

Printed On



Type : OP

Ref. Dr

Investigation BIOCHEMISTRY	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	1.3	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.5	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.8	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.6	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.7	gm/dL	3.5 - 5.2
Globulin (Serum/ <i>Derived</i>)	2.9	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.6		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	33	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	53	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	73	U/L	53 - 128
GGT(Gamma Glutamyl Transpeptidase) (Serum/SZASZ standarised IFCC)	84	U/L	< 55

: 30/03/2022 5:38 PM



VERIFIED BY



 PID No.
 : MED120924617
 Register On
 : 26/03/2022 9:02 AM

 SID No.
 : 522212695
 Collection On
 : 26/03/2022 10:50 AM

Report On

MEDALI

Type : OP

Ref. Dr : MediWheel

Age / Sex : 45 Year(s) / Male

Printed On : 30/03/2022 5:38 PM

27/03/2022 3:12 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	176	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	122	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.

part of the day.			
HDL Cholesterol (Serum/Immunoinhibition)	41	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	110.6	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	24.4	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	135.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



PID No. : MED120924617 Register On : 26/03/2022 9:02 AM : 522212695 SID No. Collection On : 26/03/2022 10:50 AM

Report On



Type : OP

Printed On : 30/03/2022 5:38 PM

Ref. Dr : MediWheel

Age / Sex : 45 Year(s) / Male

Investigation <u>Observed</u> **Unit Biological Value** Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

27/03/2022 3:12 PM

Total Cholesterol/HDL Cholesterol Ratio 4.3

(Serum/Calculated)

Optimal: < 3.3Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 3

(TG/HDL)

(Serum/Calculated)

High Risk: > 5.0

LDL/HDL Cholesterol Ratio 2.7 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated)

High Risk: > 6.0

Optimal: < 2.5Mild to moderate risk: 2.5 - 5.0



VERIFIED BY



 SID No.
 : 522212695
 Collection On
 : 26/03/2022 10:50 AM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 27/03/2022 3:12 PM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	6.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 134.11 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Un	<u>Biological</u>
	Value	Reference Interval

: 30/03/2022 5:38 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.48 ng/mL 0.7 - 2.04

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 12.37 μg/dL 4.2 - 12.0

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.87 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Microparticle

Immuno as say (CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



 PID No.
 : MED120924617
 Register On
 : 26/03/2022 9:02 AM

 SID No.
 : 522212695
 Collection On
 : 26/03/2022 10:50 AM

 Age / Sex
 : 45 Year(s) / Male
 Report On
 : 27/03/2022 3:12 PM

Printed On



Type : OP

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

Pale yellow

: 30/03/2022 5:38 PM

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

(Urine)	J		
Appearance (Urine)	Clear		Clear
Volume	15	mL	
(Urine)			
CHEMICAL EXAMINATION(Automated-			

<u>Urineanalyser)</u>

Colour

pH (Urine/AUTOMATED URINANALYSER)	6.0	4.5 - 8.0
Specific Gravity (Urine)	1.015	1.002 - 1.035
Ketones (Urine)	Negative	Negative
Urobilinogen (Urine/AUTOMATED URINANALYSER)	0.2	0.2 - 1.0
Blood (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Nitrite (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Bilirubin (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Protein	Negative	Negative



(Urine)

VERIFIED BY



: MediWheel

PID No. : MED120924617 Register On : 26/03/2022 9:02 AM : 522212695 **Collection On** : 26/03/2022 10:50 AM SID No. Age / Sex : 45 Year(s) / Male Report On : 27/03/2022 3:12 PM



Type : OP

Ref. Dr

Printed On : 30/03/2022 5:38 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine)	Negative		Negative
Leukocytes (Urine) MICROSCOPY(URINE DEPOSITS)	Negative	leuco/uL	Negative
Pus Cells (Urine/Flow cytometry)	2-4	/hpf	3-5
Epithelial Cells (Urine)	0-1	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	NIL
Others (Urine)	Nil		Nil



VERIFIED BY



Type : OP

Ref. Dr : MediWheel



InvestigationObservedUnitBiologicalValueReference Interval

: 30/03/2022 5:38 PM

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'AB' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.

Printed On



VERIFIED BY



PID No. Register On : MED120924617 : 522212695 SID No.

Age / Sex : 45 Year(s) / Male Report On 27/03/2022 3:12 PM

Printed On

Type : OP

Ref. Dr : MediWheel

: 26/03/2022 9:02 AM Collection On : 26/03/2022 10:50 AM

: 30/03/2022 5:38 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	10		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	100	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Negative Glucose Fasting - Urine Negative (Urine - F) Glucose Postprandial (PPBS) 143 mg/dL 70 - 140 (Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN) mg/dL (Serum/Urease-GLDH) 0.8 0.9 - 1.3Creatinine mg/dL

(Serum/Jaffe Kinetic)

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine

Uric Acid 6.2 3.5 - 7.2mg/dL

(Serum/Uricase/Peroxidase)



VERIFIED BY



 PID No.
 : MED120924617
 Register On
 : 26/03/2022 9:02 AM

 SID No.
 : 522212695
 Collection On
 : 26/03/2022 10:50 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Prostate specific antigen - Total(PSA) (Serum/Chemiluminescent Microparticle Immunoassay(CMIA))	1.12	ng/mL	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0

: 30/03/2022 5:38 PM



VERIFIED BY



APPROVED BY

-- End of Report --

Name	RAKESH KUMAR	ID	MED120924617
Age & Gender	45/Male	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (12.8 cms) and **shows diffusely increased echopattern.** No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER show normal shape and has clear contents. Wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size (10.2 cms) and echopattern.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cm)	Parenchymal thickness (cm)
Right Kidney	10.5	1.7
Left Kidney	10.7	2.0

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern. It measures 3.4 x 2.8 x 3.2 cm, volume: 15.6cc.

Small 5.5 mm midline cyst is noted.

No evidence of ascites.

IMPRESSION:

- Grade II fatty liver.
- No other significant abnormality detected.

Name	RAKESH KUMAR	ID	MED120924617
Age & Gender	45/Male	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



DR. H.K. ANAND DR. C.R RAMACHANDRA DR. LOHITH H.P DR. VARSHA KALE CONSULTANT RADIOLOGISTS

Lh/Ss

Name	RAKESH KUMAR	ID	MED120924617
Age & Gender	45/Male		26-03-2022 00:00:00
Ref Doctor Name	MediWheel	-	



2D ECHOCARDIOGRAPHIC STUDY

M-mode measurement:

AORTA 2.70 cms. LEFT ATRIUM 3.24 cms. AVS 1.48 cms. LEFT VENTRICLE (DIASTOLE) 4.55 cms. (SYSTOLE) 2.28 cms. **VENTRICULAR SEPTUM** (DIASTOLE) 0.96 cms. (SYSTOLE) 1.47 cms. POSTERIOR WALL (DIASTOLE) 1.00 cms. (SYSTOLE) 1.39 cms. **EDV** 95 ml. **ESV** 17 ml. FRACTIONAL SHORTENING 30 %

EJECTION FRACTION EPSS cms. **RVID** 1.80 cms.

DOPPLER MEASUREMENTS:

MITRAL VALVE: E - 0.9 m/sA - 0.8 m/sNO MR.

60

%

AORTIC VALVE: 1.0 m/sNO AR.

TRICUSPID VALVE: E - 0.5 m/s A - 0.4 m/s NO TR.

PULMONARY VALVE: 0.8 m/sNO PR.

Name	RAKESH KUMAR	ID	MED120924617
Age & Gender	45/Male		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHY FINDINGS:

Left Ventricle : Normal size, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal.Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- NORMAL SIZED CARDIAC CHAMBERS.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

Name	RAKESH KUMAR	ID	MED120924617
Age & Gender	45/Male	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



DR. ANAND KUMAR M MD DM CONSULTANT INTERVENTIONAL CARDIOLOGIST

Name	RAKESH KUMAR	ID	MED120924617
Age & Gender	45/Male		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

• No significant abnormality detected.

	Name	RAKESH KUMAR	ID	MED120924617
	Age & Gender	45/Male	Visit Date	26-03-2022 00:00:00
I.	Ref Doctor Name	MediWheel		



Test completed