

General Physical Examination

Date of Examination: 12/03/2023

Name: DHARMENDRA K. Goyal Age: 54 Sex: Male

DOB: 22/05/2023

Referred By: BOB

Photo ID: UD ID #: attached

Ht: 181 (cm) Wt: 72 (Kg)

Chest (Expiration): 85 (cm) Abdomen Circumference: 85 (cm)

Blood Pressure: 138/88 mm Hg PR: 73 / min RR: 16. / min Temp: Afebric

BMI 22.0.

Eye Examination: Vision using Spectacles both eye

Partial Colour blindness present

Other: Not significant.

On examination he/she appears physically and mentally fit: Yes / No

Signature Of Examinee : [Signature]

Name of Examinee: _____

Signature Medical Examiner : _____

Name Medical Examiner _____

Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
RMC Reg. No. 017996

9462695849



भारत सरकार



आधार

भारतीय विशिष्ट पहचान प्राधिकरण

भारत सरकार
Unique Identification Authority of India
Government of India

नामांकन क्रम / Enrollment No 1050/08393/04684

To,
धर्मेन्द्र कुमार गोयल
Dharmendra Kumar Goyal
S/O: Trilok Prakash Goyal
12/04/2013
112
bhartendu nagar
khatipura
Jaipur
Jhotwara A C Jobner Jaipur
Rajasthan 302012
9462695849

Ref: 275 / 13G / 164098 / 164814 / P



SH020079764FT



आपका आधार क्रमांक / Your Aadhaar No. :

9775 5176 7341

आधार - आम आदमी का अधिकार



भारत सरकार

Government of India



धर्मेन्द्र कुमार गोयल
Dharmendra Kumar Goyal
जन्म वर्ष / Year of Birth : 1968
पुरुष / Male




9775 5176 7341

आधार - आम आदमी का अधिकार


Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
FMC Reg. No.-017996



DR. GOYAL PATH LAB & IMAGING CENTER, JAIPUR
 4022 / MR. DHARMENDRA K GOYAL / 54 Yrs / M/ Non Smoker
 Heart Rate : 66 bpm / Tested On : 12-Mar-23 10:25:31 / HF 0.05 Hz - LF 35 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s
 / Refd By: BOB MEDIWHEEL



Vent Rate : 66 bpm
 PR Interval : 156 ms
 QRS Duration: 78 ms
 QT/QTc Int : 370/381 ms
 P-QRS-T axis: 76.00° 33.00° 56.00°



Turn

Mohanka
 33703
 RADIO (REGD. RTG)

Reported By: M. (RCCP-JKD)

Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
Sex / Age :- Male 54 Yrs 9 Mon 21 Days
Company :- MediWHEEL

Patient ID :- 122229977
Ref. By Dr:- BOB
Lab/Hosp :-



Sample Type :- EDTA

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 12:50:13

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
HAEMOGARAM			
HAEMOGLOBIN (Hb)	16.0	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	7.00	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	71.7	%	40.0 - 80.0
LYMPHOCYTE	24.0	%	20.0 - 40.0
EOSINOPHIL	1.4	%	1.0 - 6.0
MONOCYTE	2.6	%	2.0 - 10.0
BASOPHIL	0.3	%	0.0 - 2.0
NEUT#	5.02	10 ³ /uL	1.50 - 7.00
LYMPH#	1.69	10 ³ /uL	1.00 - 3.70
EO#	0.09	10 ³ /uL	0.00 - 0.40
MONO#	0.18	10 ³ /uL	0.00 - 0.70
BASO#	0.02	10 ³ /uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	5.38	x10 ⁶ /uL	4.50 - 5.50
HEMATOCRIT (HCT)	47.00	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	87.4	fL	83.0 - 101.0
MEAN CORP HB (MCH)	29.8	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	34.1	g/dL	31.5 - 34.5
PLATELET COUNT	275	x10 ³ /uL	150 - 410
RDW-CV	13.2	%	11.6 - 14.0
MENTZER INDEX	16.25		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

AJAYSINGH
Technologist

Page No: 2 of 13



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37

Patient ID :-122229977

NAME :- Mr. DHARMENDRA K GOYAL

Ref. By Dr:- BOB

Sex / Age :- Male 54 Yrs 9 Mon 21 Days

Lab/Hosp :-

Company :- MediWheel



Sample Type :- EDTA

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 12:50:13

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
Erythrocyte Sedimentation Rate (ESR)	28 H	mm/hr.	00 - 13

(ESR) Methodology : Measurement of ESR by cells aggregation.

Instrument Name : Independent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation : ESR test is a non-specific indicator of inflammatory disease and abnormal protein states.

The test is used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction)

Levels are higher in pregnancy due to hyperfibrinogenaemia.

The "3-figure ESR" $\times > 100$ value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia or connective tissue disease.

(CBC): Methodology: TLC, DLC, Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance. and MCH, MCV, MCHC, MENTZER INDEX are calculated. Instrument Name: Sysmex 6 part fully automatic analyzer XN-L, Japan

AJAYSINGH
Technologist

Page No: 3 of 13



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37

NAME :- Mr. DHARMENDRA K GOYAL

Sex / Age :- Male 54 Yrs 9 Mon 21 Days

Company :- MediWheel

Patient ID :-122229977

Ref. By Dr:- BOB

Lab/Hosp :-



Sample Type :- EDTA

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 12:50:13

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BOB PACKAGE ABOVE 40MALE GLYCOSYLATED HEMOGLOBIN (HbA1C) Method:- HPLC	6.3	H %	Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0 Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose over the period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasma glucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHb depends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to the mean of HbA1C. Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1C measurements. The effects vary depending on the specific Hb variant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

Method:- Calculated Parameter

134 H mg/dL

Non Diabetic < 100 mg/dL
 Prediabetic 100- 125 mg/dL
 Diabetic 126 mg/dL or Higher

AJAYSINGH
 Technologist

Page No: 1 of 13



Dr. Chandrika Gupta
 MBBS.MD (Path)
 RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
 Sex / Age :- Male 54 Yrs 9 Mon 21 Days
 Company :- MediWheel

Patient ID :-122229977
 Ref. By Dr:- BOB
 Lab/Hosp :-



Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 16:10:09

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	161.46	mg/dl	Desirable <200 Borderline 200-239 High > 240
TRIGLYCERIDES Method:- GPO-PAP	210.82 H	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	31.35	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	94.97	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	42.16	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	5.15 H		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	3.03		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED	594.78	mg/dl	400.00 - 1000.00
TOTAL CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders			
TRIGLYCERIDES InstrumentName:Randox Rx Imola Interpretation : Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.			
DIRECT HDLCHOLESTERO InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.			
DIRECT LDL-CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.			
TOTAL LIPID AND VLDL ARE CALCULATED			

MUKESH SINGH

Page No: 4 of 13



Dr. Chandrika Gupta
 MBBS.MD (Path)
 RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
 Sex / Age :- Male 54 Yrs 9 Mon 21 Days
 Company :- MediWHEEL

Patient ID :- 122229977
 Ref. By Dr:- BOB
 Lab/Hosp :-



Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 16:10:09

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.52	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12 1 month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2 Ref-(ACCP 2020)
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.19	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.33	mg/dl	0.30-0.70
SGOT Method:- IFCC	20.2	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	31.0	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	76.10	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC	29.40	U/L	11.00 - 50.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	6.97	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.35	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	2.62	gm/dl	2.20 - 3.50
A/G RATIO	1.66		1.30 - 2.50

Total Bilirubin Methodology: Colorimetric method InstrumentName Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCC InstrumentName Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

Alkaline Phosphatase Methodology: AMP Buffer InstrumentName Randox Rx Imola Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobiliary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology: Biuret Reagent InstrumentName Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra- or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal).

MUKESH SINGH

Page No: 5 of 13



Dr. Chandrika Gupta
 MBBS.MD (Path)
 RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
 Sex / Age :- Male 54 Yrs 9 Mon 21 Days
 Company :- MediWheel

Patient ID :-122229977
 Ref. By Dr:- BOB
 Lab/Hosp :-



Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 12:08:22

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.320	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	9.260	ug/dl	5.530 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	2.428	μIU/mL	0.400 - 4.649

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

MUKESH SINGH
Technologist

Page No: 6 of 13



Dr. Chandrika Gupta
 MBBS.MD (Path)
 RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
 Sex / Age :- Male 54 Yrs 9 Mon 21 Days
 Company :- MediWheel

Patient ID :-122229977
 Ref. By Dr:- BOB
 Lab/Hosp :-



Sample Type :- URINE

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 15:49:55

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH) Method:- Reagent Strip(Double indicator blue reaction)	5.5		5.0 - 7.5
SPECIFIC GRAVITY Method:- Reagent Strip(bromthymol blue)	1.015		1.010 - 1.030
PROTEIN Method:- Reagent Strip (Sulphosalicylic acid test)	NIL		NIL
GLUCOSE Method:- Reagent Strip (Glu.Oxidase Peroxidase Benedict)	NIL		NIL
BILIRUBIN Method:- Reagent Strip (Azo-coupling reaction)	NEGATIVE		NEGATIVE
UROBILINOGEN Method:- Reagent Strip (Modified chrlich reaction)	NORMAL		NORMAL
KETONES Method:- Reagent Strip (Sodium Nitropruside) Rothera's	NEGATIVE		NEGATIVE
NITRITE Method:- Reagent Strip (Diazotization reaction)	NEGATIVE		NEGATIVE
RBC Method:- Reagent Strip (Peroxidase like activity)	NIL		NIL
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	1-2	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT

VIJENDRAMEENA
Technologist

Page No: 7 of 13




Dr. Chandrika Gupta
 MBBS.MD (Path)
 RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur
Tele: 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com

Date :- 12/03/2023 09:24:37

NAME :- Mr. DHARMENDRA K GOYAL

Sex / Age :- Male 54 Yrs 9 Mon 21 Days

Company :- MediWheel

Patient ID :-122229977

Ref. By Dr:- BOB

Lab/Hosp :-



Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 16:10:09

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
SERUM CREATININE Method:- Colorimetric Method	1.30	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20
SERUM URIC ACID Method:- Enzymatic colorimetric	7.93 H	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7

MUKESH SINGH

Page No: 9 of 13



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37

Patient ID :-122229977

NAME :- Mr. DHARMENDRA K GOYAL

Ref. By Dr:- BOB

Sex / Age :- Male 54 Yrs 9 Mon 21 Days

Lab/Hosp :-

Company :- MediWheel



Sample Type :- EDTA, URINE

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 15:49:55

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BLOOD GROUP ABO	"B" POSITIVE		
BLOOD GROUP ABO Methodology : Haemagglutination reaction Kit Name : Monoclonal agglutinating antibodies (Span clone).			
URINE SUGAR (FASTING) Collected Sample Received	Nil		Nil

AJAYSINGH, VIJENDRAMEENA
Technologist

Page No: 11 of 13



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganeer Road, Jaipur
Tele: 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com

Date :- 12/03/2023 09:24:37

Patient ID :- 122229977

NAME :- Mr. DHARMENDRA K GOYAL

Ref. By Dr:- BOB

Sex / Age :- Male 54 Yrs 9 Mon 21 Days

Lab/Hosp :-

Company :- MediWheel



Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 16:10:09

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
BLOOD UREA NITROGEN (BUN)	10.2	mg/dl	0.0 - 23.0

MUKESH SINGH

Page No: 12 of 13



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037

Date :- 12/03/2023 09:24:37

Patient ID :-122229977



NAME :- Mr. DHARMENDRA K GOYAL

Ref. By Dr:- BOB

Sex / Age :- Male 54 Yrs 9 Mon 21 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2023 10:39:09

Final Authentication : 12/03/2023 12:08:22

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
TOTAL PSA Method:- Chemiluminescence	2.270	ng/ml	0.000 - 4.000

InstrumentName: VITROS ECI **Interpretation :** Elevated serum PSA concentrations are found in men with prostate cancer, benign prostatic hypertrophy (BHP) or inflammatory conditions of other adjacent genitourinary tissues, but not in apparently healthy men or in men with cancers other than prostate cancer. PSA has been demonstrated to be an accurate marker for monitoring advancing clinical stage in untreated patients and for monitoring response to therapy by radical prostatectomy, radiation therapy and anti-androgen therapy. PSA is also important in determining the potential and actual effectiveness of surgery or other therapies. Progressive disease is defined by an increase of at least 25%. Sampling should be repeated within two to four weeks for additional evidence. Different assay methods cannot be used interchangeably.

*** End of Report ***

MUKESH SINGH
Technologist

Page No: 13 of 13



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
Sex / Age :- Male 54 Yrs 9 Mon 21 Days
Company :- MediWheel

Patient ID :- 122229977
Ref. By Doctor:-BOB
Lab/Hosp :-

Final Authentication : 12/03/2023 12:54:10

BOB PACKAGE ABOVE 40MALE

USG WHOLE ABDOMEN

Liver is mild enlarged in size (15.7 cm). Echo-texture is minimal bright. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation. **Small calculus of size ~4.2 mm is seen in lower calyx of right kidney and ~4.8 mm is seen in lower calyx of left kidney.**

Urinary bladder is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Prostate is mild enlarged in size (33 gms) with normal echo-texture and outline.

No enlarged nodes are visualised. No retro-peritoneal lesion is identified.

No significant free fluid is seen in peritoneal cavity.

IMPRESSION:

- * Mild hepatomegaly with early fatty changes.
- * Small bilateral renal calculus.
- * Mild prostatomegaly.

Needs clinical correlation for further evaluation

Page No: 1 of 1

*** End of Report ***

ANITASHARMA

Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
RMC Reg No. 017996

Dr. Poonam Gupta
MBBS, MD (Radio Diagnosis)
RMC No. 32495

Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant
FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain
MBBS, DNB, (Radio-Diagnosis)
RMC No. 21687

Transcript by.

Print Copy



Date :- 12/03/2023 09:24:37	Patient ID :-122229977
NAME :- Mr. DHARMENDRA K GOYAL	Ref. By Doctor:-BOB
Sex / Age :- Male 54 Yrs 9 Mon 21 Days	Lab/Hosp :-
Company :- MediWheel	

Final Authentication : 12/03/2023 12:53:15

BOB PACKAGE ABOVE 40MALE
 2D ECHO OPTION TMT (ADULT/CHILD)

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	NORMAL	TRICUSPID VALVE	NORMAL
AORTIC VALVE	NORMAL	PULMONARY VALVE	NORMAL

M.MODE EXAMINATION:

AO	23	mm	LA	29	Mm	IVS-D	8	mm
IVS-S	14	mm	LVID	35	Mm	LVSD	23	mm
LVPW-D	08	mm	LVPW-S	14	Mm	RV		mm
RVWT		mm	EDV		MI	LVVS		ml
LVEF	66%		RWMA		ABSENT			

CHAMBERS:

LA	NORMAL	RA	NORMAL
LV	NORMAL	RV	NORMAL
PERICARDIUM	NORMAL		

COLOUR DOPPLER:

MITRAL VALVE					
E VELOCITY	0.90	m/sec	PEAK GRADIENT		Mm/hg
A VELOCITY	0.75	m/sec	MEAN GRADIENT		Mm/hg
MVA BY PHT		Cm2	MVA BY PLANIMETRY		Cm2
MITRAL REGURGITATION			ABSENT		
AORTIC VALVE					
PEAK VELOCITY	1.25	m/sec	PEAK GRADIENT		mm/hg
AR VMAX		m/sec	MEAN GRADIENT		mm/hg
AORTIC REGURGITATION			ABSENT		
TRICUSPID VALVE					
PEAK VELOCITY	0.36	m/sec	PEAK GRADIENT		mm/hg
MEAN VELOCITY		m/sec	MEAN GRADIENT		mm/hg
VM _{max} VELOCITY					
TRICUSPID REGURGITATION			ABSENT		
PULMONARY VALVE					
PEAK VELOCITY	0.90	M/sec.	PEAK GRADIENT		Mm/hg
MEAN VELOCITY			MEAN GRADIENT		Mm/hg
PULMONARY REGURGITATION			ABSENT		

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
Sex / Age :- Male 54 Yrs 9 Mon 21 Days
Company :- MediWheel

Patient ID :-122229977
Ref. By Doctor:-BOB
Lab/Hosp :-

Final Authentication : 12/03/2023 12:53:15

Impression--

1. Normal LV size & contractility.
2. No RWMA, LVEF 66%.
3. Normal cardiac chamber.
4. Normal valve.
5. No clot, no vegetation, no pericardial effusion.

(Cardiologist)

*** End of Report ***

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalpathlab.com | E-mail : drgoyalpiyush@gmail.com



Date :- 12/03/2023 09:24:37
NAME :- Mr. DHARMENDRA K GOYAL
Sex / Age :- Male 54 Yrs 9 Mon 21 Days
Company :- MediWheel

Patient ID :- 12222977
Ref. By Doctor:-BOB
Lab/Hosp :-

Final Authentication : 12/03/2023 12:06:29

BOB PACKAGE ABOVE 40MALE

X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression :- Normal Study

(Please correlate clinically and with relevant further investigations)

*** End of Report ***

Page No: 1 of 1

Dr. Piyush Goyal
(D.M.R.D.) BILAL

Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
RMC Reg No. 017996

Dr. Poonam Gupta
MBBS, MD (Radio Diagnosis)
RMC No. 32495

Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant

FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain
MBBS, DNB, (Radio-Diagnosis)
RMC No. 21687

Transcript by.

Dr Goyal's Path Lab, Jaipur

Name : DHARMENDRA K GOYAL

12 Mar 2023

