

From, DDRE SAL Kannue

To, Medruheel

Respected Six/Madam,

Please note itoday (08.10.2022), me have a client

named as Vegethar G Pillai from Mednucheel (Accession

no-4053V J000 652). Due to emergency leave of our optometor

me are mable to sprocess ophthal But the client is not

mulling to some another day. So, Kinolly do needful.



wh.





CLIENT CODE: CA00010147 **CLIENT'S NAME AND ADDRESS:**

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED

F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA

8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

ACCESSION NO: 4053VJ000652 AGE: 29 Years SEX: Female ABHA NO:

RECEIVED: 08/10/2022 08:33 08/10/2022 13:49 DRAWN: REPORTED:

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

MEDIWHEEL HEALTH CHECKUP BELOW 40(F)TMT

TREADMILL TEST

COMPLETED TREADMILL TEST

OPTHAL

COMPLETED **OPTHAL**

PHYSICAL EXAMINATION

COMPLETED PHYSICAL EXAMINATION







MEDICAL EXAMINATION REPORT (MER)

If the examinee is suffering from an acute life threatening situation, you may be obliged to disclose the result of the medical examination to the examinee.

 Name of the examinee Mark of Identification Age/Date of Birth Photo ID Checked 	: Mr/Mrs/Ms. mvs Gelle G. Pillar : (Mole/Scar/any other (specify location)): make below right ey : 29, 06-10-1993 Gender: F/M : (Passport/Election Card/PAN Card/Driving Licence/Company ID) Acidl
---	---

PHYSICAL DETAILS:

a. Height 16.5 (cms)	b. Weight (Kgs)	c. Girth of Abdome	en
d. Pulse Rate	e. Blood Pressure:	Systolic E	Diastolic
	1" Reading	110	70'
	2 nd Reading	110	70

FAMILY HISTORY:

Relation	Age if Living	Health Status	If deceased, age at the time and cause
Father	65	Healthy	
Mother	50	4)	
Brother(s)	21	y	
Sister(s)		HOUSE ME TO THE	PROBLEM STREET

HABITS & ADDICTIONS: Does the examinee consume any of the following?

Tobacco in any form	Sedative	Alcohol
1 10	<u>~0</u>	NO!

PERSONAL HISTORY

- a. Are you presently in good health and entirely free from any mental or Physical impairment or deformity.
 If No, please attach details.
- b. Have you undergone/been advised any surgical procedure?
- c. During the last 5 years have you been medically examined, received any advice or treatment or admitted to any hospital?
- d. Have you lost or gained weight in past 12 months?

Have you ever suffered from any of the following?

- Psychological Disorders or any kind of disorders of the Nervous System?
- · Any disorders of Respiratory system?
- Any Cardiac or Circulatory Disorders?
- · Enlarged glands or any form of Cancer/Tumour?
- · Any Musculoskeletal disorder?

- Any disorder of Gastrointestinal System?
- Unexplained recurrent or persistent fever, and/or weight loss
- Have you been tested for HIV/HBsAg / HCV before? If yes attach reports
- Are you presently taking medication of any kind?







Corp. Office: DDRC SRL Tower, G- 131, Panampilly Nagar, Ernakulam - 682 036 Ph No. 0484-2318223, 2318222, e-mail: info@ddrcsrl.com, web: www.ddrcsrl.com





ma/dl

CLIENT CODE: CA00010147
CLIENT'S NAME AND ADDRESS:

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED

F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156 DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

ACCESSION NO: 4053VJ000652 AGE: 29 Years SEX: Female ABHA NO:

DRAWN: RECEIVED: 08/10/2022 08:33 REPORTED: 08/10/2022 13:49

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

5

Test Report Status Final Results Units

MEDIWHEEL HEALTH CHECKUP BELOW 40(F)TMT

SFRIIM	RI OOD	IIRFA	NITROGEN

BLOOD LIDEA NITDOGEN

BEOOD ONEA WITHOUGH	_		3,
BUN/CREAT RATIO			
BUN/CREAT RATIO	7.14	5.00 - 15.00	
CREATININE, SERUM			
CREATININE	0.70	0.60 - 1.1	mg/dL

CREATININE	0.70	0.00 - 1.1	IIIg/uL
GLUCOSE, POST-PRANDIAL, PLASMA			

GLUCOSE, POST-PRANDIAL, PLASMA	69	Diabetes Mellitus : > or = 200 mg/dL
		ma/dl

Impaired Glucose tolerance/ Prediabetes: 140 to 199 mg/dL. Hypoglycemia: < 55 mg/dL.

GLUCOSE, FASTING, PLASMA

GLUCOSE, FASTING, PLASMA 79 Diabetes Mellitus : > or = 126 mg/dL

mg/dL.

Low 6 - 20

Impaired fasting Glucose/ Prediabetes: 101 to 125 mg/dL. Hypoglycemia: < 55 mg/dL.

GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD

GLYCOSYLATED HEMOGLOBIN (HBA1C) 5.4 Normal: 4.0 - 5.6 %. % Non-diabetic level: < 5.7%.

Non-diabetic level: < 5.7%.

More stringent goal: < 6.5 %.

General goal: < 7%.

Less stringent goal: < 8%.

Glycemic targets in CKD:
If eGFR > 60: < 7%.

If eGFR < 60: 7 - 8.5%.

CORONARY RISK PROFILE (LIPID PROFILE), SERUM

CHOLESTEROL 167 < 200 Desirable mg/dL 200 - 239 Borderline High

>/= 240 High
TRIGLYCERIDES 103 Normal : < 150 mg/dL

High: 150-199 Hypertriglyceridemia: 200-499

Very High: > 499

HDL CHOLESTEROL

38 Low 40 - 60 mg/dL

DIRECT LDL CHOLESTEROL 112 High < 100 Optimal mg/dL 100 - 129 Near or above optimal

130 - 159 Borderline High

160 - 189 High >/= 190 Very High





Page 2 Of 9

Any disorders of Urinary System?	YN	Any disorder of the Eyes, Ears, Nose, Throat or Mouth & Skin	
FOR FEMALE CANDIDATES ONLY			
a. Is there any history of diseases of breast/genital organs?	YN	d. Do you have any history of miscarriage/ abortion or MTP	YN
b. Is there any history of abnormal PAP Smear/Mammogram/USG of Pelvis or any other tests? (If yes attach reports)	YN	 e. For Parous Women, were there any complication during pregnancy such as gestational diabetes, hypertension etc 	YN
c. Do you suspect any disease of Uterus, Cervix or Ovaries?	YA	f. Are you now pregnant? If yes, how many month	s?
CONFIDENTAIL COMMENTS FROM MEDICA	AL EXA	MINER	V N
➤ Was the examinee co-operative?			
Is there anything about the examine's health, life his/her job?	estyle tha	t might affect him/her in the near future with regard	YN
> Are there any points on which you suggest furth	er inforn	nation be obtained?	YIN
 Based on your clinical impression, please provide 			
> Do you think he/she is MEDICALLY FIT or UN	NFIT for	employment.	
medically	1 4	it	
MEDICAL EXAMINER'S DECLARATION			

I hereby confirm that I have examined the above individual after verification of his/her identity and the findings stated above are true and correct to the best of my knowledge.

Name & Signature of the Medical Examiner

Dr. INDUSARATH.S, MBBS,MD,DNB

Regd. No: 41964

DDRC SRL, KANNUR

Name & Seal of DDRC SRL Branch

Seal of Medical Examiner

Date & Time

DDRC SRL Diagnostics Private Limited





CLIENT CODE: CA00010147 **CLIENT'S NAME AND ADDRESS:**

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED

F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030

DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

ACCESSION NO: 4053VJ000652 AGE: 29 Years SEX : Female ABHA NO:

RECEIVED: 08/10/2022 08:33 08/10/2022 13:49 DRAWN: REPORTED:

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Units
NON HDL CHOLESTEROL	129	Desirable-Less than 130 Above Desirable-130-159 Borderline High-160-189 High-190-219 Very High- >or =220	mg/dL
CHOL/HDL RATIO	4.4	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk	
LDL/HDL RATIO	3.0	0.5-3 Desirable/Low risk 3.1-6 Borderline/Moderate risk >6.0 High Risk	
VERY LOW DENSITY LIPOPROTEIN LIVER FUNCTION TEST WITH GGT	20.5	= 30</td <td>mg/dL</td>	mg/dL
BILIRUBIN, TOTAL	0.50	Upto 1.2	mg/dL
BILIRUBIN, DIRECT	0.14	< 0.31	mg/dL
BILIRUBIN, INDIRECT	0.36	0.00 - 0.60	mg/dL
TOTAL PROTEIN	7.5	Ambulatory: 6.4 - 8.3 Recumbant: 6 - 7.8	g/dL
ALBUMIN	4.5	3.5 - 5.2	g/dL
GLOBULIN	3.0	2.0 - 4.0	g/dL
ALBUMIN/GLOBULIN RATIO	1.5	1.0 - 2.0	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	28	< 33	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT)	36 Hig	gh < 34	U/L
ALKALINE PHOSPHATASE	67	35 - 105	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT) TOTAL PROTEIN, SERUM	19	< 40	U/L
TOTAL PROTEIN	7.5	Ambulatory : 6.4 - 8.3 Recumbant : 6 - 7.8	g/dL
URIC ACID, SERUM			
URIC ACID ABO GROUP & RH TYPE, EDTA WHOLE BLOOD	4.5	2.4 - 5.7	mg/dL
ABO GROUP	TYPE O		
RH TYPE	POSITIVE		
BLOOD COUNTS			
HEMOGLOBIN	12.2	12.0 - 15.0	g/dL
RED BLOOD CELL COUNT	4.24	3.8 - 4.8	mil/μL





Scan to View Report







CLIENT CODE: CA00010147
CLIENT'S NAME AND ADDRESS:

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED

F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156 DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

ACCESSION NO: 4053VJ000652 AGE: 29 Years SEX: Female ABHA NO:

DRAWN: RECEIVED: 08/10/2022 08:33 REPORTED: 08/10/2022 13:49

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results			Units
			4.0.40.0	
WHITE BLOOD CELL COUNT	6.10		4.0 - 10.0	thou/µL
PLATELET COUNT	217		150 - 410	thou/µL
RBC AND PLATELET INDICES	25.7	1	26 46	0/
HEMATOCRIT	35.7	LOW	36 - 46	%
MEAN CORPUSCULAR VOL	84.2		83 - 101	fL
MEAN CORPUSCULAR HGB.	28.8		27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION	34.2		31.5 - 34.5	g/dL
MEAN PLATELET VOLUME	9.8		6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT - NLR	5.0			
SEGMENTED NEUTROPHILS	46		40 - 80	%
ABSOLUTE NEUTROPHIL COUNT	2.81		2.0 - 7.0	thou/µL
LYMPHOCYTES	48	High	20 - 40	%
ABSOLUTE LYMPHOCYTE COUNT	2.93		1 - 3	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	1.0			
EOSINOPHILS	3		1 - 6	%
ABSOLUTE EOSINOPHIL COUNT	0.18		0.02 - 0.50	thou/µL
MONOCYTES	2		2 - 10	%
ABSOLUTE MONOCYTE COUNT	0.12	Low	0.20 - 1.00	thou/µL
BASOPHILS	1		0 - 2	%
ERYTHRO SEDIMENTATION RATE, BLOOD				
SEDIMENTATION RATE (ESR)	12		0 - 20	mm at 1 hr
STOOL: OVA & PARASITE				
COLOUR	BROWN			
CONSISTENCY	HARD			
ODOUR	FAECAL			
MUCUS	NOT DETECTED		NOT DETECTED	
VISIBLE BLOOD	ABSENT		ABSENT	
POLYMORPHONUCLEAR LEUKOCYTES	1-2		0 - 5	/HPF
RED BLOOD CELLS	NOT DETECTED		NOT DETECTED	/HPF
OVA	NOT SEEN			
OCCULT BLOOD	NOT DETECTED		NOT DETECTED	
SUGAR URINE - POST PRANDIAL				
SUGAR URINE - POST PRANDIAL	NOT DETECTED		NOT DETECTED	
THYROID PANEL, SERUM				





V6 V6	
IR IR	M CM
ID: GEETHU G PILLAI Female 29Years cm kg Regd No: 41964 Y O O O O C C C Standard C C C C C Standard C C C C C Standard C C C C C C Standard C C C C C C C Standard C C C C C C C C C C C C C Standard C C C C C C C C C C C C C C C C C C C	ID: GEETHU G PILLAI Female 29Years cm kg Regd. No: 41964 DDRC SRL, KANNU TO Oma (ECC) Suma Lumin HR 81 bpm P 100 ms P 100 ms PR 155 ms QRS QRS QRS GRS FQRS/T : 5442,16 ° RV5/SV1 : 1,113/0.620 mV





CLIENT CODE: CA00010147 **CLIENT'S NAME AND ADDRESS:**

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED

F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

KANNUR KERALA, INDIA

Tel: 93334 93334 Email: customercare.ddrc@srl.in

DDRC SRL DIAGNOSTICS

PATIENT NAME: GEETHU G PILLAI

GEETF0810934053

ACCESSION NO: 4053VJ000652

AGE: 29 Years

SEX: Female ABHA NO:

DRAWN:

RECEIVED: 08/10/2022 08:33

08/10/2022 13:49 REPORTED:

REFERRING DOCTOR: SELF

CLIENT PATIENT ID:

PATIENT ID:

Test Report Status <u>Final</u>	Results		Units
Т3	135.50	80.00 - 200.00	ng/dL
T4	8.53	5.10 - 14.10	μg/dl
TSH 3RD GENERATION	4.770 High	Non-Pregnant: 0.4-4.2	μIU/mL
		Pregnant Trimester-wise : 1st : 0.1-2.5 2nd : 0.2-3 3rd : 0.3-3	
URINE ANALYSIS			
COLOR	PALE YELLOW		
APPEARANCE	TURBID		
PH	5.0	4.7 - 7.5	
SPECIFIC GRAVITY	1.015	1.003 - 1.035	
GLUCOSE	NOT DETECTED	NOT DETECTED	
PROTEIN	DETECTED (SMALL)	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
CASTS	ABSENT		
CRYSTALS	ABSENT		
CHEMICAL EXAMINATION, URINE			
BILIRUBIN	NOT DETECTED	NOT DETECTED	
NITRITE	NOT DETECTED	NOT DETECTED	
MICROSCOPIC EXAMINATION, URINE			
WBC	30-40	0-5	/HPF
EPITHELIAL CELLS	30-50	0-5	/HPF
BACTERIA	DETECTED (+++)	NOT DETECTED	

Interpretation(s)

SERUM BLOOD UREA NITROGEN-

Causes of Increased levels Pre renal

• High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal • Renal Failure

Post Renal

• Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

· Liver disease









CLIENT CODE: CA00010147

CLIENT'S NAME AND ADDRESS:

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030

DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

ACCESSION NO: 4053VJ000652 AGE: 29 Years SEX: Female ABHA NO:

DRAWN: RECEIVED: 08/10/2022 08:33 REPORTED: 08/10/2022 13:49

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results Units <u>Final</u>

SIADH.

CREATININE, SERUM-

Higher than normal level may be due to:

- Blockage in the urinary tract
 Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
 Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis
- Muscular dystrophy

GLUCOSE, POST-PRANDIAL, PLASMA-

ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75grams of glucose in 300 ml water, over a period of 5 minutes. GLUCOSE, FASTING, PLASMA-ADA 2012 guidelines for adults as follows:

Pre-diabetics: 100 - 125 mg/dL

Diabetic: > or = 126 mg/dL

(Ref: Tietz 4th Edition & ADA 2012 Guidelines)

GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-Glycosylated hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood, the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells.

Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia, increased red cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of

testing such as glycated serum protein (fructosamine) should be considered.
"Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of

diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations.

References

- 1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006,
- 2. Forsham PH. Diabetes Mellitus: A rational plan for management. Postgrad Med 1982, 71,139-154.
 3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184. CORONARY RISK PROFILE (LIPID PROFILE), SERUM-

Serum cholesterol is a blood test that can provide valuable information for the risk of coronary artery disease This test can help determine your risk of the build up of plaques in your arteries that can lead to narrowed or blocked arteries throughout your body (atherosclerosis). High cholesterol levels usually don't cause any signs or symptoms, so a cholesterol test is an important tool. High cholesterol levels often are a significant risk factor for heart disease and important for diagnosis of hyperlipoproteinemia, atherosclerosis, hepatic and thyroid diseases.

Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn't need into triglycerides, which are stored in fat cells. High triglyceride levels are associated with several factors, including being overweight, eating too many sweets or drinking too much alcohol, smoking, being sedentary, or having diabetes with elevated blood sugar levels. Analysis has proven useful in the diagnosis and treatment of patients with diabetes mellitus, nephrosis, liver obstruction, other diseases involving lipid metabolism, and various endocrine disorders. In conjunction with high density lipoprotein and total serum cholesterol, a triglyceride determination provides valuable information for the assessment of coronary heart disease risk. It is done in fasting state.

High-density lipoprotein (HDL) cholesterol. This is sometimes called the ""good"" cholesterol because it helps carry away LDL cholesterol, thus keeping arteries open and blood flowing more freely. HDL cholesterol is inversely related to the risk for cardiovascular disease. It increases following regular exercise, moderate alcohol consumption and with oral estrogen therapy. Decreased levels are associated with obesity, stress, cigarette smoking and diabetes mellitus.

SERUM LDL The small dense LDL test can be used to determine cardiovascular risk in individuals with metabolic syndrome or established/progressing coronary artery disease, individuals with triglyceride levels between 70 and 140 mg/dL, as well as individuals with a diet high in trans-fat or carbohydrates. Elevated sdLDL levels are associated with metabolic syndrome and an 'atherogenic lipoprotein profile', and are a strong, independent predictor of cardiovascular disease.
Elevated levels of LDL arise from multiple sources. A major factor is sedentary lifestyle with a diet high in saturated fat. Insulin-resistance and pre-diabetes have also been implicated, as has genetic predisposition. Measurement of sdLDL allows the clinician to get a more comprehensive picture of lipid risk factors and tailor treatment accordingly. Reducing LDL levels will reduce the risk of CVD and MI.

Non HDL Cholesterol - Adult treatment panel ATP III suggested the addition of Non-HDL Cholesterol as an indicator of all atherogenic lipoproteins (mainly LDL and VLDL).

NICE guidelines recommend Non-HDL Cholesterol measurement before initiating lipid lowering therapy. It has also been shown to be a better marker of risk in both primary and secondary prevention studies.



Page 6 Of 9 Scan to View Report



Name	GEETHU.G.PILLAI	Age/Sex	29Yrs/Female
Ref: By:	MEDI WHEEL	Date	08.10.2022

Thanks for referral

CHEST X-RAY - PA VIEW

Trachea is central. Carina and principal bronchi are normal.

Cardio-thoracic ratio is within normal limits.

Both lungs show normal Broncho-vascular markings. No definite focal opacities noted.

No volume loss in either hemithorax.

No definite mediastinal widening or other abnormalities noted.

CP angles, diaphragm, bony cage and soft tissue shadows - not remarkable.

IMPRESSION:

Normal X-ray chest

DR. P. NIYAZI NASIR, MBBS, DMRD

(Because of technical and technological limitation complete diagnosis cannot be assured on imaging sonography. Clinical correlation, consultation if required repeat imaging required in the event of controversies. This document is not for legal purposes).

Dr. P. NIYAZI NASIR. MBBS, DMRD REG. No. 41419 CONSULTANT RADIOLOGIST DDRC SRL DIAGNOSTIC (P) LTD. KANNUR







CLIENT CODE: CA00010147

CLIENT'S NAME AND ADDRESS:

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030

DELHI INDIA 8800465156

DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

4053VJ000652 29 Years SEX: Female ACCESSION NO: AGE: ABHA NO:

DRAWN: RECEIVED: 08/10/2022 08:33 REPORTED: 08/10/2022 13:49

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results Units <u>Final</u>

Recommendations:

Results of Lipids should always be interpreted in conjunction with the patient's medical history, clinical presentation and other findings,

NON FASTING LIPID PROFILE includes Total Cholesterol, HDL Cholesterol and calculated non-HDL Cholesterol. It does not include triglycerides and may be best used in patients for whom fasting is difficult.

TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage),Burns,Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

URIC ACID, SERUM-

Causes of Increased levels Dietary

- High Protein Intake.
- Prolonged Fasting,Rapid weight loss.

Gout

Lesch nyhan syndrome.

Type 2 DM.

Metabolic syndrome.

Causes of decreased levels

- Low Zinc Intake
- OCP's
- · Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels

- Drink plenty of fluids
- · Limit animal proteins
- High Fibre foods
- Vit C Intake
- Antioxidant rich foods

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-

Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

BLOOD COUNTS-

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICESThe cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

WBC DIFFERENTIAL COUNT - NLRThe optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope. ERYTHRO SEDIMENTATION RATE, BLOOD-

Erythrocyte sedimentation rate (ESR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0 -1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

- 1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition
- 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin
 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition" SUGAR URINE POST PRANDIAL-METHOD: DIPSTICK/BENEDICT'S TEST

THYROID PANEL, SERUM-

Triiodothyronine T3, is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and







Name	Mrs. GEETHU.G.PILLAI	Age/Sex	29Yrs/Female
Ref: By:	MEDI WHEEL	Date	08.10.2022

ULTRASOUND SCAN OF ABDOMEN AND PELVIS

(With relevant image copies)

LIVER: Normal in size and shows diffusely increased echotexture. No e/o focal parenchymal lesions / IHBD. PV, HV & IVC are within normal limits.

GB: Partially distended, shows normal wall thickness. No e/o calculi/polyps/ pericholecystic collections.

CBD: Normal.

PANCREAS: Head and body visualized and are of normal size and echotexture. No e/o focal/diffuse parenchymal lesions/ductal dilatation/calculi. Tail cannot be visualized due to poor window.

SPLEEN: Normal in size and echotexture. Splenic vein shows normal diameter.

KIDNEY'S: Both kidneys are normal in size and echotexture. No e/o calculi/ hydronephrosis/ focal lesions/ perinephric collections.

RIGHT KIDNEY: Measures 108×42 mms shows a tiny calculus measuring upto 3.5 mm in the upper pole of calyx.

LEFT KIDNEY: Measures 106 x 48 mms

UB: Well distended, shows normal wall thickness. No e/o calculi/growth/diverticulae. Both UV junctions are within normal limits.

UTERUS: AV, measures 79 x 38 x 50 mms. Normal in size and echotexture.

EMT: 6 mm, normal.

OVARIES: Both ovaries are normal in size and echotexture.

RIGHT OVARY: measures 32 x 18 mms

LEFT OVARY: measures 29 x 16 mms

POD: No free fluid.

No e/o intraperitoneal free fluid/ abdominal lymphadenopathy/ mass lesion.

IMPRESSION

- > TINY RIGHT RENAL CALCULUS.
- GRADE I FATTY LIVER.

Dr. P.NIYAZI NASIR MBBS, DMRD

(Because of technical and technological limitation complete diagnosis cannot be assured on imaging sonography. Clinical correlation, consultation if required repeat imaging required in the event of controversies. This document is not for legal purposes).

Dr. P. NIYAZI NASIR. MBBS, DMRD REG. No. 41419 CONSULTANT RADIOLOGIST DDRC SRL DIAGNOSTIC (P) LTD. KANNUR







CLIENT CODE: CA00010147

CLIENT'S NAME AND ADDRESS:

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED

F701A, LADO SARAI, NEW DELHI, SOUTH DELHI, DELHI,

SOUTH DELHI 110030 **DELHI INDIA**

8800465156

DDRC SRL DIAGNOSTICS

KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

ACCESSION NO: 4053VJ000652 29 Years SEX: Female AGE: ABHA NO:

DRAWN: RECEIVED: 08/10/2022 08:33 REPORTED: 08/10/2022 13:49

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results Units <u>Final</u>

heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.
Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is

hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.

Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3

Levels in TOTAL T4 TSH3G TOTAL T3

(μIU/mL) 0.1 - 2.5 0.2 - 3.0 0.3 - 3.0 Pregnancy First Trimester (µg/dL) 6.6 - 12.4 (ng/dL) 81 - 190 6.6 - 15.5 6.6 - 15.5 100 - 260 100 - 260 2nd Trimester 3rd Trimester

Below mentioned are the guidelines for age related reference ranges for T3 and T4. T3

(ng/dL) (µg/dL) 1-3 day: 8.2 - 19.9 1 Week: 6.0 - 15.9 New Born: 75 - 260

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group.

Kindly note: Method specific reference ranges are appearing on the report under biological reference range.

- 1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.
 2. Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.
 3. Behrman R.E. Kilegman R.M., Jenson H. B. Nelson Text Book of Pediatrics, 17th Edition
 MICROSCOPIC EXAMINATION, URINE-

Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus. Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia






DDRC SRL KANNUR

GEETHU: 08_10_2022_13_03_47

20221008









CLIENT CODE: CA00010147 **CLIENT'S NAME AND ADDRESS:**

MEDIWHEEL ARCOFEMI HEALTHCARE LIMITED

F701A, LADO SARAI, NEW DELHI,

SOUTH DELHI, DELHI, SOUTH DELHI 110030 DELHI INDIA 8800465156

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

DDRC SRL DIAGNOSTICS

PATIENT NAME: GEETHU G PILLAI PATIENT ID: GEETF0810934053

ACCESSION NO: 4053VJ000652 AGE: 29 Years SEX: Female ABHA NO:

RECEIVED: 08/10/2022 08:33 08/10/2022 13:49 DRAWN: REPORTED:

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Units <u>Final</u>

MEDIWHEEL HEALTH CHECKUP BELOW 40(F)TMT

ECG WITH REPORT

REPORT

COMPLETED

USG ABDOMEN AND PELVIS

REPORT

COMPLETED

CHEST X-RAY WITH REPORT

REPORT

COMPLETED

End Of Report Please visit www.srlworld.com for related Test Information for this accession

JINSHA KRISHNAN

LAB TECHNICIAN

RESHMA RAJAN LAB TECHNICIAN

DR.INDUSARATH S CONSULTANT PATHOLOGIST

KTRAN K **Msc Medical Biochemistry**



