PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	14.5	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	45.4	%	42 - 52
RBC Count (EDTA Blood/Impedance Variation)	5.23	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	87.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	27.7	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	31.9	g/dL	32 - 36
RDW-CV (Derived from Impedance)	14.7	%	11.5 - 16.0
RDW-SD (Derived from Impedance)	44.76	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	9720	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	49.12	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	36.87	%	20 - 45
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	6.87	%	01 - 06



VERIFIED BY



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood/Impedance Variation & Flow Cytometry)	6.82	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.31	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.77	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.58	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.67	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.66	10^3 / μ1	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.03	10^3 / μ1	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	378.2	10^3 / μl	150 - 450
MPV (Blood/Derived from Impedance)	9.07	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.34	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	31	mm/hr	< 15



VERIFIED BY



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>BIOCHEMISTRY</u>			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.5	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.3	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.3	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.3	gm/dL	3.5 - 5.2
Globulin (Serum/ <i>Derived</i>)	3.0	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.4		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	27	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	34	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	110	U/L	53 - 128
GGT(Gamma Glutamyl Transpeptidase) (Serum/SZASZ standarised IFCC)	35	U/L	< 55



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	223	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	150	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

part of the day.			
HDL Cholesterol (Serum/Immunoinhibition)	39	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	154	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	30	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	184.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219



Very High: >= 220

 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u> <u>Value</u> <u>Reference Interval</u>

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio 5.7 Optimal: < 3.3 (Serum/Calculated) Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 3.8 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio 3.9 Optimal: 0.5 - 3.0

(Serum/Calculated)
Borderline: 3.1 - 6.0
High Risk: > 6.0



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	7.7	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 174.29 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.18 ng/mL 0.7 - 2.04

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Thyroxine) - Total 7.00 μg/dL 4.2 - 12.0

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.26 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Microparticle

Immunoassay(CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

Pale yellow

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

(Urine)			
Appearance	Clear		Clear
(Urine)			
Volume	20	mL	

Volume (Urine)

Protein

(Urine)

Colour

CHEMICAL EXAMINATION(Automated-

<u>Urineanalyser)</u>

5.0	4.5 - 8.0
1.025	1.002 - 1.035
Negative	Negative
0.2	0.2 - 1.0
Negative	Negative
Negative	Negative
Negative	Negative
	1.025 Negative 0.2 Negative Negative

Negative



VERIFIED BY



APPROVED BY

Negative

 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine)	Negative		Negative
Leukocytes (Urine) <u>MICROSCOPY(URINE DEPOSITS)</u>	Negative	leuco/uL	Negative
Pus Cells (Urine/Flow cytometry)	2-3	/hpf	3-5
Epithelial Cells (Urine)	1-2	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	2-3
Others (Urine)	Nil		Nil



VERIFIED BY



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'O' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



VERIFIED BY



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
BIOCHEMISTRY			
BUN / Creatinine Ratio	9		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	124	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Fasting - Urine	Negative		Negative
(Urine - F)			
Glucose Postprandial (PPBS)	113	mg/dL	70 - 140
(Plasma - PP/GOD - POD)			

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Glucose Postprandial - Urine (Urine - PP)	Sample Not Given		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	8	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.9	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 6.7 mg/dL 3.5 - 7.2 (Serum/Uricase/Peroxidase)



 PID No.
 : MED120813880
 Register On
 : 17/02/2022 9:17 AM

 SID No.
 : 522209069
 Collection On
 : 17/02/2022 12:40 PM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/02/2022 10:27 AM

 Type
 : OP
 Printed On
 : 08/03/2022 12:07 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Prostate specific antigen - Total(PSA) (Serum/Chemiluminescent Microparticle Immunoassay(CMIA))	1.22	ng/mL	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0



-- End of Report --

Name	MR. KUMARESHAN	ID	MED120813880
Age & Gender	48Y/MALE	Visit Date	17 Feb 2022
Ref Doctor Name	MediWheel	-	

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (14.8cm) and shows increased echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is partially contracted. Grossly appears normal.

CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size (9.5cm) and echopattern.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cm)	Parenchymal thickness (cm)
Right Kidney	10.6	1.6
Left Kidney	10.6	2.2

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

Prevoid: 200 cc Postvoid: 2cc.

PROSTATE shows normal shape, size and echopattern. It measures 3.3 x 4.2 x 2.9 cm volume: 22cc.

No evidence of ascites.

IMPRESSION:

• Grade II fatty infiltration of liver.

DR. H.K. ANAND DR. C.R RAMACHANDRA DR. LOHITH H.P DR. VARSHA KALE CONSULTANT RADIOLOGISTS

Vk/sm

Name	MR. KUMARESHAN	ID	MED120813880
Age & Gender	48Y/MALE	Visit Date	17 Feb 2022
Ref Doctor Name	MediWheel	-	

Name	MR. KUMARESHAN	ID	MED120813880
Age & Gender	48Y/MALE	Visit Date	17 Feb 2022
Ref Doctor Name	MediWheel		

2D ECHOCARDIOGRAPHIC STUDY

M-mode measurement:

AORTA 2.83 cms. LEFT ATRIUM 2.40 cms. **AVS** 1.45 cms. LEFT VENTRICLE (DIASTOLE) 4.53 cms. (SYSTOLE) 2.83 cms. **VENTRICULAR SEPTUM** (DIASTOLE) 0.93 cms. (SYSTOLE) 1.49 cms. **POSTERIOR WALL** (DIASTOLE) 1.08 cms. (SYSTOLE) 1.34 cms. **EDV** 93 ml. **ESV** ml. 30 % FRACTIONAL SHORTENING 30 **EJECTION FRACTION** % 60 **EPSS** cms. **RVID** 1.85 cms.

DOPPLER MEASUREMENTS:

MITRAL VALVE: E - 0.7 m/s A - 0.8 m/s TRIVIAL MR.

AORTIC VALVE: 1.1 m/s NO AR.

TRICUSPID VALVE: E - 0.3 m/s A - 0.4 m/s TRIVIAL TR.

PULMONARY VALVE: 0.6 m/s NO PR.

Name	MR. KUMARESHAN	ID	MED120813880
Age & Gender	48Y/MALE	Visit Date	17 Feb 2022
Ref Doctor Name	MediWheel		

2D ECHOCARDIOGRAPHY FINDINGS:

Left Ventricle : Normal size, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal.Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- NORMAL SIZED CARDIAC CHAMBERS.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- TRIVIAL MR.
- TRIVIAL TR.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. ANAND KUMAR M, MD DM CONSULTANT INTERVENTIONAL CARDIOLOGIST

Name	MR. KUMARESHAN	ID	MED120813880
Age & Gender	48Y/MALE	Visit Date	17 Feb 2022
Ref Doctor Name	MediWheel	-	



Name	KUMARESHAN	ID	MED120813880
Age & Gender	48Y/M	Visit Date	Feb 17 2022 9:14AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

• No significant abnormality detected.

DR. H.K. ANAND DR. VARSHA KALE DR. LOHITH H.P

CONSULTANT RADIOLOGISTS

DR. C.R. RAMACHANDRA