

# बैंक ऑफ़ बड़ौदा Bank of Baroda

जाम

प्रदीप कुमार सैनी

Name

PRADEEP KUMAR SAINI

कर्मवारी कूट क्र.

E.C. No. 78939





जारीकर्ता प्राधिकारी Issuing Authority




.धारक के ढरताक्षर Signature of Holder

Dr. U/C. GUPTA MBBS, MD (Physician) RMC No. 281



 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

9 +91 141 4824885 maxcarediagnostics1@gmail.com





# **General Physical Examination**

Date of Examination: <u>Q4/1@/</u> QQ
Name: PRADEEP KUMAR SAINI Age: 36/YRDOB: 11/03/1986Sex: Mole
Referred By: BANK OF BARODA
Photo ID: TD CARD ID#: 78939
Ht: 168 (cm) Wt: 60 (Kg)
Chest (Expiration): & (cm)
Blood Pressure: 25/ mm Hg PR: 78/min RR: 17/min Temp: Afelon le
BMI &&
Eye Examination: RIEY G/G, N/G, NCB LIEJ G/G, N/G, NCB
Other: N/A
On examination he/she appears physically and mentally fit: Yes/No  Signature Of Examine: Name of Examinee: PRADEEP ROMAR SAIN
Signature Medical Examiner:  Dr. U. C. GUPTA  MBBS, MD (Physician)  RMC No. 291



© +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. PRADEEP KUMAR SAINI

Age:- 36 Yrs 9 Mon 15 Days

Sex :- Male



Patient ID :-12222724

Date :- 24/12/2022

09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company :-

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:34

# **HAEMATOLOGY**

Test Name	Value	Unit	<b>Biological Ref Interval</b>
FULL BODY HEALTH CHECKUP BELOW 40	MALE		
HAEMOGARAM	W/ YCL		
HAEMOGLOBIN (Hb)	14.3	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	2.90 L	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	51.0	%	40.0 - 80.0
LYMPHOCYTE	40.0	%	20.0 - 40.0
EOSINOPHIL	3.0	%	1.0 - 6.0
MONOCYTE	6.0	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	4.93	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	44.90	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	91.0	£ £	83.0 - 101.0
MEAN CORP HB (MCH)	29.0	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	31.8	g/dL	31.5 - 34.5
PLATELET COUNT	209	x10^3/uL	150 - 410
RDW-CV	14.4 H	%	11.6 - 14.0
MENTZER INDEX A complete blood picture (CBP) is a kind of blood test	18.46 H	es a person's overall health a	0.00 - 0.00

A complete blood picture (CBP) is a kind of blood test that is done to assess a person's overall health and diagnose a wide range of health disorders like leukemia, anemia and other infections.

A complete blood count (CBC) is a complete blood test that diagnose many components and features of a persons blood which includes: -

(CBC): Methodology: TLC,TRBC,PCV,PLT Impedance method, HB Calorimetric method, and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: MINDRAY BC-3000 Plus 3 part automatic analyzer,

VIKARANTJI

Technologist

Page No: 1 of 15

DR.TANU RUNGTA

<sup>\*</sup>Red Blood Cells (RBC), which carry oxygen -

<sup>\*</sup>White Blood Cells (WBC), which help in fighting against infections -

<sup>\*</sup>Hemoglobin, which is the oxygen carrying protein in the red blood cells -

<sup>\*</sup>Hematocrit (HCT), the proportion of RBC to the fluid component, or plasma present in blood -

<sup>\*</sup>Platelets, which aid in blood clotting



© +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. PRADEEP KUMAR SAINI

Age:- 36 Yrs 9 Mon 15 Days

Sex :- Male



Patient ID :-12222724 Date :- 24/12/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company :-

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:34

09:40:38

### HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

08

mm in 1st hr

00 - 15

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases.ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein.ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



VIKARANTJI

Technologist Page No: 2 of 15 DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



© +91 141 4824885 ⊚ maxcarediagnostics1@gmail.com



Date :- 24/12/2022 09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Patient ID :-12222724

Company :- Mr.MEDIWHEEL

NAME :- Mr. PRADEEP KUMAR SAINI Age :- 36 Yrs 9 Mon 15 Days

Sex :- Male

Compan

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



VIKARANTJI

Page No: 3 of 15



⊕ +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. PRADEEP KUMAR SAINI

Age :-36 Yrs 9 Mon 15 Days

Sex :-Male



Date :- 24/12/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company:-

Patient ID: -12222724

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:34

09:40:38

**BIOCHEMISTRY** 

**Test Name** Value Unit **Biological Ref Interval** 

FASTING BLOOD SUGAR (Plasma)

Methord:- GOD POD

91.7

mg/dl

70.0 - 115.0

Impaired glucose tolerance (IGT) 111 - 125 mg/dL Diabetes Mellitus (DM) > 126 mg/dL

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm,

hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result

from excessive insulin

therapy or various liver diseases.



VIKARANTJI

**Technologist** Page No: 4 of 15

form DR.TANU RUNGTA MD (Pathology) RMC No. 17226




# **P3 HEALTH SOLUTIONS LLP**

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 maxcarediagnostics1@gmail.com



Age :-

36 Yrs 9 Mon 15 Days

Sex :- Male



Patient ID :-12222724

Date :- 24/12/2022

09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:54:05

# **BIOCHEMISTRY**

Test Name	Value	Unit	Biological Ref Interval

### FULL BODY HEALTH CHECKUP BELOW 40 MALE

BLOOD SUGAR PP (Plasma) Methord:- GOD PAP

96.4

mg/dl

70.0 - 140.0

**Instrument Name: HORIBA Interpretation:** Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.



VIKARANTJI

Technologist

Page No: 1 of 1

DR.TANU RUNGTA



NAME :- Mr. PRADEEP KUMAR SAINI

Age :-36 Yrs 9 Mon 15 Days

Male Sex :-



Patient ID :-12222724

Date :- 24/12/2022

09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Company:-

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:34

# HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (HbA1C) Methord:- CAPILLARY with EDTA	5.0	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Methord:- Calculated Parameter	103	mg/dL	68 - 125

#### INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA) Reference Group HbA1c in % Non diabetic adults >=18 years < 5.7 At risk (Prediabetes) 5.7 - 6.4 Diagnosing Diabetes >= 6.5

#### CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings. Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al ]

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis.
   Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease
- 2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.
- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH
   Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH
- 4. Erythrocyte destruction
- Increased HbA1c: increased erythrocyte life span: Splenectomy.
   Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use chronic renal failure
- Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

1. Shortened RBC life span -HbA1c test will not be accurate when a person has a condition that affects the average lifespan of red blood cells (RBCs), such as hemolytic anemia or blood loss. When the lifespan of RBCs in circulation is shortened, the A1c result is faisely low and is an unreliable measurement of a person's average glucose over time 2. Abnormal forms of hemoglobin – The presence of some hemoglobin variants, such as hemoglobin S in sickle cell anemia, may affect certain methods for measuring A1c. In these cases, fructosamine can be used to monitor glucose control.

### Advised:

1.To follow patient for glycemic control test like fructosamine or glycated albumin may be performed instead.

2. Hemoglobin HPLC screen to analyze abnormal hemoglobin variant.
estimated Average Glucose (eAG): based on value calculated according to National Glycohemoglobin Standardization Program (NGSP) criteria.

VIKARANTJI

**Technologist** 

Page No: 5 of 15

Janu DR.TANU RUNGTA



© +91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. PRADEEP KUMAR SAINI

Age:- 36 Yrs 9 Mon 15 Days

Sex :- Male



Patient ID :-12222724

Date :- 24/12/2022

09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:34

# HAEMATOLOGY

BLOOD GROUP ABO Methord:- Haemagglutination reaction "A" POSITIVE



VIKARANTJI

Technologist

Page No: 6 of 15

DR.TANU RUNGTA



+91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. PRADEEP KUMAR SAINI

Age :-36 Yrs 9 Mon 15 Days

Sex :-Male



Patient ID: -12222724

Date :- 24/12/2022

09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:34

# **BIOCHEMISTRY**

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE TOTAL CHOLESTEROL Methord:- CHOD-PAP methodology	135.00	mg/dl	Desirable <200 Borderline 200-239 High> 240
InstrumentName:MISPA PLUS Interpretation: disorders.	: Cholesterol measurements	s are used in the diagnosis a	and treatments of lipid lipoprotein metabolism
TRIGLYCERIDES Methord:- GPO-TOPS methodology	106.00	mg/dl	Normal <150 Borderline high 150-199 High 200-499
A STATE OF THE STA			Very high >500

InstrumentName: MISPA PLUS Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL Methord:- Selective inhibition Method

mg/dl

Male 35-80

Female 42-88

Instrument Name: MISPA PLUS Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to

precipitation methods. LDL CHOLESTEROL Optimal <100 47.23 mg/dl Methord:- Calculated Method Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190 VLDL CHOLESTEROL 21.20 mg/dl 0.00 - 80.00Methord:- Calculated T.CHOLESTEROL/HDL CHOLESTEROL RATIO 1.93 0.00 - 4.90Methord:- Calculated LDL / HDL CHOLESTEROL RATIO 0.67 0.00 - 3.50

TOTAL LIPID

Methord:- Calculated

429.89

mg/dl

400.00 - 1000.00

1. Measurements in the same patient can show physiological analytical variations. Three scrialsamples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.

2. As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended

3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated fromperipheral tissues.

Comments: 1- ATP III suggested the addition of Non HDL Cholesterol (Total Cholesterol - HDL Cholesterol) as an indicator of all VIKARANTJI Janu

**Technologist** 

Page No: 7 of 15

DR.TANU RUNGTA



# P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

© +91 141 4824885 maxcarediagnostics1@gmail.com



Date :- 24/12/2022

09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Patient ID: -12222724

Company:- Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:35

# NAME :- Mr. PRADEEP KUMAR SAINI

Age :- 36

36 Yrs 9 Mon 15 Days

Sex :- Male

# **BIOCHEMISTRY**

LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Methord:- DMSO/Diazo	0.59	mg/dL	Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Methord:- DMSO/Diazo	0.24	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Methord:- Calculated	0.35	mg/dl	0.30-0.70
SGOT Methord:- IFCC	29.5	U/L	Men- Up to - 37.0 Female - Up to - 31.0
SGPT Methord:- IFCC	40.0	U/L	Men- Up to - 40.0 Female- Up to - 31.0
SERUM ALKALINE PHOSPHATASE Methord:- DGKC - SCE	94.80	U/L	53.00 - 141.00
SERUM GAMMA GT Methord:- Szasz methodology Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than thos	19.00	U/L s in cases of obstructive jaundice and	10.00 - 45.00
metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post- hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times r	normal)are observed with i	nfectious hepatitis.	z
SERUM TOTAL PROTEIN Methord:- Direct Biuret Reagent	6.69	g/dl	5.10 - 8.00
SERUM ALBUMIN Methord:- Bromocresol Green	4.48	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	2.21	gm/dl	2.20 - 3.50
A/G RATIO	2.03		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note:- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B,C, paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.

VIKARANTJI

Technologist

Page No: 9 of 15

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226

form



♦ +91 141 4824885 
maxcarediagnostics1@gmail.com



Date :- 24/12/2022 09:40:38

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:35

NAME :- Mr. PRADEEP KUMAR SAINI Age :-36 Yrs 9 Mon 15 Days

Sex :-

Company :-

Patient ID: -12222724

BIOCHEMISTRY

## RFT / KFT WITH ELECTROLYTES

SERUM UREA Methord:- Urease/GLDH 18.30

mg/dl

10.00 - 50.00

InstrumentName: HORIBA CA 60 Interpretation: Urea measurements are used in the diagnosis and treatment of certain renal and metabolic

SERUM CREATININE Methord:- Jaffe's Method

0.85

mg/dl

Males: 0.6-1.50 mg/dl

Females : 0.6 -1.40 mg/dl

Interpretation:

Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not

clinically significant. SERUM URIC ACID

4.02

mg/dl

2.40 - 7.00

InstrumentName: HORIBA YUMIZEN CA60 Daytona plus Interpretation: Elevated Urate: High purine diet, Alcohol. Renal insufficiency, Drugs, Polycythaemia vera, Malignancies, Hypothyroidism, Rare enzyme defects, Downs syndrome, Metabolic syndrome, Pregnancy, Gout.

**SODIUM** Methord: - ISE 135.2

mmol/L

Interpretation: Decreased sodium - Hyponatraemia Causes include: fluid or electrolyte loss, Drugs, Oedematous states, Legionnaire's disease and other chest infections, pseudonatremia, Hyperlipidaemias and paraproteinaemias, endocrine diseases, SIADH.

**POTASSIUM** 

Methord:- ISE

4.63

mmol/L

3.50 - 5.50

A. Elevated potassium (hyperkalaemia). Artefactual, Physiological vation, Drugs, Pathological states, Renal failure Interpretation: Adrenocortical insufficiency, metabolic acidoses, very high platelet or white cell counts B. Decreased potassium (hypokalaemia)Drugs, Liquoric, Diarrhoea and vomiting, Metabolic alkalosis, Corticosteroid excess, Oedematous state, Anorexia nervosa/bulimia

**CHLORIDE** 

105.5

mmol/L

94.0 - 110.0

Interpretation: Used for Electrolyte monitoring.

SERUM CALCIUM

Methord:- Arsenazo III Method

8.64 L

mg/dL

8.80 - 10.20

InstrumentName: MISPA PLUS Interpretation: Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia. Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN VNEARIA NIFCI Biuret Reagent

6.69

g/dl

5.10 - 8.00

form

**Technologist** 

Page No: 10 of 15

DR.TANU RUNGTA



⊕ +91 141 4824885 maxcarediagnostics1@gmail.com

Patient ID :-12222724

Date :- 24/12/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Mr.MEDIWHEEL Company :-

Final Authentication: 24/12/2022 17:39:35

# NAME: - Mr. PRADEEP KUMAR SAINI

36 Yrs 9 Mon 15 Days Age :-

Sex :-

# **BIOCHEMISTRY**

SERUM ALBUMIN Methord:- Bromocresol Green	4.48	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	2.21	gm/dl	2.20 - 3.50
A/G RATIO	2.03		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

### **INTERPRETATION**

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR .in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.

VIKARANTJI

**Technologist** Page No: 11 of 15

DR.TANU RUNGTA MD (Pathology) RMC No. 17226



# **P3 HEALTH SOLUTIONS LLP**

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

S +91 141 4824885 S maxcarediagnostics1@gmail.com

NAME :- Mr. PRADEEP KUMAR SAINI Pati

Age:- 36 Yrs 9 Mon 15 Days

Sex :- Male

Patient ID :-12222724 Date :- 24/12/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company:- Mr.MEDIWHEEL

Final Authentication: 24/12/2022 17:39:35

09:40:38

# **CLINICAL PATHOLOGY**

Test Name	Value Unit	Biological Ref Interval
Urine Routine		
PHYSICAL EXAMINATION		
COLOUR	PALE YELLOW	PALE YELLOW
APPEARANCE	Clear	Clear
<b>CHEMICAL EXAMINATION</b>		
REACTION(PH)	5.0	5.0 - 7.5
SPECIFIC GRAVITY	1.030	1.010 - 1.030
PROTEIN	NIL	NIL
SUGAR	NIL	NIL
BILIRUBIN	NEGATIVE	NEGATIVE
UROBILINOGEN	NORMAL	NORMAL
KETONES	NEGATIVE	NEGATIVE
NITRITE	NEGATIVE	NEGATIVE
MICROSCOPY EXAMINATION		
RBC/HPF	NIL /HPF	NIL
WBC/HPF	2-3 /HPF	2-3
EPITHELIAL CELLS	2-3 /HPF	2-3
CRYSTALS/HPF	ABSENT	ABSENT
CAST/HPF	ABSENT	ABSENT
AMORPHOUS SEDIMENT	ABSENT	ABSENT
BACTERIAL FLORA	ABSENT	ABSENT
YEAST CELL	ABSENT	ABSENT
OTHER	ABSENT	

VIKARANTJI

Technologist

Page No: 12 of 15

Jane

DR.TANU RUNGTA MD (Pathology) RMC No. 17226



🕓 +91 141 4824885 🖨 maxcarediagnostics1@gmail.com



Date :- 24/12/2022

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp:-

Mr.MEDIWHEEL Company:-

Final Authentication: 24/12/2022 17:39:35

NAME :- Mr. PRADEEP KUMAR SAINI

36 Yrs 9 Mon 15 Days Age :-Sex :-

Male

### TOTAL THYROID PROFILE

### **IMMUNOASSAY**

Test Name	Value	Unit	Biological Ref Interval
THYROID-TRIIODOTHYRONINE T3	<b>4.74</b> H	ng/mL	0.70 - 2.04

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions simoultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by †serum T3 & T4 values along with "TSH level 2.Low TSH,high FT4 and TSH receptor antibody(TRAb) +ve seen in patients with Graves disease 3.Low TSH,high FT4 and TSH receptor antibody(TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.HighTSH,Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis 5.HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency 6.Low TSH,Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism 7.Primary hypothyroidism is accompanied by † serum T3 and T4 values & 'serum TSH levels Normal T4 levels accompanied by \*T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis9 Normal or T3 & T4 along with "TSH indicate mild / Subclinical Hypothyroidism .11.Normal T3 & "T4 along with "TSH is seen in Hypothyroidism .12.Normal T3 & T4 levels with 'TSH indicate Mild / Subclinical Hypothyroidism .13 & "T4 along with "TSH indicate Mild / Subclinical Hypothyroidism .14.Normal T3 & "T4 along with "TSH indicate Mild / Subclinical Hypothyroidism .15 with T3 wit

DURING PREGNANCY - REFERENCE RANGE for TSH IN ulu/mL (As per American Thyroid Association) 1st Trimester: 0.10-2.50 ullu/mL 2nd Trimester: 0.20-3.00 ullu/mL 3rd Trimester: 0.30-3.00

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher by the condition of the condition Methord:- ECLIA

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration.Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simoultaneous measuremen of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1. Primary hyperthyroidism is accompanied by tserum T3 & T4 values along with \*TSH level.2.Low TSH,high FT4 and TSH receptor antibody(TRAb) +ve seen in patients with Graves disease 3.Low TSH,high FT4 and TSH receptor antibody(TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4. HighTSH,Low FT4 and Thyroid microsomal The seep in patients with Taxic adenomal Taxic Multinodular gotter 4. High 1 SH, Low F14 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis 5. High TSH, Low F14 and Thyroid microsomal antibody normal seen in patients with Iddine deficiency/Congenital T4 synthesis deficiency 6. Low TSH, Low F14 and TRH stimulation test - Delayed response seen in patients with Tertiary hypothyroidism 7. Primary hypothyroidism is accompanied by 1 serum T3 and T4 values & 'serum TSH levels8. Normal T4 levels accompanied by 'T3 levels and low TSH are seen in patients with 'TSH indicate mild / Subclinical Hypothyroidism .12. Normal T3 & T4 along with 'TSH indicate mild / Subclinical Hypothyroidism .11. Normal T3 & T4 along with 'TSH indicate mild / Subclinical Hypothyroidism .12. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .12. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .12. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .12. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .12. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild / Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild 'Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mild 'Subclinical Hypothyroidism .13. Normal T3 & T4 levels with 'TSH indicate mil

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester: 0.10-2.50 uIU/mL 2nd Trimester: 0.20-3.00 uIU/mL 3rd Trimester: 0.30-3.00 ulU/mL. The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

TSH Methord:- ECLIA 0.020 L

μIU/mL

0.350 - 5.500

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simoultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay VIKARANAThybidism is accompanied by †serum T3 & T4 values along with [ TSH level

Technologist

Page No: 14 of 15

Janu DR.TANU RUNGTA



O B-14, Vidhyadhar Enclave - II, Near Axis Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023 \$\infty\$ +91 141 4824885 \$\infty\$ maxcarediagnostics1@gmail.com



NAME:	MR. PRADEEP KUMAR SAINI	AGE	36 YRS/M
REF.BY	BANK OF BARODA	DATE	24/12/2022

# **CHEST X RAY (PA VIEW)**

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected.

Shallni


DR.SHALINI GOEL M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954











MR. PRADEEP KUMAR SAINI	36 Y/Male
Registration Date: 24/12/2022	Ref. by: BANK OF BARODA

# **ULTRASOUND OF WHOLE ABDOMEN**

**Liver** is of normal size (11.2 cm). Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

**Gall bladder** is well distended. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

**Pancreas** is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape (10.4 cm). Echotexture is normal. No focal lesion is seen.

**Kidneys** are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Right kidney is measuring approx. 10.3 x 4.6 cm.

**Left kidney** is measuring approx. 10.8 x 5.6 cm.

Urinary bladder does not show any calculus or mass lesion.

Prostate is normal in size with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified. No significant free fluid is seen in pelvis.

# **IMPRESSION:**



**DR.SHALINI GOEL** 

M.B.B.S, D.N.B (Radiodiagnosis)

RMC no.: 21954

Ref.: BANK OF BARODA Test Date: 24-Dec-2022(10:31:51) Notch: 50Hz 0.05Hz - 100Hz 12229451322732/Mr Pradeep Kumar Saini 36Yrs/Male B-14, Vidhyanagar Nagar, Enclave, Phase-2, Jaipur P3 HEALTH SOLUTIONS LLP P-QRS-T axis: 0 - 177 - 109 • (Deg) FINDINGS: Normal Variant with Non Specific ST Changes Comments: Vent Rate: 107 bpmPR Interval: ms; avR QRS Duration: 200 ms; QT/QTc Int: 423/565 ms Kgs/ Cms  $\leq$ 10mm/mV 25mm/Sec mmHg 5 6 Bonderline winus RMC No.: 35703
MBBS, DIP. CARDIO (ESCORTS)
D.E.M. (RCGP-UK) Dr. Naresh Kumar Mohanka QRS Duration: 200 ms QT/QTc: 423/565ms P-QRS-T Axis: 0 - -177 - -109 (Deg) PR Interval: 182 ms AB 3S tachyc andia 0

3-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

322301/MR PRADEEP KUMAR SAINI 36 Yrs/Male 0 Kg/0 Cms ate: 24-Dec-2022 10:34:26 AM ef.By : BANK OF BARODA


Protocol : BRUCE History :

oummary

itage 2 : sguibni: stage 1 supine itage Recovery *lecovery* :xStart itanding Recovery <sup>3</sup>eakEx ibjective: dvice/Comments: *\ecover* edication: Max WorkLoad attained :7.7(Fair Effort Tolerance) Max BP : 155/90(mmHg) Max HR Attained Exercise Time StageTime PhaseTime Speed
(Min:Sec) (Min:Sec) (mph) 3:00 2:00 0:36 1:00 3:01 3:01 6:37 6:02 3:02 :190 bpm 110% of Max Predictable HR 173 0.0 0.0 0.0 Grade 14.0 12.0 0.0 0.0 0.0 0.0 10 1.0 1.0 1.0 1.0 METS 1.0 17 RMC No.: 35703 //BBS, DIP. CARDIO (ESCORTS) D.E.M. (RCGP-UK) 5 0 aresh Kumar Mohanka 143 174 118 109 H.R. 137 190 176 129 142 129 109 135/85 145/90 145/90 135/85 125/85 155/90 145/90 145/90 125/85 125/85 125/85 B.P. R.P.P. 221 252 255 147 136 136 275 191 161 198 PVC -1 1 . Base Wine Changes week STE MAKEDO legal Comments 7690 000 ٧6 PreEx ٧6 PeakEx 0.1 0.2 6 بالاله ع أما ا how & thyendian avf avL avR ٧6 12 **5** 4 5 < III MANIMANTA III **TTS** J. W. W. W. D. 0.5 mm/Div 12 15 18 21 Min. heavesare

6410J





Average

