Age / Sex : 49 Year(s) / Female

 PID No.
 : MED111023576
 Register On
 : 17/03/2022 8:36 AM

 SID No.
 : 422021489
 Collection On
 : 17/03/2022 9:05 AM

**Report On** : 19/03/2022 1:10 PM

Ref. Dr : MediWheel

Type



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<b>HAEMATOLOGY</b>			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	11.1	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	35.0	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.10	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	85.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	27.1	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	31.8	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	14.8	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	44.03	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	7700	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	51.1	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	37.1	%	20 - 45





PID No. : MED111023576 : 422021489 SID No.

Age / Sex : 49 Year(s) / Female Report On : 19/03/2022 1:10 PM

Type : OP

Ref. Dr : MediWheel Register On : 17/03/2022 8:36 AM

Collection On : 17/03/2022 9:05 AM

**Printed On** : 26/03/2022 12:31 PM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	4.1	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	6.7	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	1.0	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.93	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.86	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.32	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.52	10^3 / μ1	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.08	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	318	10^3 / μl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	9.1	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.29	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate)	25	mm/hr	< 20



(Citrated Blood/Modified Westergren)



 PID No.
 : MED111023576
 Register On
 : 17/03/2022 8:36 AM

 SID No.
 : 422021489
 Collection On
 : 17/03/2022 9:05 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 19/03/2022 1:10 PM

**Printed On** 

MEDALI

Type : OP

Ref. Dr : MediWheel

Investigation	<u>Observed</u> Value	<u>Unit</u>	<u>Biological</u> Reference Interval

: 26/03/2022 12:31 PM

	<u>Value</u>		Reference Interval
<b>BIOCHEMISTRY</b>			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.4	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.2	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.8	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.4	gm/dL	3.5 - 5.2
Globulin (Serum/Derived)	3.4	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.3		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	22	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	22	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	90	U/L	42 - 98
GGT(Gamma Glutamyl Transpeptidase)	34	U/L	< 38



(Serum/SZASZ standarised IFCC)



 PID No.
 : MED111023576
 Register On : 17/03/2022 8:36 AM

 SID No.
 : 422021489
 Collection On : 17/03/2022 9:05 AM

**Printed On** 

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	257	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	244	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 26/03/2022 12:31 PM

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.

r			
HDL Cholesterol (Serum/Immunoinhibition)	39	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	169.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	48.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	218.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220





Age / Sex : 49 Year(s) / Female

 PID No.
 : MED111023576
 Register On : 17/03/2022 8:36 AM

 SID No.
 : 422021489
 Collection On : 17/03/2022 9:05 AM

Report On

**Printed On** 



Type : OP

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

**INTERPRETATION:** 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 19/03/2022 1:10 PM

: 26/03/2022 12:31 PM

Total Cholesterol/HDL Cholesterol Ratio 6.6 Optimal: < 3.3

Low Risk: 3.4 - 4.4

Average Risk: 4.5 - 7.1

Moderate Risk: 7.2 - 11.0

High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 6.3 Optimal: < 2.5 (TG/HDL) Mild to moderate risk: 2.5 - 5.0 (Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio 4.3 Optimal: 0.5 - 3.0 (Serum/Calculated) Borderline: 3.1 - 6.0 High Risk: > 6.0





 PID No.
 : MED111023576
 Register On
 : 17/03/2022 8:36 AM

 SID No.
 : 422021489
 Collection On
 : 17/03/2022 9:05 AM

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	6.2	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 26/03/2022 12:31 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

**Printed On** 

Estimated Average Glucose 131.24 mg/dL

(Whole Blood)

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.





PID No. : MED111023576 Register On : 17/03/2022 8:36 AM

**Printed On** 

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Unit</u>	<u>Biological</u>
	Value	Reference Interval

: 26/03/2022 12:31 PM

#### **IMMUNOASSAY**

#### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.24 ng/mL 0.7 - 2.04

(Serum/CMIA)

#### INTERPRETATION:

#### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Thyroxine) - Total 7.06 μg/dL 4.2 - 12.0

(Serum/CMIA)

#### INTERPRETATION:

#### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 4.61 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Microparticle

Immunoassay(CMIA))

#### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

#### Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&amplt,0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



PID No. : MED111023576 **Register On** : 17/03/2022 8:36 AM : 422021489 SID No. Collection On : 17/03/2022 9:05 AM Age / Sex : 49 Year(s) / Female Report On : 19/03/2022 1:10 PM



Type : OP

**Printed On** : 26/03/2022 12:31 PM Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

## **CLINICAL PATHOLOGY**

#### PHYSICAL EXAMINATION

Colour (Urine)	Pale Yellow		
Appearance (Urine)	Clear		Clear
Volume	15	mL	
(Urine)			
CHEMICAL EXAMINATION(Automated-			

#### <u>Urineanalyser)</u>

рН	6.5	4.5 - 8.0
(Urine/AUTOMATED URINANALYSER)		
Specific Gravity (Urine)	1.005	1.002 - 1.035
Ketones (Urine)	Negative	Negative
Urobilinogen (Urine/AUTOMATED URINANALYSER)	0.2	0.2 - 1.0
Blood (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Nitrite (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Bilirubin (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Protein (Urine)	Negative	Negative



**VERIFIED BY** 



 PID No.
 : MED111023576
 Register On Register On Register On SID No.
 : 17/03/2022 8:36 AM SID No.

 SID No.
 : 422021489
 Collection On Report On SID No.
 : 17/03/2022 9:05 AM SID No.

**Printed On** 



Ref. Dr : MediWheel

: OP

Type

<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine)	Negative		Negative
Leukocytes (Urine)	Negative	leuco/uL	Negative
MICROSCOPY(URINE DEPOSITS)			
Pus Cells (Urine/Flow cytometry)	4-6	/hpf	3-5
Epithelial Cells (Urine)	4-6	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	2-3
Others (Urine)	Nil		Nil
Casts (Urine/Flow cytometry)	Nil	/hpf	0 - 1
Crystals (Urine)	Nil		NIL

: 26/03/2022 12:31 PM



**VERIFIED BY** 



PID No. : MED111023576 **Register On** : 17/03/2022 8:36 AM : 422021489 SID No. Collection On : 17/03/2022 9:05 AM

Age / Sex : 49 Year(s) / Female Report On : 19/03/2022 1:10 PM

**Printed On** 

Type : OP

Ref. Dr : MediWheel

: 26/03/2022 12:31 PM

Investigation <u>Observed</u> <u>Unit</u> **Biological** <u>Value</u> Reference Interval

#### **IMMUNOHAEMATOLOGY**

BLOOD GROUPING AND Rh TYPING 'A' 'Positive'

(EDTA Blood/Agglutination)



**VERIFIED BY** 



**Printed On** 

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	12.9		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	90	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 26/03/2022 12:31 PM

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Fasting - Urine (Urine - F)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD - POD)	89	mg/dL	70 - 140

#### INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Glucose Postprandial - Urine (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	9	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.7	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 3.9 mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)





**APPROVED BY** 

-- End of Report --

Name	BHAVANI B S	ID	MED111023576
Age & Gender	49/FeMale	Visit Date	17-03-2022 00:00:00
Ref Doctor Name	MediWheel		



#### 2 D ECHOCARDIOGRAPHIC STUDY

#### M mode measurement:

AORTA : 1.9cms

LEFT ATRIUM : 2.4cms

AVS :----

LEFT VENTRICLE (DIASTOLE) : 4.3cms

(SYSTOLE) : 2.9cms

VENTRICULAR SEPTUM (DIASTOLE) : 0.8cms

(SYSTOLE) : 1.1cms

POSTERIOR WALL (DIASTOLE) : 0.8cms

(SYSTOLE) : 1.2cms

EDV : 84ml

ESV : 33ml

FRACTIONAL SHORTENING : 31%

EJECTION FRACTION : 61%

EPSS :---

RVID : 1.5cms

#### **DOPPLER MEASUREMENTS:**

MITRAL VALVE : E' 0.88 m/s A' 0.71 m/s NO MR

AORTIC VALVE : 0.98 m/s NO AR

TRICUSPID VALVE : E' 2.13 m/s A' - m/s NO TR

PULMONARY VALVE : 0.66 m/s NO PR

Name	BHAVANI B S	ID	MED111023576
Age & Gender	49/FeMale	Visit Date	17-03-2022 00:00:00
Ref Doctor Name	MediWheel		



#### **2D ECHOCARDIOGRAPHY FINDINGS:**

Left ventricle : Normal size, Normal systolic function.

No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral valve : Normal, No mitral valve prolapsed.

Aortic valve : Normal, Trileaflet.

Tricuspid valve : Normal.

Pulmonary valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

#### **IMPRESSION:**

- > NORMAL SIZED CARDIAC CHAMBERS.
- > NORMAL LV SYSTOLIC FUNCTION. EF:61 %.
- > NO REGIONAL WALL MOTION ABNORMALITIES.
- > NORMAL VALVES.
- > NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. K.S. SUBRAMANI. MBBS, MD, DM (CARDIOLOGY) FESC SENIOR CONSULTANT INTERVENTIONAL CARDIOLOGIST Kss/da

Name	BHAVANI B S	ID	MED111023576
Age & Gender	49/FeMale	Visit Date	17-03-2022 00:00:00
Ref Doctor Name	MediWheel		



- Note:

  \* Report to be interpreted by qualified medical professional.

  \* To be correlated with other clinical findings.

  \* Parameters may be subjected to inter and intra observer variations.

Name	BHAVANI B S	ID	MED111023576
Age & Gender	49/FeMale		17-03-2022 00:00:00
Ref Doctor Name	MediWheel		



#### ABDOMINO-PELVIC ULTRASONOGRAPHY (TAS + TVS)

**LIVER** shows normal in shape, size and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

**GALL BLADDER** shows normal shape and has clear contents. Gall bladder wall is of normal thickness. CBD is of normal calibre.

**PANCREAS** has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

**SPLEEN** shows normal shape, size and echopattern. Spleen measures 6.9 cms in long axis and 3.2 cms in short axis.

No demonstrable Para -aortic lymphadenopathy.

**KIDNEYS** move well with respiration and have normal shape, size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

·	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	9.5	1.1
Left Kidney	11.4	1.2

**URINARY BLADDER** shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

Prevoid: 270cc Postvoid: 20cc

**UTERUS** is retroverted and normal in size. It has uniform myometrial echopattern.

**Endometrial thickness measures 3mm** 

Uterus measures as follows: LS: 8.2cms AP: 4.0cms TS: 5.2cms.

Few nabothian cysts are noted in the cervix, two larger ones with internal echoes measuring about  $14 \times 14$ mm.

**OVARIES** are normal in size, shape and echotexture. No focal lesion seen.

Ovaries measure as follows: **Right ovary**: 1.9 x 1.2cms **Left ovary**: 1.9 x 1.0cms

POD & adnexae are free.

No evidence of ascites/pleural effusion.

#### **IMPRESSION:**

Name	BHAVANI B S	ID	MED111023576
Age & Gender	49/FeMale		17-03-2022 00:00:00
Ref Doctor Name	MediWheel		



#### > NO SIGNIFICANT ABNORMALITY.

DR. H.K. ANAND CONSULTANT RADIOLOGISTS MS/an DR. MEERA S

Name	BHAVANI B S	ID	MED111023576
Age & Gender	49/FeMale	Visit Date	17-03-2022 00:00:00
Ref Doctor Name	MediWheel		



# X-ray mammogram (mediolateral oblique and craniocaudal views) followed by Sonomammography was performed.

#### MAMMOGRAPHY OF BOTH BREASTS

Both breasts show symmetrical fibro glandular fatty tissue.

No evidence of focal soft tissue lesion.

No evidence of cluster micro calcification.

Subcutaneous fat deposition is within normal limits.

Right axillary lymph nodes noted.

#### SONOMAMMOGRAPHY OF BOTH BREASTS

Both breasts show normal echopattern.

No evidence of focal solid / cystic areas in either breast.

No evidence of ductal dilatation.

Few lymphnodes with maintained fatty hilum are noted in both axillae.

#### **IMPRESSION:**

#### > NO SIGNIFICANT ABNORMALITY.

**ASSESSMENT: BI-RADS CATEGORY -1** 

DR. H.K. ANAND CONSULTANT RADIOLOGISTS DR. MEERA S

#### BI-RADS CLASSIFICATION

DI MIDO CEMBRITANI	1011
CATEGORY	RESULT
0	Assessment incomplete. Need additional imaging evaluation
1	Negative. Routine mammogram in 1 year recommended.
2	Benign finding. Routine mammogram in 1 year recommended.
3	Probably benign finding. Short interval follow-up suggested.
4	Suspicious. Biopsy should be considered.
5	Highly suggestive of malignancy. Appropriate action should be taken.

Name	BHAVANI B S	ID	MED111023576
Age & Gender	49/FeMale	Visit Date	17-03-2022 00:00:00
Ref Doctor Name	MediWheel		



### X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: Essentially normal study.