PID No.
 : MED111451757
 Register On : 12/01/2023 8:36 AM

 SID No.
 : 663879862
 Collection On : 12/01/2023 9:35 AM

 Age / Sex : 34 Year(s) / Male
 Report On : 12/01/2023 1:55 PM

MEDALI

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING	'O' 'Positive'		
(Blood/Agglutination)			
Complete Blood Count With - ESR			
Haemoglobin (Blood/Spectrophotometry)	10.9	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (Blood/ <i>Numeric Integration of MCV</i>)	36.9	%	42 - 52
RBC Count (Blood/Electrical Impedance)	5.75	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (Blood/Calculated)	64.2	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (Blood/Calculated)	19.0	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (Blood/Calculated)	29.6	g/dL	32 - 36
RDW-CV (Calculated)	20.9	%	11.5 - 16.0
RDW-SD (Calculated)	46.96	fL	39 - 46
Total Leukocyte Count (TC) (Blood/Electrical Impedance)	6890	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance and absorbance)	52.36	%	40 - 75
Lymphocytes (Blood/Impedance and absorbance)	33.21	%	20 - 45
Eosinophils (Blood/Impedance and absorbance)	5.83	%	01 - 06
Monocytes (Blood/Impedance and absorbance)	8.33	%	01 - 10





APPROVED BY

 PID No.
 : MED111451757
 Register On
 : 12/01/2023 8:36 AM

 SID No.
 : 663879862
 Collection On
 : 12/01/2023 9:35 AM

 Age / Sex
 : 34 Year(s) / Male
 Report On
 : 12/01/2023 1:55 PM

Printed On



Type : OP

Ref. Dr : MediWheel

Investigation	Observed	Unit	Biological
IIIVOSI.ganori	<u>Value</u>	<u>OTIM</u>	Reference Interval
Basophils	0.27	%	00 - 02
(Blood/Impedance and absorbance)			
INTERPRETATION: Tests done on Automated	Five Part cell count	er. All abnormal results are	e reviewed and confirmed microscopically.
Absolute Neutrophil count (Blood/Impedance and absorbance)	3.61	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (Blood/ <i>Impedance</i>)	2.29	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (Blood/Impedance)	0.40	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (Blood/ <i>Impedance</i>)	0.57	10^3 / μl	< 1.0
Absolute Basophil count (Blood/ <i>Impedance</i>)	0.02	10^3 / μl	< 0.2
Platelet Count (Blood/Impedance)	3.34	lakh/cu.mm	1.4 - 4.5
INTERPRETATION: Platelet count less than 1.5	5 lakhs will be confi	rmed microscopically.	
MPV (Blood/Derived from Impedance)	8.67	fL	7.9 - 13.7
PCT (Calculated)	0.29	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated ESR analyser)	08	mm/hr	< 15
BUN / Creatinine Ratio	7.7		
Glucose Fasting (FBS) (Plasma - F/Glucose oxidase/Peroxidase)	92	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 21/01/2023 3:47 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)	Negative		Negative
(Urine - F)			
Glucose Postprandial (PPBS)	140	mg/dL	70 - 140
(Dlacma DD/COD DOD)			





APPROVED BY

The results pertain to sample tested.

Page 2 of 7

 PID No.
 : MED111451757
 Register On : 12/01/2023 8:36 AM

 SID No.
 : 663879862
 Collection On : 12/01/2023 9:35 AM

 Age / Sex : 34 Year(s) / Male
 Report On : 12/01/2023 1:55 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Unit	Biological
	Value	Reference Interval

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

: 21/01/2023 3:47 PM

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Calculated)	7.0	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe ⁻ Alkaline Picrate)	0.9	mg/dL	0.9 - 1.3
Uric Acid (Serum/Uricase/Peroxidase)	6.1	mg/dL	3.5 - 7.2
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulphanilic acid)	1.3	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulphanilic acid)	0.5	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Calculated)	0.80	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC without P-5-P)	25	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC without P-5-P)	37	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC AMP Buffer)	76	U/L	53 - 128
Total Protein (Serum/Biuret)	6.9	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.8	gm/dl	3.5 - 5.2
Globulin (Serum/Calculated)	3.10	gm/dL	2.3 - 3.6





APPROVED BY

 PID No.
 : MED111451757
 Register On : 12/01/2023 8:36 AM

 SID No.
 : 663879862
 Collection On : 12/01/2023 9:35 AM

 Age / Sex : 34 Year(s) / Male
 Report On : 12/01/2023 1:55 PM



Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
A : G RATIO (Serum/ <i>Calculated</i>)	1.23		1.1 - 2.2
INTERPRETATION: Enclosure: Graph			
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	22	U/L	< 55
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	105	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol-phosphate oxidase/Peroxidase)	94	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	36	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	50.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	18.8	mg/dL	< 30





APPROVED BY

PID No. Register On : 12/01/2023 8:36 AM : MED111451757 : 663879862 SID No. Collection On : 12/01/2023 9:35 AM Age / Sex : 34 Year(s) / Male Report On : 12/01/2023 1:55 PM



Type : OP

Printed On : 21/01/2023 3:47 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Non HDL Cholesterol (Serum/Calculated)	69.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol	2.9	Optimal: < 3.3
Ratio		Low Risk: 3.4 - 4.4
(Serum/Calculated)		Average Risk: 4.5 - 7.1
		Moderate Risk: 7.2 - 11.0
		High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio	2.6	Optimal: < 2.5
(TG/HDL)		Mild to moderate risk: 2.5 - 5.0
(Serum/Calculated)		High Risk: > 5.0
LDL/HDL Cholesterol Ratio	1.4	Optimal: 0.5 - 3.0
(Serum/Calculated)		Borderline: 3.1 - 6.0
		High Risk: > 6.0

Glycosylated Haemoglobin (HbA1c)

Normal: 4.5 - 5.6 HbA1C 6.5 % Prediabetes: 5.7 - 6.4 (Whole Blood/HPLC-Ion exchange) Diabetic: \geq 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

139.85 mg/dL Mean Blood Glucose

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values. Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.





APPROVED BY

The results pertain to sample tested.

Page 5 of 7

Lab Address: MEDALL HEALTH CARE PVT LTD#17-11-3/4,DR.GKS MANSION,OFFICIAL COLONY,MAHARANI PETA. VIZAG 530002...

 PID No.
 : MED111451757
 Register On
 : 12/01/2023 8:36 AM

 SID No.
 : 663879862
 Collection On
 : 12/01/2023 9:35 AM

Age / Sex : 34 Year(s) / Male Report On : 12/01/2023 1:55 PM

Ref. Dr : MediWheel

MEDALL

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
THYROID PROFILE / TFT T3 (Triiodothyronine) - Total (Serum/Chemiluminescent Immunometric Assay (CLIA))	1.22	ng/ml	0.7 - 2.04

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 10.25 µg/dl 4.2 - 12.0

 $(Serum/{\it Chemiluminescent\ Immunometric\ Assay}$

(CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.65 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescence)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

Urine Analysis - Routine

Others

(Urine/Microscopy)

INTERPRETATION: Note: Done with Automated Urine Analyser & microscopy

Physical Examination(Urine Routine)

Colour PALE YELLOW Yellow to Amber

(Urine/Physical examination)





APPROVED BY

The results pertain to sample tested.

Page 6 of 7

Lab Address: MEDALL HEALTH CARE PVT LTD#17-11-3/4,DR.GKS MANSION,OFFICIAL COLONY,MAHARANI PETA. VIZAG 530002..

PID No. : MED111451757 Register On : 12/01/2023 8:36 AM : 663879862 SID No. Collection On : 12/01/2023 9:35 AM Age / Sex : 34 Year(s) / Male Report On : 12/01/2023 1:55 PM



Type : OP

Ref. Dr

Printed On : 21/01/2023 3:47 PM

Investigation **Observed** <u>Unit</u> **Biological Value** Reference Interval Appearance Clear Clear (Urine/Physical examination) <u>Chemical Examination(Urine Routine)</u>

: MediWheel

Protein Negative Negative

(Urine/Dipstick-Error of indicator/ Sulphosalicylic acid method)

Negative Negative Glucose

(Urine/Dip Stick Method / Glucose Oxidase -Peroxidase / Benedict s semi quantitative method.)

Microscopic Examination(Urine *Routine*)

0 - 5 Pus Cells 3-5 /hpf

(Urine/Microscopy exam of urine sediment)

NIL **Epithelial Cells** 1-2 /hpf

(Urine/Microscopy exam of urine sediment) **NIL** /hpf 0 - 5

(Urine/Microscopy exam of urine sediment)



Consultant Pathologist Reg No. 070707

APPROVED BY

-- End of Report --



Name	PERUBOINA MANI KUMAR	ID	MED111451757
Age & Gender	34Y/M	Visit Date	Jan 12 2023 8:34AM
Ref Doctor	MediWheel		

ULTRASOUND WHOLE ABDOMEN

Liver : Mildly enlarged in size (16.5 cm) shows diffuse increase in

echotexture.

There is no evidence of IHBR / EHBR dilatation seen.

No focal space occupying lesions seen.

CBD is normal. PV normal.

Gall Bladder : Normal in volume and wall thickness.

No e/o intraluminal calculi seen.

Pancreas : Head, body and tail are identified with normal echopattern and

smooth outlines.

Spleen : Measured 11.7 cm, in size with normal echotexture.

Right kidney : Measured 10.6 x 5.3 cm in size.

Left kidney : Measured 11.4 x 5.6 cm in size.

Both kidneys are normal in size, position, with well preserved cortico medullary differentiation and normal pelvicalyceal

anatomy.

No e/o calculi / space occupying lesion seen. No e/o suprarenal / retroperitoneal masses noted.

Urinary : Normal in volume and wall thickness.
bladder No e/o intraluminal calculi / masses seen.

Prostate : Measured 3.1 x 3.1 x 3.1 cm in size (Vol : 16.6 cc) with normal

echotexture.

No e/o ascites / pleural effusion seen. No e/o detectable bowel pathology seen.

IMPRESSION:

• Mild hepatomegaly with Grade II steatosis – To correlate with LFT.

- For clinical correlation.



Name	PERUBOINA MANI KUMAR	ID	MED111451757
Age & Gender	34Y/M	Visit Date	Jan 12 2023 8:34AM
Ref Doctor	MediWheel		

Dr.Jahnavi Barla MD (RD), DGO. Consultant Radiologist