PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	9.7	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	31.3	%	37 - 47
RBC Count (EDTA Blood)	4.60	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood)	68.1	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	21.2	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	31.1	g/dL	32 - 36
RDW-CV	19.1	%	11.5 - 16.0
RDW-SD	45.52	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	7500	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	46.5	%	40 - 75
Lymphocytes (Blood)	45.7	%	20 - 45
Eosinophils (Blood)	0.9	%	01 - 06





 PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood)	6.4	%	01 - 10
Basophils (Blood)	0.5	%	00 - 02
INTERPRETATION: Tests done on Automated Five Pa	art cell counter. All a	abnormal results are	reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	3.49	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	3.43	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.07	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.48	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.04	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	417	10^3 / μl	150 - 450
MPV (Blood)	7.7	fL	8.0 - 13.3
PCT (Automated Blood cell Counter)	0.32	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	19	mm/hr	< 20





 PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.26	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.12	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.14	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	16.88	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	12.31	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	16.87	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	74.1	U/L	42 - 98
Total Protein (Serum/Biuret)	7.18	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.25	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.93	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.45		1.1 - 2.2



VERIFIED BY



 PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	194.80	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	149.42	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	36.55	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	128.4	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	29.9	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	158.3	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



PID No. : MED121154909 Register On : 25/06/2022 8:46 AM : 522219850 SID No. Collection On : 25/06/2022 10:39 AM Age / Sex : 26 Year(s) / Female Report On 25/06/2022 8:00 PM **Type** : OP

Printed On

Ref. Dr : MediWheel

Investigation Observed **Unit Biological** Reference Interval **Value**

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 29/06/2022 4:20 PM

Total Cholesterol/HDL Cholesterol Ratio 5.3 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated)

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio Optimal: < 2.54.1

Mild to moderate risk: 2.5 - 5.0 (TG/HDL)

High Risk: > 5.0(Serum/Calculated)

LDL/HDL Cholesterol Ratio 3.5 Optimal: 0.5 - 3.0

Borderline: 3.1 - 6.0 (Serum/Calculated) High Risk: > 6.0



VERIFIED BY



 PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	6.5	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 139.85 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



PID No. Register On : 25/06/2022 8:46 AM : MED121154909 : 522219850 SID No. Collection On : 25/06/2022 10:39 AM Age / Sex : 26 Year(s) / Female Report On : 25/06/2022 8:00 PM **Type** : OP

Printed On

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>L</u>	<u>Biological</u>
	Value	Reference Interval

: 29/06/2022 4:20 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.38 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

4.2 - 12.0T4 (Tyroxine) - Total 9.93 μg/dl

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

 $\mu IU/mL$ 0.35 - 5.50TSH (Thyroid Stimulating Hormone) 3.42

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



 PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u>
Value Reference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION (URINE COMPLETE)

Colour Pale Yellow Yellow to Amber

(Urine)

Appearance Clear Clear

(Urine)

Volume(CLU) 20

(Urine)

CHEMICAL EXAMINATION (URINE

COMPLETE)

pH 5.0 4.5 - 8.0

(Urine)

Specific Gravity 1.012 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

(Urine)

Blood Negative Negative

Nitrite Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)





Negative

PID No. Register On : 25/06/2022 8:46 AM : MED121154909 : 522219850 SID No. Collection On : 25/06/2022 10:39 AM Age / Sex : 26 Year(s) / Female Report On 25/06/2022 8:00 PM : OP **Type Printed On** : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

Investigation **Observed** <u>Unit</u> **Biological** Reference Interval <u>Value</u> Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

MICROSCOPIC EXAMINATION (URINE COMPLETE)

NIL 0-2/hpf Pus Cells (Urine) **Epithelial Cells** 0-1/hpf **NIL**

(Urine)

Nil /hpf **NIL RBCs**

(Urine)

Nil Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.





 PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'O' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



VERIFIED BY



 PID No.
 : MED121154909
 Register On
 : 25/06/2022 8:46 AM

 SID No.
 : 522219850
 Collection On
 : 25/06/2022 10:39 AM

 Age / Sex
 : 26 Year(s) / Female
 Report On
 : 25/06/2022 8:00 PM

 Type
 : OP
 Printed On
 : 29/06/2022 4:20 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	12.4		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	100.88	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)	Negative		Negative
(Urine - F/GOD - POD)			
Glucose Postprandial (PPBS)	82.07	mg/dL	70 - 140
(Plasma - PP/GOD-PAP)			

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti-diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	8.1	mg/dL	7.0 - 21
(Serum/ <i>Urease UV / derived</i>)			
Creatinine	0.65	mg/dL	0.6 - 1.1
(Serum/Modified Jaffe)			

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine

Uric Acid 3.94 mg/dL 2.6 - 6.0

(Serum/Enzymatic)



VERIFIED BY



APPROVED BY

-- End of Report --

Name	MS.MANASA	ID	MED121154909
Age & Gender	26Y/FEMALE	Visit Date	25 Jun 2022
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (12.4 cm) and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER show normal shape and has clear contents. Wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN show normal shape, size (9.5 cm) and echopattern.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cm)	Parenchymal thickness (cm)
Right Kidney	10.9	1.7
Left Kidney	12.1	1.7

URINARY BLADDER show normal shape and wall thickness. It has clear contents. No evidence of diverticula.

UTERUS is anteverted and has normal shape and size. It has uniform myometrial echopattern. Endometrial echo is of normal thickness - 5.2 mm.

Uterus measures LS: 5.4 cm AP: 3.3 cm TS: 4.8 cm.

OVARIES are normal in size, shape and echotexture

Right ovary measures 2.7 x 1.8 cm Left ovary measures 2.8 x 1.5 cm.

POD & adnexa are free.

No evidence of ascites.

IMPRESSION:

• No significant abnormality detected.

Name	MS.MANASA	ID	MED121154909
Age & Gender	26Y/FEMALE	Visit Date	25 Jun 2022
Ref Doctor Name	MediWheel		

DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST

Hn/mj

Name	MS.MANASA	ID	MED121154909
Age & Gender	26Y/FEMALE	Visit Date	25 Jun 2022
Ref Doctor Name	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

• No significant abnormality detected.

DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST

Hn/mj