Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787


Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



### **General Physical Examination**

Date of Examination: 08-10-2022
Name: Ram Singh Age: 58 DOB: 09-07-1964 Sex: Male
Referred By: BOB (Mcoliwheel)
Photo ID: AADHAR ID#: catterned.
Ht: 172 (cm) Wt: 85 (Kg)
Chest (Expiration): 48 (cm) Abdomen Circumference: 102 (cm)
Blood Pressure: 131/98 mm Hg PR: 80 / min RR: 18 / min Temp: Afternile
BMI 28-7
Eye Examination: Dis Vision 6/9 (with specs B/c eyes)  Near Vision M/6 (with specs)  Near Vision M/6 (with specs)
Other: Dot significant-
On examination he/she appears physically and mentally fit: Yes / No
Signature Of Examine: Name @Examinee: D.M.R.D.  N.B. B. S. S. No017996  RMC Rag. No017996
Signature Medical Examiner : Name Medical Examiner






पता: S/O नर सिंह, प.न.४४, माँ करनी विहार, कालवाड रोड, झोटवाडा, जयपुर, राजस्थान - 302012

Address: S/O Nar Singh, p.n.44, maa kami vihar, kalwad road, JHOTWARA, Jaipur, Rajasthan - 302012

9442 8426 4956 VID: 9175 0556 3355 4160



help@uidai.gcv.in



www. www.uidai.gov.in Allengers ECG (Pisces)(PIS212160118) 102220585 / MR RAMSINGH / 58 Yrs / M/ Non Smoker
Heart Rate: 72 bpm / / Refd By.: BOB / Tested On: 08-Oct-22 11:31:59 / HF 0.05 Hz - LF 100 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s **V2** 50 5

DR. GOYALS PATH LAB & IMAGING CENTER

ECG



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sangant Final Januar Tele: 0141-2293346, 4049787, 9887049787

Date :- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male 58 Yrs 3 Mon 1 Days

Company :- MediWheel

Patient ID :-12222717

Ref. By Doctor:-

Lab/Hosp :-



Final Authentication: 08/10/2022 12:23:19

BOB PACKAGE ABOVE 40MALE

#### USG WHOLE ABDOMEN

Liver is mild enlarge size(14.5cm). Echo-texture is minimal bright. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

**Pancreas** is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

**Kidneys** are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

**Urinary bladder** is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Pre void - 259ml, Post void - 36(Insignificant)

Prostate is moderate enlarge in size( 42gms) with normal echo-texture and outline.

No enlarged nodes are visualised. No retro-peritoneal lesion is identified No significant free fluid is seen in peritoneal cavity.

#### IMPRESSION:

\*Mild hepatomegaly with early fatty changes.

\*Moderate prostatomegaly.

Needs clinical correlation for further evaluation

\*\*\* End of Report \*\*\*

Page No: 1 of 1

**GEETASAINI** 

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996 Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495 Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163 Transcript by.



MITRAL VALVE

## Dr. Goyal's

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

#### 2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

TRICUSPID VALVE

AORTIC VALVE NORN		MAL	PULMO	PULMONARY VALVE			NORMAL		
			1	M.MODE EXAMI	TATION:			alles a constant	
AO	26	mm	LA	36	Mm	IVS-D	09	mm	
IVS-S	12	mm	LVID	45	Mm	LVSD	28	mm	
LVPW-D	09	mm	LVPW-S	15	Mm	RV		mm	
RVWT	66%	mm	EDV		MI	LVVS		ml	
LVEF				RWMA		ABSENT			

NORMAL

**CHAMBERS:** 

LA	NORMAL	RA	NORMAL
LV	NORMAL	RV	NORMAL
PERICARDIUM		NORMAL	

#### COLOUR DOPPLER:

	MITRA	AL VALVE					
E VELOCITY	0.90	m/sec	PEAK	PEAK GRADIENT			Mm/hg
A VELOCITY	0.72	m/sec	MEA	N GRADIEN	T		Mm/hg
MVA BY PHT		Cm2	MVA	BY PLANIN	IETRY		Cm2
MITRAL REGURGITATION	ON				ABSENT		
	AORT	C VALVE					
PEAK VELOCITY	1.3	m/	sec	PEAK G	RADIENT		mm/hg
AR VMAX		m/	sec	MEAN	GRADIENT		mm/hg
AORTIC REGURGITATION	ON			ABSENT			
	TRICUS	PID VALVE					
PEAK VELOCITY	0.3	5	m/sec	PEAK G	RADIENT		mm/hg
MEAN VELOCITY			m/sec	MEAN	GRADIENT		mm/hg
VMax VELOCITY							
TRICUSPID REGURGITATION			ABSENT				
	PULM	ONARY VAI	LVE				
PEAK VELOCITY		0.98		M/sec.	PEAK GRADI	ENT	Mm/hg
MEAN VALOCITY					MEAN GRAD	IENT	Mm/hg
PULMONARY REGURO	SITATION				ABSENT		***************************************

### Impression--

- Normal LV size & contractility
- No RWMA, LVEF 66 %.
- Normal cardiac chamber.
- Normal valve
- No clot, no vegetation, no pericardial effusion. (Cardiologist)

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996 Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495 Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant
FMF ID - 260517 | RMC No 22430

Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163

NORMAL

Transcript by.

## Dr. Goya

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male

Sample Type :- EDTA

58 Yrs 3 Mon 1 Days

Company :- MediWheel

Patient ID :-12222717

Ref. By Dr:-

Lab/Hosp :-

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:50:11

#### HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BOB PACKAGE ABOVE 40MALE			
HAEMOGARAM		¥.	
HAEMOGLOBIN (Hb)	13.9	- / IT	12.2 90 0.0000
TOTAL LEUCOCYTE COUNT		g/dL	13.0 - 17.0
DIFFERENTIAL LEUCOCYTE COUNT	6.71	/cumm	4.00 - 10.00
NEUTROPHIL			
	50.0	%	40.0 - 80.0
LYMPHOCYTE	41.2 H	% .	20.0 - 40.0
EOSINOPHIL	5.5	%	1.0 - 6.0
MONOCYTE	3.1	%	2.0 - 10.0
BASOPHIL	0.2	%	0.0 - 2.0
NEUT#	3.36	10^3/uL	1.50 - 7.00
LYMPH#	2.77	10^3/uL	1.00 - 3.70
EO#	0.37	10^3/uL	0.00 - 0.40
MONO#	0.20	10^3/uL	0.00 - 0.70
BASO#	0.01	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	4.73	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	39.90 └	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	84.3	fL	83.0 - 101.0
MEAN CORP HB (MCH)	29.4	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	34.4	g/dL	31.5 - 34.5
PLATELET COUNT	135 └	x10^3/uL	150 - 410
RDW-CV	14.0	%	
MENTZER INDEX	17.82	70	11.6 - 14.0

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

**AJAYSINGH Technologist** 

Page No: 1 of 13



## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male

58 Yrs 3 Mon 1 Days

Company :- MediWheel

Patient ID :-12222717

Ref. By Dr:-

Lab/Hosp :- '

Sample Type :- EDTA

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:50:11

HAEMATOLOGY

**Test Name** 

Unit

**Biological Ref Interval** 

Erythrocyte Sedimentation Rate (ESR)

28 H

Value

mm/hr.

00 - 13

(ESR) Methodology: Measurment of ESR by cells aggregation.

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation : ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia (C BC) in the thodology of the the thodology of the thodology of

**AJAYSINGH Technologist** 

Page No: 2 of 13



## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

58 Yrs 3 Mon 1 Days

Patient ID :-12222717 Ref. By Dr:-

Lab/Hosp :-

Sex / Age :- Male Company :- MediWheel

**Test Name** 

Sample Type :- EDTA, KOx/Na FLUORIDE-F, KSax/Nat-FCbl@RiezETTPRe DBITME2022 10:06:22

HAEMATOLOGY

Final Authentication: 08/10/2022 15:15:21

Value

**Biological Ref Interval** 

BLOOD GROUP ABO

"B" POSITIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone)

FASTING BLOOD SUGAR (Plasma)

Method:- GOD PAP

110.7

mg/dl

75.0 - 115.0

Impaired glucose tolerance (IGT) 111 - 125 mg/dL Diabetes Mellitus (DM) > 126 mg/dL

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases

BLOOD SUGAR PP (Plasma)

Method:- GOD PAP

153.8 H

mg/dl

70.0 - 140.0

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil

AJAYSINGH, KAUSHAL, POOJABOHRA **Technologist** DR.HANSA Page No: 3 of 13



Dr. Piyush Goyal (D.M.R.D.) Dr. Rashmi Bakshi Dr. Chandrika Gupta

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male

Company :- MediWheel

58 Yrs 3 Mon 1 Days

Ref. By Dr:-

Lab/Hosp:-

Patient ID :-12222717

Sample Type :- PLAIN/SERUM

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:41:27

BIOCHEMISTR	

	DIOCILLIA		
Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	139.42	mg/dl	Desirable <200 Borderline 200-239 High> 240
TRIGLYCERIDES Method:- GPO-PAP	135.68	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	37.49	mg/dl	Low < 40 High > 60
BONCO DIAGRAM WAS A DESCRIPTION OF THE SECOND OF THE SECON	79.32	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	27.14	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	3.72		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	2.12		0.00 - 3.50
TOTAL LIPID Method:-CALCULATED TOTAL CHOLESTEROL InstrumentName:Randox Rx Imola Int	469.61	mg/dl	400.00 - 1000.00

terol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and

DIRECT HDLCHOLESTERO InstrumentName: Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROLInstrumentName: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid crosclerosis or reduce its progress and to avoid plaque rupture TOTAL LIPID AND VLDL ARE CALCULATED

**MKSHARMA** 

Page No: 5 of 13



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male

58 Yrs 3 Mon 1 Days

Company :- MediWheel Sample Type :- PLAIN/SERUM Patient ID: -12222717

Ref. By Dr:-

Lab/Hosp :- .

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13 41.27

DI	10	0	т	EN	A	ST	CD	1
D			п.	1,11	V 1		ıĸ	·

BIOCHEMISTRY							
Test Name	Value	Unit	Biological Ref Interval				
LIVER PROFILE WITH GGT	9						
SERUM BILIRUBIN (TOTAL) Method:-Colorimetric method	0.88	mg/dl	Up to - 1.0 Cord blood <2 mg/dL Premature < 6 days <16mg/dL Full-term < 6 days= 12 mg/dL 1month - <12 months <2 mg/dL 1-19 years <1.5 mg/dL Adult - Up to - 1.2 Ref-(ACCP 2020)				
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.36	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL				
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.52	mg/dl	0.30-0.70				
SGOT Method:- IFCC	18.2	U/L	Men- Up to - 37.0 Women - Up to - 31.0				
SGPT Method:- IFCC	19.1	U/L	Men- Up to - 40.0 Women - Up to - 31.0				
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	78.90	IU/L	30.00 - 120.00				
SERUM GAMMA GT Method:- IFCC	· 32:10	U/L ·	11.00 - 50.00				
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.25	g/dl	6.40 - 8.30				
SERUM ALBUMIN Method:- Bromocresol Green	4.71	g/dl	3.80 - 5.00				
SERUM GLOBULIN Method:- CALCULATION	2.54	gm/dl	2.20 - 3.50				
A/G RATIO	1.85		1.30 - 2.50				

Total Bilirubin Methodology: Colorimetric method InstrumentName: Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName:Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular destrophy and organ damage.

Alkaline Phosphatase Methodology: AMP Buffer InstrumentName: Randox Rx Imola Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of TOTAL PROTEIN Methodology: Birret Reagent InstrumentName: Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology: Birret Reagent InstrumentName: Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName:Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving

**MKSHARMA** 

Page No: 6 of 13



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Sex / Age :- Male

Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

58 Yrs 3 Mon 1 Days

Lab/Hosp :-

Company :- MediWheel Sample Type :- PLAIN/SERUM

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:41:27

#### DIOCHEMICTRY

BIOCHEMISTRY					
Test Name	Value	Unit	Biological Ref Interval		
SERUM CREATININE Method:- Colorimetric Method	-1.12	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20		
SERUM URIC ACID Method:- Enzymatic colorimetric	6.69	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7		

Ref. By Dr:-

Patient ID :-12222717

**MKSHARMA** 

Page No: 8 of 13



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13 Date

NAME :- Mr. RAMSINGH

Patient ID :-12222717

Ref. By Dr:-

Lab/Hosp :-

Company :- MediWheel

Sex / Age :- Male

Sample Type :- PLAIN/SERUM

58 Yrs 3 Mon 1 Days

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:41:27

#### RIOCHEMISTRY

	DIOCHEN			
Test Name	Value	Unit	Biological Re	f Interval
BLOOD UREA NITROGEN (BUN)	16.5	mg/dl	0.0 - 23.0	

**MKSHARMA** 

Page No: 9 of 13



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Sample Type :- EDTA

Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13 Date

NAME :- Mr. RAMSINGH

Sex / Age :- Male

Company :- MediWheel

58 Yrs 3 Mon 1 Days

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:50:11

HAEMATOLOGY

**Test Name** Value Unit Biological Ref Interval GLYCOSYLATED HEMOGLOBIN (HbA1C) 6.1 H 9/0 Non-diabetic: < 5.7 Method:- HPLC Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0 Action suggested: > 6.5

Ref. By Dr:-

Lab/Hosp :-

Patient ID: -12222717

Instrument name: ARKRAY'S ADAMS Lite HA 8380V, JAPAN

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb.High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1cmeasurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

MEAN PLASMA GLUCOSE Method:- Calculated Parameter

128 H

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

**AJAYSINGH** Technologist

Page No: 10 of 13



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male 58 Yrs 3 Mon 1 Days

Company :- MediWheel

Sample Type :- URINE

Patient ID: -12222717

Ref. By Dr:-

Lab/Hosp :-

Final Authentication: 08/10/2022 12:08:29

**CLINICAL PATHOLOGY** 

Sample Collected Time 08/10/2022 10:06:22

**Test Name** Value Unit **Biological Ref Interval** 

**Urine Routine** 

NITRITE

PHYSICAL EXAMINATION

COLOUR PALE YELLOW PALE YELLOW **APPEARANCE** Clear Clear

CHEMICAL EXAMINATION

REACTION(PH) 5.5 5.0 - 7.5SPECIFIC GRAVITY 1.015 1.010 - 1.030 **PROTEIN** NIL. NIL **SUGAR** NIL NIL BILIRUBIN NEGATIVE **NEGATIVE** UROBILINOGEN NORMAL NORMAL KETONES **NEGATIVE NEGATIVE** 

·NEGATIVE

**NEGATIVE** MICROSCOPY EXAMINATION RBC/HPF NIL /HPF NIL WBC/HPF 2-3 /HPF 2-3 EPITHELIAL CELLS 2-3 /HPF 2-3 CRYSTALS/HPF ABSENT **ABSENT** CAST/HPF ABSENT ABSENT AMORPHOUS SEDIMENT ABSENT ABSENT BACTERIAL FLORA ABSENT ABSENT YEAST CELL ABSENT ABSENT **OTHER** ABSENT

**POOJABOHRA** Technologist DR.HANSA Page No: 11 of 13



## Dr. Goya

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



:- 08/10/2022 09:55:13 Date

NAME :- Mr. RAMSINGH

Ref. By Dr:-

Patient ID: -12222717

Sex / Age :- Male 58 Yrs 3 Mon 1 Days Lab/Hosp :-

Company :- MediWheel Sample Type :- PLAIN/SERUM

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:02:42

#### **IMMUNOASSAY**

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.320	ng/ml	0.600 - 1.810
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	7.130	ug/dl	4.500 - 10.900
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	1.840	μIU/mL	0.550 - 4.780

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease.T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions; such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation (TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

#### INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	. 0.30-3.00

NARENDRAKUMAR **Technologist** 

Page No: 12 of 13



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur - 302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male

58 Yrs 3 Mon 1 Days

Ref. By Dr:-

Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sample Collected Time 08/10/2022 10:06:22

Final Authentication: 08/10/2022 13:02:42

**IMMUNOASSAY** 

**Test Name** 

TOTAL PSA

Method:- Chemiluminescence

Value

Unit

Patient ID: -12222717

**Biological Ref Interval** 

1.130

ng/ml

0.000 - 4.000

InstrumentName: ADVIA CENTAUR CP Interpretation: Elevated serum PSA concentrations are found in men with prostate cancer, benign prostatic hypertrophy (BHP) or inflammatory conditions of other adjacent genitourinary tissues, but not in apparently healthy men or in men with cancers other than prostate cancer PSA has been demonstrated to be an accurate marker for monitoring advancing clinical stage in untreated patients and for monitoring response to therapy by radical prostatectomy, radiation therapy and anti-androgen therapy. PSA is also important in determining the potential and actual effectiveness of surgery or other therapies. Progressive disease is defined by an increase of at least 25%. Sampling should be repeated within two to four weeks for additional evidence. Different assay methods cannot be used interchangeably.

\*\*\* End of Report \*\*\*

NARENDRAKUMAR **Technologist** 

Page No: 13 of 13





B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 08/10/2022 09:55:13

NAME :- Mr. RAMSINGH

Sex / Age :- Male 58 Yrs 3 Mon 1 Days

Company :- MediWheel

Patient ID :-12222717

Ref. By Doctor:-

Lab/Hosp :-

Final Authentication: 08/10/2022 14:12:48

BOB PACKAGE ABOVE 40MALE

### X RAY CHEST PA VIEW:

### Bronchovascular markings are prominent.

Otherwise lung fields are clear.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Aortic knuckle calcification is seen.

(Please correlate clinically and with relevant further investigations.)

\*\*\* End of Report \*\*\*

Page No: 1 of 1

Dr. Piyush Goyal
M.B.B.S., D.M.R.D.
RMC Reg No. 017996

Dr. Poon
MBBS, MD R
RMC N

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Ashish Choudhary
MBBS, MD (Radio Diagnosis)
Fetal Medicine Consultant
FMF ID - 260517 | RMC No 22430

Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163

Dr. Piyush Goyal

Transcript by.

BILAL

