PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	14.18	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	43.2	%	42 - 52
RBC Count (EDTA Blood)	5.10	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	84.7	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	27.8	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	32.8	g/dL	32 - 36
RDW-CV	14.0	%	11.5 - 16.0
RDW-SD	41.50	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	9810	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	44.72	%	40 - 75
Lymphocytes (Blood)	38.72	%	20 - 45
Eosinophils (Blood)	8.63	%	01 - 06





 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood)	7.74	%	01 - 10
Basophils (Blood)	0.19	%	00 - 02
INTERPRETATION: Tests done on Automated Five P	art cell counter. All	abnormal results are	reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	4.39	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	3.80	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.85	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.76	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.02	$10^3 / \mu l$	< 0.2
Platelet Count (EDTA Blood)	194.6	$10^3 / \mu l$	150 - 450
MPV (Blood)	9.08	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.18	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	13	mm/hr	< 15





 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.37	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.14	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.23	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	13.37	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	17.62	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	18.03	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	66.4	U/L	53 - 128
Total Protein (Serum/Biuret)	6.91	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.60	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.31	gm/dL	2.3 - 3.6
A: G RATIO (Serum/Derived)	1.99		1.1 - 2.2



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	183.41	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	168.26	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	29.10	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	120.6	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >=190
VLDL Cholesterol (Serum/Calculated)	33.7	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	154.3	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u> <u>Value</u> <u>Reference Interval</u>

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio 6.3 Optimal: < 3.3 (Serum/Calculated) Low Risk: 3.4 - 4.4

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 5.8 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio 4.1 Optimal: 0.5 - 3.0

(Serum/Calculated)
Borderline: 3.1 - 6.0
High Risk: > 6.0



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	5.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 105.41 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
_	Value	Reference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.27 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Tyroxine) - Total 8.84 µg/dl 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.60 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION (URINE COMPLETE)

Colour Yellow Yellow Yellow to Amber

(Urine)

Appearance Clear Clear

(Urine)

Volume(CLU) 15

(Urine)

CHEMICAL EXAMINATION (URINE

<u>COMPLETE</u>)

pH 5 4.5 - 8.0

(Urine)

Specific Gravity 1.012 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)

Nitrite Negative Negative

(Urine)

Bilirubin Negative Negative Negative

(Urine)

Protein Negative Negative

(Urine)



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

<u>Investigation</u> <u>Observed</u> <u>Unit</u> <u>Biological</u> <u>Value</u> <u>Reference Interval</u>

Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

(Urine)

MICROSCOPIC EXAMINATION (URINE COMPLETE)

Pus Cells 0-1 /hpf NIL

(Urine)

Epithelial Cells **0-1** /hpf NIL

(Urine)

RBCs Nil /HPF NIL

(Urine)

Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'A' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	12.5		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	81.48	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)	Negative		Negative
(Urine - $F/GOD - POD$)			
Glucose Postprandial (PPBS)	62.03	mg/dL	70 - 140

(Plasma - PP/GOD-PAP)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	16.8	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	1.34	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc

Uric Acid 7.24 mg/dL 3.5 - 7.2

(Serum/Enzymatic)



VERIFIED BY



 PID No.
 : MED121139086
 Register On
 : 18/06/2022 9:27 AM

 SID No.
 : 522219368
 Collection On
 : 18/06/2022 11:36 AM

 Age / Sex
 : 48 Year(s) / Male
 Report On
 : 18/06/2022 4:21 PM

 Type
 : OP
 Printed On
 : 29/06/2022 2:43 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Prostate specific antigen - Total(PSA) (Serum/Manometric method)	1.45	ng/ml	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0



VERIFIED BY



APPROVED BY

-- End of Report --

Name	MR.PASHA SHAIK MUNAWAR	ID	MED121139086
Age & Gender	48Y/MALE	Visit Date	18 Jun 2022
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (13.9 cms) and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is partially distended. CBD is not dilated.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size (11.4 cms) and echopattern.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

·	Bipolar length (cm)	Parenchymal thickness (cm)
Right Kidney	10.4	2.1
Left Kidney	9.4	1.5

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern. It measures 3.5 x 2.9 x 4.9 cm, volume: 25cc.

No evidence of ascites.

IMPRESSION:

• No significant abnormality detected.

DR. HEMANANDINI V.N

CONSULTANT RADIOLOGIST

Hn/Ss

Name	MR.PASHA SHAIK MUNAWAR	ID	MED121139086
Age & Gender	48Y/MALE	Visit Date	18 Jun 2022
Ref Doctor Name	MediWheel		

Name	MR.PASHA SHAIK MUNAWAR	ID	MED121139086
Age & Gender	48Y/MALE	Visit Date	18 Jun 2022
Ref Doctor Name	MediWheel		

2D ECHOCARDIOGRAPHIC STUDY

M-mode measurement:

AORTA 3.19 cms. LEFT ATRIUM : 3.50 cms. AVS 1.48 cms. LEFT VENTRICLE (DIASTOLE) 4.78 cms. (SYSTOLE) 2.57 cms. **VENTRICULAR SEPTUM** (DIASTOLE) 1.34 cms. (SYSTOLE) 1.80 cms. **POSTERIOR WALL** (DIASTOLE) 1.39 cms. (SYSTOLE) 1.74 cms. **EDV** 106 ml. **ESV** 23 ml. FRACTIONAL SHORTENING 30 % **EJECTION FRACTION** 60 % : **EPSS** cms. **RVID** 1.80 cms.

DOPPLER MEASUREMENTS:

MITRAL VALVE: $E - 0.8 \text{ m/s} \quad A - 0.7 \text{ m/s} \quad \text{MILD MR}.$

AORTIC VALVE: 1.1 m/s NO AR.

TRICUSPID VALVE: E - 0.4 m/s A - 0.3 m/s, MILD TR, PASP 10mmHg.

PULMONARY VALVE: 0.8 m/s NO PR.

Name	MR.PASHA SHAIK MUNAWAR	ID	MED121139086
Age & Gender	48Y/MALE	Visit Date	18 Jun 2022
Ref Doctor Name	MediWheel		

2D ECHOCARDIOGRAPHY FINDINGS:

Left Ventricle : Concentric LVH, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal.Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- CONCENTRIC LVH.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- MILD MR.
- MILD TR, PASP 10mmHg.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

Name	MR.PASHA SHAIK MUNAWAR	ID	MED121139086
Age & Gender	48Y/MALE	Visit Date	18 Jun 2022
Ref Doctor Name	MediWheel		

DR. ANAND KUMAR M MD DM CONSULTANT INTERVENTIONAL CARDIOLOGIST

Name	MR.PASHA SHAIK MUNAWAR	ID	MED121139086
Age & Gender	48Y/MALE	Visit Date	18 Jun 2022
Ref Doctor Name	MediWheel		

X - RAY CHEST PA VIEW

Bilateral costophrenic angles are blunted.

Rest of the lungs appear clear.

Cardiac size is within normal limits. Unfolding of aorta is seen.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm are normal.

Visualised bones and soft tissues appear normal.

Suggested: Clinical correlation.

DR. HEMANANDINI. V.N

CONSULTANT RADIOLOGIST

Hn/ra