PID No.
 : MED120924626
 Register On
 : 26/03/2022 9:07 AM

 SID No.
 : 522212698
 Collection On
 : 26/03/2022 10:49 AM

MEDALL

Type : OP

Ref. Dr : MediWheel

Age / Sex : 42 Year(s) / Female

OP **Printed On** : 30/03/2022 5:50 PM

Report On

: 27/03/2022 3:40 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	12.9	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	38.8	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.45	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	87.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	28.8	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood'Derived from Impedance)	33.1	g/dL	32 - 36
RDW-CV (Derived from Impedance)	14.3	%	11.5 - 16.0
RDW-SD (Derived from Impedance)	44.2	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	5100	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	46.8	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	42.6	%	20 - 45
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	1.0	%	01 - 06



 PID No.
 : MED120924626
 Register On
 : 26/03/2022 9:07 AM

 SID No.
 : 522212698
 Collection On
 : 26/03/2022 10:49 AM

Printed On

Type : OP

Ref. Dr : MediWheel

MEDALI

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes	9.1	%	01 - 10
(Blood/Impedance Variation & Flow Cytometry)	0. #	0/	00.00
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.5	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.39	10^3 / μΙ	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.17	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.05	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.46	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.03	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	261	10^3 / μl	150 - 450
MPV (Blood/Derived from Impedance)	10.0	fL	8.0 - 13.3
PCT (Automated Blood cell Counter)	0.260	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	18	mm/hr	< 20

: 30/03/2022 5:50 PM



 PID No.
 : MED120924626
 Register On
 : 26/03/2022 9:07 AM

 SID No.
 : 522212698
 Collection On
 : 26/03/2022 10:49 AM

 Age / Sex
 : 42 Year(s) / Female
 Report On
 : 27/03/2022 3:40 PM

Printed On

MEDALI

Type : OP

Ref. Dr	: MediWheel			

: 30/03/2022 5:50 PM

<u>Investigation</u>	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.8	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.3	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.5	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.6	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.5	gm/dL	3.5 - 5.2
Globulin (Serum/Derived)	3.1	gm/dL	2.3 - 3.6
A: G Ratio (Serum/Derived)	1.5		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	15	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	17	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	75	U/L	42 - 98
GGT(Gamma Glutamyl Transpeptidase) (Serum/SZASZ standarised IFCC)	13	U/L	< 38



VERIFIED BY



 PID No.
 : MED120924626
 Register On
 : 26/03/2022 9:07 AM

 SID No.
 : 522212698
 Collection On
 : 26/03/2022 10:49 AM

 Age / Sex
 : 42 Year(s) / Female
 Report On
 : 27/03/2022 3:40 PM

Printed On

MEDALL

Type : OP

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>

30/03/2022 5:50 PM

Lipid Profile 198 Cholesterol Total mg/dL Optimal: < 200 Borderline: 200 - 239 (Serum/Cholesterol oxidase/Peroxidase) High Risk: ≥ 240 Triglycerides 104 mg/dL Optimal: < 150 Borderline: 150 - 199 (Serum/Glycerol phosphate oxidase / peroxidase) High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	48	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	129.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	20.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	150.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



PID No. : MED120924626 Register On : 26/03/2022 9:07 AM

: 522212698 SID No. Collection On : 26/03/2022 10:49 AM

Printed On

Age / Sex : 42 Year(s) / Female Report On : 27/03/2022 3:40 PM

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
-	<u>Value</u>		Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 30/03/2022 5:50 PM

Total Cholesterol/HDL Cholesterol Ratio 4.1 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated)

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 2.2 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

High Risk: > 5.0(Serum/Calculated)

LDL/HDL Cholesterol Ratio 2.7 Optimal: 0.5 - 3.0 (Serum/Calculated)

Borderline: 3.1 - 6.0 High Risk: > 6.0



VERIFIED BY



 PID No.
 : MED120924626
 Register On
 : 26/03/2022 9:07 AM

 SID No.
 : 522212698
 Collection On
 : 26/03/2022 10:49 AM

0.49 AW

: 30/03/2022 5:50 PM

Printed On

PM

Type : OP

Investigation

Ref. Dr : MediWheel

Observed Unit Biological

Glycosylated Haemoglobin (HbA1c)

HbA1C **5.8** % Normal: 4.5 - 5.6

(Whole Blood/HPLC) Prediabetes: 5.7 - 6.4

<u>Value</u>

Diabetic: >= 6.5

Reference Interval

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 119.76 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



PID No. : MED120924626 Register On : 26/03/2022 9:07 AM

: 522212698 SID No. Collection On : 26/03/2022 10:49 AM

Age / Sex : 42 Year(s) / Female Report On : 27/03/2022 3:40 PM

Printed On

Type : OP

Ref. Dr

: MediWheel

: 30/03/2022 5:50 PM



Investigation	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
•	<u>Value</u>		Reference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.24 ng/mL 0.7 - 2.04

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

4.2 - 12.0T4 (Thyroxine) - Total 8.06 μg/dL

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

 $\mu IU/mL$ 0.35 - 5.50TSH (Thyroid Stimulating Hormone) 2.75

(Serum/Chemiluminescent Microparticle

Immunoassay(CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



PID No. : MED120924626 **Register On** : 26/03/2022 9:07 AM : 522212698 **Collection On** : 26/03/2022 10:49 AM SID No.

Age / Sex : 42 Year(s) / Female Report On : 27/03/2022 3:40 PM

Type

: OP

Printed On : 30/03/2022 5:50 PM

Ref. Dr : MediWheel



Investigation CLINICAL PATHOLOGY PHYSICAL EXAMINATION	Observed <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Colour (Urine)	Pale Yellow		
Appearance (Urine)	Clear		Clear
Volume (Urine)	20	mL	
<u>CHEMICAL EXAMINATION(Automated-Urineanalyser)</u>			
pH (Urine/AUTOMATED URINANALYSER)	6.0		4.5 - 8.0
Specific Gravity	1.015		1.002 - 1.035

(Urine/AUTOMATED URINANALYSER)		
Specific Gravity (Urine)	1.015	1.002 - 1.035
Ketones (Urine)	Negative	Negative
Urobilinogen (Urine/AUTOMATED URINANALYSER)	0.2	0.2 - 1.0
Blood (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Nitrite (Urine/AUTOMATED URINANALYSER)	Negative	Negative
Bilirubin (Urine/AUTOMATED URINANALYSER)	Negative	Negative

Negative



Protein (Urine)



Negative

 PID No.
 : MED120924626
 Register On
 : 26/03/2022 9:07 AM

 SID No.
 : 522212698
 Collection On
 : 26/03/2022 10:49 AM

 Age / Sex
 : 42 Year(s) / Female
 Report On
 : 27/03/2022 3:40 PM

Printed On



Type : OP

Ref. Dr : MediWheel

Investigation	Observed	Linit	Piological

: 30/03/2022 5:50 PM

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine)	Negative		Negative
Leukocytes (Urine)	Negative	leuco/uL	Negative
MICROSCOPY(URINE DEPOSITS)			
Pus Cells (Urine/Flow cytometry)	2-4	/hpf	3-5
Epithelial Cells (Urine)	2-4	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	NIL
Others (Urine)	Nil		Nil





SID No.

PID No. : MED120924626 **Register On** : 26/03/2022 9:07 AM

: 522212698 Collection On : 26/03/2022 10:49 AM

Age / Sex : 42 Year(s) / Female Report On

: 27/03/2022 3:40 PM Type : OP

Ref. Dr : MediWheel

: 30/03/2022 5:50 PM **Printed On**



Investigation <u>Observed</u> <u>Unit</u> **Biological** Reference Interval **Value**

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'O' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



VERIFIED BY



PID No. Register On : MED120924626 : 26/03/2022 9:07 AM : 522212698 Collection On : 26/03/2022 10:49 AM SID No. Age / Sex : 42 Year(s) / Female

Report On

Type : OP

Ref. Dr

: MediWheel

27/03/2022 3:40 PM

Printed On 30/03/2022 5:50 PM

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	7.14		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	92	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Fasting - Urine (Urine - F)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD - POD)	75	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Glucose Postprandial - Urine (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	5	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.7	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists, N-acetylcyteine , chemotherapeutic agent such as flucytosine

Uric Acid 5.3 2.6 - 6.0 mg/dL

(Serum/Uricase/Peroxidase)



VERIFIED BY



APPROVED BY

-- End of Report --

Name	SIPRA	ID	MED120924626
Age & Gender	42/FeMale		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is collapsed and shows calculi in the lumen, largest measuring about 9 - 10mm causing posterior acoustic shadowing. Wall thickness is normal. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN show normal shape, size (9.2 cms) and echopattern.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

THE HIGHE HIGHER WE TO HE WE TO HE HE				
	Bipolar length (cm)	Parenchymal thickness (cm)		
Right Kidney	9.9	1.7		
Left Kidney	10.5	2.0		

URINARY BLADDER show normal shape and wall thickness. It has clear contents. No evidence of diverticula.

UTERUS is anteverted and has normal shape and size. It has uniform myometrial echopattern. Endometrial echo is of normal thickness - 6.5 mm.

Uterus measures LS: 7.0 cm AP: 4.1 cm TS: 5.9 cm.

OVARIES are normal in size, shape and echotexture

POD & adnexa are free.

No evidence of ascites.

IMPRESSION:

• Collapsed gall bladder with calculi in the lumen. No biliary dilatation.

Name	SIPRA	ID	MED120924626
Age & Gender	42/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



DR. H.K. ANAND DR. C.R RAMACHANDRA DR. LOHITH H.P DR. VARSHA KALE CONSULTANT RADIOLOGISTS

Lh/Ss

Name	SIPRA	ID	MED120924626
Age & Gender	42/FeMale		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHIC STUDY

M-mode measurement:

AORTA 2.65 cms. LEFT ATRIUM 3.04 cms. AVS 1.45 cms. LEFT VENTRICLE (DIASTOLE) 4.82 cms. (SYSTOLE) 2.65 cms. **VENTRICULAR SEPTUM** (DIASTOLE) 1.01 cms. (SYSTOLE) 1.35 cms. POSTERIOR WALL (DIASTOLE) 1.01 cms. (SYSTOLE) 1.35 cms. **EDV** 108 ml. **ESV** 25 ml. FRACTIONAL SHORTENING 30 % **EJECTION FRACTION** % 60

DOPPLER MEASUREMENTS:

EPSS

RVID

MITRAL VALVE: E - 0.9 m/s A - 0.7 m/s NO MR.

cms.

cms.

1.80

AORTIC VALVE: 1.0 m/s NO AR.

TRICUSPID VALVE: E - 0.4 m/s A - 0.3 m/s NO TR.

PULMONARY VALVE: 0.8 m/s NO PR.

	Name	SIPRA	ID	MED120924626
	Age & Gender	42/FeMale		26-03-2022 00:00:00
- 1	Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHY FINDINGS:

Left Ventricle : Normal size, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal.Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- NORMAL SIZED CARDIAC CHAMBERS.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

Name	SIPRA	ID	MED120924626
Age & Gender	42/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



DR. ANAND KUMAR M MD DM CONSULTANT INTERVENTIONAL CARDIOLOGIST

Name	SIPRA	ID	MED120924626
Age & Gender	42/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

• No significant abnormality detected.

N	Name	SIPRA	ID	MED120924626
A	Age & Gender	42/FeMale	Visit Date	26-03-2022 00:00:00
- I	Ref Doctor Iame	MediWheel		



TEST COMPLETED