PID No.
 : MED121189269
 Register On
 : 09/07/2022 8:52 AM

 SID No.
 : 522220886
 Collection On
 : 09/07/2022 10:41 AM

 Age / Sex
 : 52 Year(s) / Female
 Report On
 : 09/07/2022 2:32 PM

(*) MEDALI

Type : OP

Ref. Dr : MediWheel

Printed On : 15/07/2022 11:29 AM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	12.44	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	39.4	%	37 - 47
RBC Count (EDTA Blood)	4.98	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood)	79.1	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	25.0	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	31.6	g/dL	32 - 36
RDW-CV	14.6	%	11.5 - 16.0
RDW-SD	40.42	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	5680	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	57.84	%	40 - 75
Lymphocytes (Blood)	31.98	%	20 - 45
Eosinophils (Blood)	5.03	%	01 - 06



VERIFIED BY



 PID No.
 : MED121189269
 Register On
 : 09/07/2022 8:52 AM

 SID No.
 : 522220886
 Collection On
 : 09/07/2022 10:41 AM

 Age / Sex
 : 52 Year(s) / Female
 Report On
 : 09/07/2022 2:32 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood)	5.01	%	01 - 10
Basophils (Blood)	0.14	%	00 - 02
INTERPRETATION: Tests done on Automated Five Pa	art cell counter. All a	abnormal results are	reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	3.29	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	1.82	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.29	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.28	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.01	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	184.4	10^3 / μl	150 - 450
MPV (Blood)	8.62	fL	8.0 - 13.3
PCT (Automated Blood cell Counter)	0.16	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	28	mm/hr	< 30

: 15/07/2022 11:29 AM



VERIFIED BY



: MediWheel

 PID No.
 : MED121189269
 Register On
 : 09/07/2022 8:52 AM

 SID No.
 : 522220886
 Collection On
 : 09/07/2022 10:41 AM

 Age / Sex
 : 52 Year(s) / Female
 Report On
 : 09/07/2022 2:32 PM

Printed On



Type : OP

Ref. Dr

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.71	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.26	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.45	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	22.49	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	23.54	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	29.33	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	63.8	U/L	53 - 141
Total Protein (Serum/Biuret)	7.72	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.47	gm/dl	3.5 - 5.2
Globulin	3.25	gm/dL	2.3 - 3.6

1.38

: 15/07/2022 11:29 AM



(Serum/Derived)
A:GRATIO

(Serum/Derived)

VERIFIED BY



1.1 - 2.2

PID No. : 09/07/2022 8:52 AM : MED121189269 Register On : 522220886 Collection On : 09/07/2022 10:41 AM SID No. Age / Sex : 52 Year(s) / Female Report On : 09/07/2022 2:32 PM

Printed On



Type : OP

Ref. Dr

: MediWheel

: 15/07/2022 11:29 AM

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	227.99	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	234.26	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_ circulating level of triglycerides during most part of the day.

Part of the and			
HDL Cholesterol (Serum/Immunoinhibition)	35.64	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	145.5	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	46.9	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	192.4	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



 PID No.
 : MED121189269
 Register On
 : 09/07/2022 8:52 AM

 SID No.
 : 522220886
 Collection On
 : 09/07/2022 10:41 AM

MEDALL

Type : OP

P Printed On : 15/07/2022 11:29 AM

Report On

Ref. Dr : MediWheel

Age / Sex : 52 Year(s) / Female

InvestigationObservedUnitBiologicalValueReference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

09/07/2022 2:32 PM

Total Cholesterol/HDL Cholesterol Ratio

(Serum/Calculated)

6.4 Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio

(TG/HDL)

(Serum/Calculated)

LDL/HDL Cholesterol Ratio 4.1

(Serum/Calculated)

6.6 Optimal: < 2.5

Mild to moderate risk: 2.5 - 5.0

High Risk: > 5.0

Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0



VERIFIED BY



Printed On

Type : OP

PID No.

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i>)	7.6	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 15/07/2022 11:29 AM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 171.42 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



VERIFIED BY



Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>L</u>	<u>Jnit</u> <u>Biological</u>
	Value	Reference Interval

: 15/07/2022 11:29 AM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.15 ng/ml 0.4 - 1.81

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 9.62 µg/dl 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 7.81 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



 PID No.
 : MED121189269
 Register On
 : 09/07/2022 8:52 AM

 SID No.
 : 522220886
 Collection On
 : 09/07/2022 10:41 AM

 Age / Sex
 : 52 Year(s) / Female
 Report On
 : 09/07/2022 2:32 PM

Printed On



Type : OP

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>Un</u>	<u>Biological</u>
	Value	Reference Interval

: 15/07/2022 11:29 AM

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION (URINE COMPLETE)

Colour	Yellow	Yellow to Amber

(Urine)

Appearance Clear Clear

(Urine)

Volume(CLU) 20

(Urine)

CHEMICAL EXAMINATION (URINE

COMPLETE)

рН	5.5	4.5 - 8.0
----	-----	-----------

(Urine)

Specific Gravity 1.019 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)

Nitrite Negative Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)



VERIFIED BY



PID No. : MED121189269 Register On : 09/07/2022 8:52 AM : 522220886 SID No. Collection On : 09/07/2022 10:41 AM Age / Sex : 52 Year(s) / Female Report On : 09/07/2022 2:32 PM



Type : OP

Printed On : 15/07/2022 11:29 AM

Ref. Dr : MediWheel

Investigation **Observed** <u>Unit</u> **Biological** Reference Interval <u>Value</u>

Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

MICROSCOPIC EXAMINATION (URINE COMPLETE)

NIL 0-1/hpf Pus Cells

(Urine)

Epithelial Cells 0-1/hpf **NIL**

(Urine)

NIL /hpf **NIL RBCs**

(Urine)

NIL Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.



VERIFIED BY



PID No. : MED121189269 : 09/07/2022 8:52 AM Register On

: 522220886 SID No. Collection On : 09/07/2022 10:41 AM

Age / Sex : 52 Year(s) / Female Report On : 09/07/2022 2:32 PM **Type** : OP

Ref. Dr : MediWheel

: 15/07/2022 11:29 AM **Printed On**



Investigation <u>Unit</u> <u>Observed</u> **Biological Value** Reference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'A' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



VERIFIED BY



: MED121189269 PID No. Register On : 09/07/2022 8:52 AM

: 522220886 SID No. Collection On : 09/07/2022 10:41 AM

Age / Sex : 52 Year(s) / Female Report On 09/07/2022 2:32 PM Type : OP

Ref. Dr : MediWheel **Printed On** : 15/07/2022 11:29 AM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	14.3		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	148.46	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Negative Glucose, Fasting (Urine) Negative (Urine - F/GOD - POD)

Glucose Postprandial (PPBS) 196.47 mg/dL 70 - 140

(Plasma - PP/GOD-PAP)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

8.9 7.0 - 21Blood Urea Nitrogen (BUN) mg/dL (Serum/*Urease UV* / *derived*) 0.6 - 1.1 0.62 Creatinine mg/dL

(Serum/Modified Jaffe)

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine

Uric Acid 6.06 2.6 - 6.0mg/dL

(Serum/Enzymatic)



VERIFIED BY



APPROVED BY

-- End of Report --



Name	MRS.CHANDRAKALA D C	ID	MED121189269
Age & Gender	52/FEMALE	Visit Date	09/07/2022
Ref Doctor Name	MediWheel		

X-ray mammogram (mediolateral oblique & craniocaudal views) followed by Sonomammography.

BILATERAL MAMMOGRAPHY

Breast composition Type B (there are scattered areas of fibroglandular density)

No evidence of focal soft tissue lesion.

No evidence of cluster microcalcification.

Subcutaneous fat deposition is within normal limits.

Axillary lymph nodes are noted in left axilla.

BILATERAL SONOMAMMOGRAPHY

A well defined hypoechoic oval cystic lesion is noted at 8 o' clock position measuring 12.0 x 4.0 mm in subcutaneous plane in right breast - folliculitis.

Both the breasts show normal echopattern.

No evidence of focal solid areas.

No evidence of ductal dilatation.

Bilateral axillary lymphnodes are noted with preserved fatty hilum.

IMPRESSION:

- Well defined hypoechoic oval cystic lesion in subcutaneous plane in right breast folliculitis.
- No breast lesions.
- Bilateral axillary lymph nodes.

ASSESSMENT: BI-RADS CATEGORY - 2

BI-RADS CLASSIFICATION

CATEGORY RESULT

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- $11. Disputes, if any\ , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.\\$



Name	MRS.CHANDRAKALA D C	ID	MED121189269
Age & Gender	52/FEMALE	Visit Date	09/07/2022
Ref Doctor Name	MediWheel		

Benign finding. Routine mammogram in 1 year recommended.

DR. HEMANANDINI V.N
CONSULTANT RADIOLOGIST
Hn/mj

2

REPORT DISCLAIMER

^{1.}This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.

^{2.} The results reported here in are subject to interpretation by qualified medical professionals only.

^{3.} Customer identities are accepted provided by the customer or their representative.

^{4.}information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.

^{5.}If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.

^{6.}Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.

^{7.}Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,

^{8.}If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.

^{9.}Liability is limited to the extend of amount billed.

^{10.}Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.

^{11.}Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.