Name	SUNITHA P	ID	OPCS15-101525
Age & Gender	34Year(s)/FEMALE	Visit Date	3/26/2022 12:00:00 AM
Ref Doctor Name	MediWheel		

ECHO CARDIOGRAPHY REPORT

Measurements:-

M Mode:

IVS d	1.1 cm	IVS s	1.2 cm
LVID d	5.4 cm	LVID s	3.4 cm
LVPW d	1.3 cm	LVPW s	1.1 cm
AO	3.1 cm	LA	2.9 cm

Doppler study:

Location	m/sec	Location	m/sec
MP A vel	0.9	MV E	0.8
PGT	3 mmHg	Α	0.4
AV vel	1.5	Ratio	1.72
PGT	9 mmHg	TV VEL	1.9
EF	65 %	PGT	14 mmHg
FS	36 %		

<u>2D:</u>

LA	:	NORMAL	F	RA :	NORMAL
LV	:	NORMAL	RV	:	NORMAL
AV	:	NORMAL	PV :	NOR	MAL
MV	:	NORMAL	TV :	NOR	MAL
AO	:	NORMAL	PA	:	NORMAL

Name	SUNITHA P	ID	OPCS15-101525
Age & Gender	34Year(s)/FEMALE	Visit Date	3/26/2022 12:00:00 AM
Ref Doctor Name	MediWheel		

Observations:

- Cardiac chambers dimension-normal
- No regional wall motion abnormality
- Normal LV systolic and diastolic function
- Doppler flow pattern normal
- Trivial AR / MR
- No pulmonary hypertension
- Normal Pericardium
- IAS/ IVS appear Intact
- No mass

CONCLUSIONS:

- NORMAL CARDIAC DIMENSIONS.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- GOOD LV SYSTOLIC FUNCTION.
- LVEF 65 %
- NORMAL STUDY.

An

Name	SUNITHA P	ID	OPCS15-101525
Age & Gender	34Year(s)/FEMALE	Visit Date	3/26/2022 12:00:00 AM
Ref Doctor Name	MediWheel		

Prof. N. Subramanian MD, DM(CARD) FRCP, FACC Consultant Cardiologist

Done By :-M.Padmapushani Cardiac technologist

Name	: Ms. SUNITHA P					
PID No.	: OPCS15-101525	Register On	: 26/03/2022 10:27 AM	\sim		
SID No.	: 1802211766	Collection On	: 26/03/2022 10:47 AM			
Age / Sex	: 34 Year(s) / Female	Report On	26/03/2022 7:24 PM	MEDALL		
Туре	: OP	Printed On	31/03/2022 3:04 PM			
Ref. Dr	: MediWheel					
Investiga	Investigation		tigation <u>Observed Un</u> <u>Value</u>		Unit	Biological Reference Interval
TYPING (EDTA BI INTERPI	ood/Agglutination) RETATION: Reconfirm the Blood g	'B' 'Positiv roup and Typing be				
<u>Complet</u>	e Blood Count With - ESR					
Haemog (EDTA Bl	lobin ood/Spectrophotometry)	11.8	g/dL	12.5 - 16.0		
	Cell Volume(PCV)/Haematocrit ood/Derived from Impedance)	35.8	%	37 - 47		
RBC Co (EDTA Bl	unt ood/Impedance Variation)	4.60	mill/cu.mm	4.2 - 5.4		
	orpuscular Volume(MCV) ood/Derived from Impedance)	77.9	fL	78 - 100		
	orpuscular Haemoglobin(MCH) ood/Derived from Impedance)	25.7	pg	27 - 32		
concentr	orpuscular Haemoglobin ation(MCHC) ood/Derived from Impedance)	33.1	g/dL	32 - 36		
RDW-C (EDTA Bl	V ood/Derived from Impedance)	13.8	%	11.5 - 16.0		
RDW-SI (EDTA Bl) ood/Derived from Impedance)	37.63	fL	39 - 46		
	ukocyte Count (TC) ood/Impedance Variation)	5390	cells/cu.mm	4000 - 11000		
Neutroph (EDTA Ble Cytometry)	ood/Impedance Variation & Flow	58.2	%	40 - 75		
Lympho (EDTA Ble Cytometry)	ood/Impedance Variation & Flow	30.9	%	20 - 45		
Eosinopl (EDTA Bl Cytometry)	ood/Impedance Variation & Flow	4.8	%	01 - 06		
Monocyt (EDTA Ble Cytometry)	ood/Impedance Variation & Flow	5.7	%	01 - 10		



APPROVED BY

The results pertain to sample tested.

Page 1 of 7

Name	: Ms. SUNITHA P		
PID No.	: OPCS15-101525	Register On : 26/03/2022 10:27 AM	\mathbf{M}
SID No.	: 1802211766	Collection On : 26/03/2022 10:47 AM	
Age / Sex	: 34 Year(s) / Female	Report On : 26/03/2022 7:24 PM	MEDALL
Туре	: OP	Printed On : 31/03/2022 3:04 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.4	%	00 - 02
INTERPRETATION: Tests done on Automated	Five Part cell count	er. All abnormal results a	are reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.14	10^3 / µl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.67	10^3 / µl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.26	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.31	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.02	10^3 / µl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	307	10^3 / µl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	8.8	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.27	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	6	mm/hr	< 20
BUN / Creatinine Ratio	18.8		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	85.4	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	115.0	mg/dL	70 - 140



APPROVED BY

The results pertain to sample tested.

Page 2 of 7

Name	: Ms. SUNITHA P			
PID No.	: OPCS15-101525	Register On : 2	6/03/2022 10:27 AM	m
SID No.	: 1802211766	Collection On : 2	6/03/2022 10:47 AM	
Age / Sex	: 34 Year(s) / Female	Report On : 2	26/03/2022 7:24 PM	MEDALL
Туре	: OP	Printed On : 3	1/03/2022 3:04 PM	
Ref. Dr	: MediWheel			
Investiga	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Factors su Fasting ble	ood glucose level may be higher that	n Postprandial glucose,	because of physiological	nd drugs can influence blood glucose level. surge in Postprandial Insulin secretion, Insulin cation during treatment for Diabetes.
Urine Gl (Urine - PF	ucose(PP-2 hours)	Negative		Negative
	rea Nitrogen (BUN) ease UV/derived)	14.3	mg/dL	7.0 - 21
Creatinir (Serum/Ma	ne odified Jaffe)	0.76	mg/dL	0.6 - 1.1
ingestion of	of cooked meat, consuming Protein/	Creatine supplements, I	Diabetic Ketoacidosis, pro	evere dehydration, Pre-eclampsia, increased blonged fasting, renal dysfunction and drugs he, chemotherapeutic agent such as flucytosine
Uric Aci (Serum/En		4.0	mg/dL	2.6 - 6.0
	unction Test			
Bilirubin (Serum/DC	n(Total) CA with ATCS)	0.59	mg/dL	0.1 - 1.2
Bilirubin (Serum/Dia	n(Direct) azotized Sulfanilic Acid)	0.12	mg/dL	0.0 - 0.3
Bilirubin (Serum/De	n(Indirect) erived)	0.47	mg/dL	0.1 - 1.0
Aminotra	ST (Aspartate ansferase) odified IFCC)	30.8	U/L	5 - 40
	LT (Alanine Aminotransferase)) 38.2	U/L	5 - 41
	umma Glutamyl Transpeptidase CC / Kinetic)) 30.0	U/L	< 38
	Phosphatase (SAP) <i>podified IFCC</i>)	121.6	U/L	42 - 98
Total Pro (Serum/Bin		6.65	gm/dl	6.0 - 8.0
Albumin (Serum/Bro	omocresol green)	3.89	gm/dl	3.5 - 5.2



APPROVED BY

The results pertain to sample tested.

Page 3 of 7

Name	: Ms. SUNITHA P		
PID No.	: OPCS15-101525	Register On : 26/03/2022 10:27 AM	\mathbf{C}
SID No.	: 1802211766	Collection On : 26/03/2022 10:47 AM	
Age / Sex	: 34 Year(s) / Female	Report On : 26/03/2022 7:24 PM	MEDALL
Туре	: OP	Printed On : 31/03/2022 3:04 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Globulin (Serum/Derived)	2.76	gm/dL	2.3 - 3.6
A : G RATIO (Serum/ <i>Derived</i>) <i>Lipid Profile</i>	1.41		1.1 - 2.2
Cholesterol Total (Serum/CHOD-PAP with ATCS)	140.6	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	91.3	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	41.7	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/ <i>Calculated</i>)	80.6	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	18.3	mg/dL	< 30
Non HDL Cholesterol (Serum/ <i>Calculated</i>)	98.9	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

h Dayanand Kinha Pathologis No: 142072 Rea

APPROVED BY

The results pertain to sample tested.

Page 4 of 7

Name	: Ms. SUNITHA P		
PID No.	: OPCS15-101525	Register On : 26/03/202	2 10:27 AM 💦 🐴 👔
SID No.	: 1802211766	Collection On : 26/03/202	
Age / Sex	: 34 Year(s) / Female	Report On : 26/03/202	22 7:24 PM MEDALL
Туре	: OP	Printed On : 31/03/202	22 3:04 PM
Ref. Dr	: MediWheel	- -	
Investiga	ation	<u>Observed</u> <u>Unit</u> <u>Value</u>	Biological Reference Interval
2.It is the		roteins including LDL, IDL, VLD	rdiovascular risk marker than LDL Cholesterol. L and chylomicrons and it is the "new bad cholesterol" and is a
Total Ch Ratio (Serum/Ca	olesterol/HDL Cholesterol	3.4	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglycer (TG/HD) (Serum/Ca	<i>'</i>	2.2	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HD (Serum/Ca	DL Cholesterol Ratio	1.9	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

Glycosylated Haemoglobin (HbA1c)

HbA1C	5.7	%	Normal: 4.5 - 5.6
(Whole Blood/HPLC)			Prediabetes: 5.7 - 6.4
			Diabetic: ≥ 6.5

INTERPRETATION: If Diabetes - Good control : 6.1 - 7.0 %, Fair control : 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

mg/dL

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

116.89

hypertriglyceridemia,hyperbilirubinemia,Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values. Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly,Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

<u>THYROID PROFILE / TFT</u>

 T3 (Triiodothyronine) - Total
 2.05
 ng/ml
 0.7 - 2.04

 (Serum/Chemiluminescent Immunometric Assay (CLIA))
 0.7
 0.7

INTERPRETATION:

Comment :

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.



APPROVED BY

The results pertain to sample tested.

Page 5 of 7

Name	: Ms. SUNITHA P			
PID No.	: OPCS15-101525	Register On	: 26/03/2022 10:27 AM	m
SID No.	: 1802211766	Collection On	: 26/03/2022 10:47 AM	
Age / Sex	: 34 Year(s) / Female	Report On	: 26/03/2022 7:24 PM	MEDALL
Туре	: OP	Printed On	: 31/03/2022 3:04 PM	
Ref. Dr	: MediWheel			
<u>Investiga</u>	ation	<u>Observed</u> <u>Value</u>	<u>I Unit</u>	Biological Reference Interval
T4 (Tyro	oxine) - Total	9.52	μg/dl	4.2 - 12.0
(Serum/Ch (CLIA))	emiluminescent Immunometric Assay			
INTERPI	RETATION:			

Comment :

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) (Serum/Chemiluminescent Immunometric Assay (CLIA))	1.60	µIU/mL	0.35 - 5.50
INTERPRETATION: Reference range for cord blood - upto 20 1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0 (Indian Thyroid Society Guidelines) Comment :			

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI. 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM.The variation can be of the order of 50%,hence time of the day has influence on the measured serum TSH concentrations.

3. Values&lt,0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

Urine Analysis - Routine

COLOUR (Urine)	Pale yellow		Yellow to Amber
APPEARANCE (Urine)	Clear		Clear
Protein (Urine/Protein error of indicator)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Pus Cells (Urine/Automated – Flow cytometry)	1 - 2	/hpf	NIL
Epithelial Cells (Urine/Automated – Flow cytometry)	2 - 4	/hpf	NIL
RBCs (Urine/Automated – Flow cytometry)	NIL	/hpf	NIL
Casts (Urine/Automated – Flow cytometry)	NIL	/hpf	NIL



APPROVED BY

The results pertain to sample tested.

Page 6 of 7

<u>Investiga</u>	ation	<u>Observed</u> <u>Unit</u> <u>Value</u>	Biological Reference Interval
Ref. Dr	: MediWheel		
Туре	: OP	Printed On : 31/03/2022 3:04 PM	
Age / Sex	: 34 Year(s) / Female	Report On : 26/03/2022 7:24 PM	MEDALL
SID No.	: 1802211766	Collection On : 26/03/2022 10:47 AM	
PID No.	: OPCS15-101525	Register On : 26/03/2022 10:27 AM	M
Name	: Ms. SUNITHA P		

NIL

NIL

(Urine/Automated – Flow cytometry) Others

(Urine)

Crystals

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

/hpf

Dayanand Kinha Patholo ... 42072

NIL

APPROVED BY

-- End of Report --

The results pertain to sample tested.

Page 7 of 7

Name	SUNITHA P	ID	OPCS15-101525
Age & Gender	34Year(s)/FEMALE	Visit Date	3/26/2022 12:00:00 AM
Ref Doctor Name	MediWheel		

SONOGRAM REPORT - WHOLE ABDOMEN

Indication: General check up

The liver is normal in size (11.2 cms) and shows uniform echotexture with no focal abnormality.

The gall bladder is normal sized. **Sludge noted in the lumen of gall bladder.**

There is no intra or extra hepatic biliary ductal dilatation.

The pancreas shows a normal configuration and echotexture.

The pancreatic duct is normal.

The portal vein and IVC are normal.

The spleen is normal in size and measures 9.1 cms.

There is no free or loculated peritoneal fluid.

No para aortic lymphadenopathy is seen.

No abnormality is seen in the region of the adrenal glands.

The right kidney measures 8.9 x 4.2 cms.

The left kidney measures 9.6 x 4.2 cms.

Both kidneys are normal in size, shape and position. Cortical echoes are normal bilaterally.

There is no calculus or calyceal dilatation.

The ureters are not dilated.

Name	SUNITHA P	ID	OPCS15-101525
Age & Gender	34Year(s)/FEMALE	Visit Date	3/26/2022 12:00:00 AM
Ref Doctor Name	MediWheel		

The urinary bladder is smooth walled and uniformly transonic. There is no intravesical mass or calculus.

The uterus is anteverted, and measures 8.0 x 4.5 x 4.5 cms.

Myometrial echoes are homogeneous.

The endometrial thickness is 6.9 mm.

The right ovary measures 3.6 x 2.3 cms.

The left ovary measures 2.9 x 1.7 cms.

No significant mass or cyst is seen in the ovaries.

Parametria are free.

Iliac fossae are normal.

No mass or fluid collection is seen in the right iliac fossa. The appendix is not visualized.

IMPRESSION:

Sludge in gall bladder.

--- Suggested Clinical Correlation

Hara

DR. T. SANA

CONSULTANT SONOLOGIST

SH

Name	SUNITHA P	ID	OPCS15-101525
Age & Gender	34Year(s)/FEMALE	Visit Date	3/26/2022 12:00:00 AM
Ref Doctor Name	MediWheel		

Name	SUNITHA P	Customer ID	OPCS15-101525
Age & Gender	34Y/F	Visit Date	Mar 26 2022 8:51AM
Ref Doctor	MediWheel		

X-RAY CHEST (PA VIEW)

The heart size and configuration are within normal limits. The aortic arch is normal.

The lung fields show normal broncho-vascular markings.

Both the pulmonary hila are normal in size.

The costophrenic and cardiophrenic recesses and the domes of diaphragm are normal.

The bones and soft tissues of the chest wall show no abnormality.

IMPRESSION:

• No significant abnormality demonstrated.

Reben Schene

DR.REKHA S.CHERIAN, DMRD.DNB.FRCR., CONSULTANT RADIOLOGIST