

CERTIFICATE OF MEDICAL FITNESS

NAME: Ms. Spoorthi G.

AGE/ GENDER: 23y / F

HEIGHT: 154cm

WEIGHT: 59.5kg

IDENTIFICATION MARK: -

BLOOD PRESSURE: 110/70 mm/Hg.

PULSE: 76 / min

CVS: } Normal
RS:P }

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Nil

ALLERGIES, IF ANY: Nil

LIST OF PRESCRIBED MEDICINES: Nil

ANY OTHER REMARKS: Nil

I Certify that I have carefully examined Mr/Mrs. Spoorthi G. son/daughter of Mr. Gowindaraju. who has signed in my presence. He/ she has no physical disease and is fit for employment.

Spoorthi

Signature of candidate

Dr. BINDURAJ. R
MBBS, MD
Internal Medicine
Reg. No. 42805

Signature of Medical Officer

Place: Spectrum diagnostic & health care

Date: 23/09/23

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined



Dr. Ashok S
Bsc., MBBS., D.O.M.S
Consultant Ophthalmologist
KMC No: 31827

DATE: 23/09/23

EYE EXAMINATION

NAME: SP00RTHI.g

AGE: 23

GENDER: F/M

	RIGHT EYE	LEFT EYE
Vision	6/6	6/6
Vision With glass	6/6	6/6
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ impression	Normal	Normal

Dr. ASHOK SARODHE
B.Sc., M.B.B.S., D.O.M.S.
Eye Consultant & Surgeon
KMC No: 31827
Consultant (Ophthalmologist)

SCAN FOR LOCATION



NAME	AGE	GENDER
Spoorthi G.	23 yrs	Female

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY → DDC in 4el to 7; DC in 4el to 8
Both need perforators.

M: MISSING

O: OTHERS

ADVISED: ✓ ✓

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:


23/09/23

SIGNATURE OF THE DENTAL SURGEON

SEAL

Dr. SACHDEV NAGARKAR
B.D.S., F.A.G.E., F.P.F.A. (USA)
Reg. No : 2247/A

DATE

SCAN FOR LOCATION



SPECTRUM DIAGNOSTICS & HEALTH CARE

#9/1 TEJAS ARCADE, DR. RAJKUMAR ROAD, RAJAJINAGAR-560010 AUDIOGRAM



Patient ID : 0868

Name : SPOORTHI G


CR Number : 20230923101631

Registration Date : 23-Sep-2023

Age : 23

Gender : Female

Operator : spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 Hz	1500 Hz	2000 Hz	3000 Hz	4000 Hz	6000 Hz	8000 Hz
X - Air Left	25	30	40	25	30	25	15	10	10	15	10
O - Air Right	40	35	35	35	30	30	20	20	20	20	15
> - Bone Left											
< - Bone Right											

Clinical Notes :

Right Ear : Normal
Left Ear : Normal




MS SPOORTHI G
Female 23Years

Diagnosis Information:
Sinus Arrhythmia

HR	: 70	bpm
P	: 93	ms
PR	: 144	ms
QRS	: 67	ms
QT/QTc	: 370/400	ms
P/QRS/T	: 52/59/-6	°
RV5/SV1	: 0.682/0.642	mV

Report Confirmed by:



NAME : MRS.SPOORTHI G	DATE :23/09/2023
AGE/SEX : 23 YEARS/FEMALE	REG NO:2309230026
REF BY : APOLO CLINIC	

CHEST PA VIEW


Lung fields are clear.

Cardiovascular shadows are within normal limits.

Both CP angles are free.

Domes of diaphragm and bony thoracic cage are normal.

IMPRESSION: NORMAL CHEST RADIOGRAPH.



**DR.RAM PRAKASH G MDRD
CONSULTANT RADIOLOGIST**

Your suggestion / feedback is a valuable input for improving our services

SCAN FOR LOCATION



NAME AND LAB NO	MRS SPOORTHI G	Reg : 30026
AGE & SEX	23 YRS	FEMALE
DATE AND AREA OF INTEREST	23.09.2023	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

LIVER: Measures 12.5 cm. Normal in size with echotexture. No e/o IHBR dilatation. No evidence of SOL. Portal vein appears normal. CBD appears normal. . No e/o calculus / SOL

GALL BLADDER: Well distended. Wall appears normal.No e/o calculus/ neoplasm. **Sludge noted.**

SPLEEN: Measures 9.3 cm. Normal in size and echotexture. No e/o SOL/ calcification.

PANCREAS: Normal in size and echotexture. Pancreatic duct appears normal. No e/o calculus / calcifications.

RETROPERITONEUM: Poor window.

RIGHT KIDNEY: Measures 9.0 X 3.6 cm. Right kidney is normal in size & echotexture No evidence of calculus/ hydronephrosis.

LEFT KIDNEY: Measures 9.0 X4.3 cm .Left kidney is normal in size & echotexture No evidence of calculus/ hydronephrosis.

URETERS: Bilateral ureters are not dilated.

URINARY BLADDER: Partially distended. No wall thickening/ calculi.

UTERUS: Anteverted,Normal in size and echotexture Endometrium is normal.ET -9mm.

OVARIES: Right ovary - normal in size and echotexture. Left ovary – obscured by bowel gas shadows.

- No evidence of ascites/pleural effusion.

IMPRESSION:

- No significant sonological abnormality detected in the abdomen and pelvis.


DR AKSHATHA R BHAT
MDRD DNB FRCR

SCAN FOR LOCATION



PATIENT NAME	MRS SPOORTHY G	ID NO	2309230026
AGE	23YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	23.09.2023

2D ECHO CARDIOGRAHIC STUDY

M-MODE

AORTA	25mm
LEFT ATRIUM	25mm
RIGHT VENTRICLE	18mm
LEFT VENTRICLE (DIASTOLE)	37mm
LEFT VENTRICLE(SYSTOLE)	28mm
VENTRICULAR SEPTUM (DIASTOLE)	09mm
VENTRICULAR SEPTUM (SYSTOLE)	08mm
POSTERIOR WALL (DIASTOLE)	09mm
POSTERIOR WALL (SYSTOLE)	10mm
FRACTIONAL SHORTENING	30%
EJECTION FRACTION	60%

DOPPLER /COLOUR FLOW

MITRAL VALVE	E-0.65 m/sec	A-0.32m/sec	NO MR
AORTIC VALVE	1.12 m/sec		NO AR
PULMONARY VALVE	1.20 m/sec		NO PR
TRISCUSPID VALVE		26mmHg	MILD TR



PATIENT NAME	MRS SPOORTHI G	ID NO	2309230026
AGE	23YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	23.09.2023

2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE : NORMAL
LEFT ATRIUM : NORMAL
RIGHT ATRIUM : NORMAL
MITRAL VALVE : NORMAL
AORTIC VALVE : NORMAL
PULMONARY VALVE: NORMAL
TRICUSPID VALVE : NORMAL
INTER ATRIAL SEPTUM :INTACT
INTER VENTRICULAR SEPTUM: INTACT
PERICARDIUM : NORMAL
OTHERS : - NIL

IMPRESSION

- NORMAL CARDIAC CHAMBER DIMENSIONS
- NO RWMA OF LV AT REST
- NORMAL CARDIAC VALVES
- NORMAL LV FUNCTION, LVEF-60%
- MILD TR/ MILD PAH [RVSP-26mmHg]
- NO CLOT / PERICARDIAL EFFUSION
- NO ASD / VSD / PDA / CoA SEEN


V. DURGA
ECHO TECHNICIAN

The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.

SCAN FOR LOCATION



Name: MRS. SPOORTHY G	Age/Sex: 23Y Female	Date of receipt: 23.09.2023 Date of report: 23.09.2023
Ref DR: Dr. APOLO CLINIC	LABREFNO: 2309230026	PAP No: 306 /23

CERVICAL PAP SMEAR REPORT

Clinical history : Health check

Specimen : 2 Conventional PAP smears.

Specimen Adequacy : Adequate for evaluation.

Description : Seen are mixture of intermediate squamous cells, superficial squamous cells and endocervical cell clusters.

Inflammation : Neutrophilic exudate is noted.

Organism : Dodderlein bacilli are seen.

Reactive changes : Nil

Dysplastic changes : Nil

Impression : **Negative for Squamous Intraepithelial Lesion/Malignancy.**


Note: Enclosed: 2slides: preserve them carefully.

---End of report---



Dr. Nithun Reddy C, MD
Consultant Pathologist



Name : MRS. SPOORTHI G	UHID : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026	2309230026	Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Blood EDTA				
Haemoglobin (HB)	14.60	g/dL	Female: 12.0 - 15.0	Spectrophotometer
Red Blood Cell (RBC)	4.75	million/cumm	3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	42.10	%	Female: 36.0 - 45.0	Electronic Pulse
Mean corpuscular volume (MCV)	88.80	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	30.80	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	34.80	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	38.60	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	13.70	%	Female: 12.20 - 16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	8.60	fL	8.0-15.0	Volumetric Impedance
Platelet	2.77	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	10.60	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	9650.00	cells/cumm	Female: 4000.0 - 11000.0	Volumetric Impedance
Neutrophils	55.70	%	40.0-75.0	Light scattering/Manual
Lymphocytes	36.90	%	20.0-40.0	Light scattering/Manual
Eosinophils	4.80	%	0.0-8.0	Light scattering/Manual
Monocytes	2.50	%	0.0-10.0	Light scattering/Manual
Basophils	0.10	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	5.37	10 ³ /uL	2.0- 7.0	Calculated



Name : MRS. SPOORTHI G	UHID : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC	2309230026	Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Absolute Lymphocyte Count	3.00	10 ³ /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.24	10 ³ /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	470.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.01	10 ³ /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	27	mm/hr	Female: 0.0 - 20.0	Westergren

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.
WBC'S : Are normal in total number, morphology and distribution.
Platelets : Adequate in number and normal in morphology.
No abnormal cells or hemoparasites are present.
Impression : Normocytic Normochromic Blood picture.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Name : MRS. SPOORTHI G	UHD : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC	2309230026	Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA	4.90	%	Non diabetic adults : <5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes : >= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control : 8-10 Poor Control : >10	HPLC
Glycosylated Haemoglobin (HbA1c)				
Estimated Average Glucose(eAG)	93.93	mg/dL		Calculated


Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemc control as compared to blood and urinary glucose determinations.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C.MD.Consultant Pathologist



Name : MRS. SPOORTHI G	UHID : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female	 2309230026	Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)
Fasting Blood Sugar (FBS)- Plasma	85	mg/dL	60.0-110.0	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist


SCAN FOR LOCATION



Page 4 of 12

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010
+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985

Name : MRS. SPOORTHI G	UHID : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC	2309230026	Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test) :				
Blood Urea Nitrogen (BUN)-Serum	14.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.77	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	4.10	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	140.4	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.54	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	101.60	mmol/L	94.0-110.0	Ion-Selective Electrodes (ISE)



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Page 5 of 12

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010

+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985

Name	: MRS. SPOORTHI G	UHID	: 2309230026	Bill Date	: 23-Sep-2023 08:31 AM
Age / Gender	: 23 Years / Female			Sample Col. Date	: 23-Sep-2023 08:31 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 23-Sep-2023 03:25 PM
Reg. No.	: 2309230026			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	200.00	mg/dL	Female: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	67.00	mg/dL	Female: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	40.00	mg/dL	Female: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	160	mg/dL	Female: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	116.00	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	13	mg/dL	Female: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	5.00	Ratio	Female: 0.0 - 5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Name : MRS. SPOORTHI G	UHID : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC	2309230026	Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)- Serum				
Tri-Iodo Thyronine (T3)-Serum	1.13	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	10.00	µg/dL	Female: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	1.59	µIU/mL	Female: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester : 0.81-1.90, Second Trimester : 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).

Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5, Females : 5.5-11.0, > 60 Years: 5.0-10.70, Cord : 7.40-13.10, Children: 1-3 Days : 11.80-22.60, 1-2 Weeks : 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years : 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days : >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy.

Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0


Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.

Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency




Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Name : MRS. SPOORTHI G	UHD : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC	2309230026	Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serum				
Bilirubin Total-Serum	0.76	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.18	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.58	mg/dL	Female: 0.0 - 1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	30.00	U/L	Female: 15.0 - 37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	26.00	U/L	Female: 14.0 - 59.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	71.00	U/L	Female: 45.0 - 117.0	PNPP,AMP-Buffer
Protein, Total-Serum	7.89	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	4.75	g/dL	Female: 3.40 - 5.50	Bromocresol Purple
Globulin-Serum	3.14	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.51	Ratio	0.80-1.20	Calculated



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Name : MRS. SPOORTHI G	UHID : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female	 2309230026	Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Calcium, Total- Serum	9.30	mg/dL	8.50-10.10	Spectrophotometry (O- Cresolphthalein complexone)
Gamma-Glutamyl Transferase (GGT)-Serum	18.00	U/L	Female: 5.0 - 55.0	Other g-Glut-3- carboxy-4 nitro

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Page 9 of 12

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010
+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 | +91 6361 253 097 | 080-2991 6944 | 080-49511985

Name : MRS. SPOORTHI G	UHID : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026	2309230026	Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Post Prandial Urine Sugar	Negative		Negative	Dipstick/Benedicts(Mar
Post prandial Blood Glucose (PPBS)-Plasma	90	mg/dL	70-140	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Name : MRS. SPOORTHI G	UHD : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female	 2309230026	Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-Urine				
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	6.00		5.0-7.5	Dipstick
Specific Gravity	1.025		1.000-1.030	Dipstick
Biochemical Examination				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination				
Pus Cells	6-8	hpf	0.0-5.0	Microscopy
Epithelial Cells	2-4	hpf	0.0-10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C,MD, Consultant Pathologist

SCAN FOR LOCATION



Name : MRS. SPOORTHI G	UHD : 2309230026	Bill Date : 23-Sep-2023 08:31 AM
Age / Gender : 23 Years / Female		Sample Col. Date : 23-Sep-2023 08:31 AM
Ref. By Dr. : Dr. APOLO CLINIC	2309230026	Result Date : 23-Sep-2023 03:25 PM
Reg. No. : 2309230026		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Blood Group & Rh Typing-Whole Blood EDTA				
Blood Group	O			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum
Printed On : 23 Sep, 2023 03:36 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Page 12 of 12

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010

+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985