



बैंक ऑफ बड़ौदा
Bank of Baroda

नाम : ललित कुमार
Name : Lalit Kumar
कर्मचारी कूट क. :
E.C. No. : 122553



जारीकर्ता प्राधिकारी
Issuing Authority

Lalit
धारक के हस्ताक्षर
Holder's Signature

Lalit



Dr. U. C. GUPTA
MBBS, MD (Physician)
RMC No. 281

If found please return to:
Asst. General Manager (Security)
Bank of Baroda, Baroda Corporate Centre,
C-26, G-Block, Bandra-Kurla Complex, Mumbai-4000 051
Phone : 91 22 6698 5196 Fax : 91 22 2652 5747

मिलने पर निम्नलिखित को लौटायें :
सहायक महाप्रबंधक (सुरक्षा)
बैंक ऑफ बड़ौदा, बड़ौदा कॉरपोरेट सेन्टर,
सी-26, जी-ब्लॉक, बांद्रा-कुर्ला कॉम्प्लेक्स, मुम्बई-400051
फोन नंबर : 91 22 6698 5196 फैक्स : 91 22 2652 5747
Blood Group B+



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com

General Physical Examination

Date of Examination: 28/01/2023

Name: LALIT KUMAR Age: 30 DOB: 30/08/1992 Sex: MALE

Referred By: BANK OF BARODA

Photo ID: EMPLOYEE ID ID #: 122553

Ht: 186 (cm) Wt: 118 (Kg)

Chest (Expiration): 122 (cm) Abdomen Circumference: 127 (cm)

Blood Pressure: 120/80 mm Hg PR: 78 / min RR: 18 / min Temp: Afebrile

BMI _____

Eye Examination: RIE } 6/6 , N/6 , NCB
LE } 6/6 , N/6 , NCB

Other: NO

On examination he/she appears physically and mentally fit: Yes / No

Signature Of Examinee : Lalit Name of Examinee: MRS. LALIT KUMAR

Signature Medical Examiner : [Signature] Name Medical Examiner Dr. U.C. Gupta

Dr. U. C. GUPTA
MBBS, MD (Physician)
RMC No. 291



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
91-141-4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

FULL BODY HEALTH CHECKUP BELOW 40 MALE

HAEMOGARAM

HAEMOGLOBIN (Hb) 15.2 g/dL 13.0 - 17.0

TOTAL LEUCOCYTE COUNT 7.90 /cumm 4.00 - 10.00

DIFFERENTIAL LEUCOCYTE COUNT

NEUTROPHIL 59.0 % 40.0 - 80.0

LYMPHOCYTE 34.0 % 20.0 - 40.0

EOSINOPHIL 3.0 % 1.0 - 6.0

MONOCYTE 4.0 % 2.0 - 10.0

BASOPHIL 0.0 % 0.0 - 2.0

TOTAL RED BLOOD CELL COUNT (RBC) 4.94 x10⁶/uL 4.50 - 5.50

HEMATOCRIT (HCT) 46.40 % 40.00 - 50.00

MEAN CORP VOLUME (MCV) 94.0 fL 83.0 - 101.0

MEAN CORP HB (MCH) 30.8 pg 27.0 - 32.0

MEAN CORP HB CONC (MCHC) 32.7 g/dL 31.5 - 34.5

PLATELET COUNT 283 x10³/uL 150 - 410

RDW-CV 13.6 % 11.6 - 14.0

MENTZER INDEX 19.03 H 0.00 - 0.00

A complete blood picture (CBP) is a kind of blood test that is done to assess a person's overall health and diagnose a wide range of health disorders like leukemia, anemia and other infections.

A complete blood count (CBC) is a complete blood test that diagnose many components and features of a persons blood which includes: -

- *Red Blood Cells (RBC), which carry oxygen -
- *White Blood Cells (WBC), which help in fighting against infections -
- *Hemoglobin, which is the oxygen carrying protein in the red blood cells -
- *Hematocrit (HCT), the proportion of RBC to the fluid component, or plasma present in blood -
- *Platelets, which aid in blood clotting

(CBC): Methodology: TLC,TRBC,PCV,PLT Impedance method, HB Calorimetric method, and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: MINDRAY BC-3000 Plus 3 part automatic analyzer,

MGR

Technologist

Page No: 1 of 15

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

09

mm in 1st hr

00 - 15

Method:- Westergreen

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases. ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



MGR

Technologist

Page No: 2 of 15

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:30:34

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------


FASTING BLOOD SUGAR (Plasma) Method:- GOD POD	110.0	mg/dl	70.0 - 115.0
--	-------	-------	--------------

Impaired glucose tolerance (IGT)	111 - 125 mg/dL
Diabetes Mellitus (DM)	> 126 mg/dL

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .

BLOOD SUGAR PP (Plasma) Method:- GOD PAP	97.5	mg/dl	70.0 - 140.0
---	------	-------	--------------

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .



MGR

Technologist
Page No: 4 of 15

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- MR. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

GLYCOSYLATED HEMOGLOBIN (HbA1C)

Method:- CAPILLARY with EDTA 5.2 mg%

MEAN PLASMA GLUCOSE 106 mg/dL 0 - 140
Method:- Calculated Parameter

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

- Reference Group HbA1c in %
- Non diabetic adults >=18 years < 5.7
- At risk (Prediabetes) 5.7 - 6.4
- Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings.
Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al]

1. Erythropoiesis

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis.
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease.

2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH

4. Erythrocyte destruction

- Increased HbA1c: increased erythrocyte life span: Splenectomy.
- Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

5. Others

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
- Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Note:

- Shortened RBC life span -HbA1c test will not be accurate when a person has a condition that affects the average lifespan of red blood cells (RBCs), such as hemolytic anemia or blood loss. When the lifespan of RBCs in circulation is shortened, the A1c result is falsely low and is an unreliable measurement of a person's average glucose over time.
- Abnormal forms of hemoglobin - The presence of some hemoglobin variants, such as hemoglobin S in sickle cell anemia, may affect certain methods for measuring A1c. In these cases, fructosamine can be used to monitor glucose control.

Advised:

- To follow patient for glycemic control test like fructosamine or glycated albumin may be performed instead.
 - Hemoglobin HPLC screen to analyze abnormal hemoglobin variant.
- estimated Average Glucose (eAG) : based on value calculated according to National Glycohemoglobin Standardization Program (NGSP) criteria.

MGR

Technologist

Page No: 5 of 15

Tanu

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- **MR. LALIT KUMAR**

Age :- 30 Yrs 5 Mon

Sex :- Male

Patient ID :-12222951

Date :- 28/01/2023

10:13:17

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 28/01/2023 16:24:42

HAEMATOLOGY

BLOOD GROUP ABO

Method:- Haemagglutination reaction

"B" POSITIVE



MGR

Technologist

Page No: 6 of 15

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 744 4824885 maxcarediagnostics1@gmail.com



NAME :- MR. LAIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

LIPID PROFILE

TOTAL CHOLESTEROL Method:- CHOD-PAP methodology	149.00	mg/dl	Desirable <200 Borderline 200-239 High > 240
---	--------	-------	--

InstrumentName:MISPA PLUS **Interpretation:** Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.

TRIGLYCERIDES Method:- GPO-TOPS methodology	122.00	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
---	--------	-------	--

InstrumentName:MISPA PLUS **Interpretation :** Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL Method:- Selective inhibition Method	64.00	mg/dl	Male 35-80 Female 42-88
---	-------	-------	----------------------------

Instrument Name:MISPA PLUS **Interpretation:** An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

LDL CHOLESTEROL Method:- Calculated Method	64.67	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
--	-------	-------	--

VLDL CHOLESTEROL Method:- Calculated	24.40	mg/dl	0.00 - 80.00
--	-------	-------	--------------

T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	2.33		0.00 - 4.90
---	------	--	-------------

LDL / HDL CHOLESTEROL RATIO Method:- Calculated	1.01		0.00 - 3.50
---	------	--	-------------

TOTAL LIPID Method:- CALCULATED	477.67	mg/dl	400.00 - 1000.00
---	--------	-------	------------------

1. Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.
2. As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended
3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

Comments: 1- ATP III suggested the addition of Non HDL Cholesterol (Total Cholesterol – HDL Cholesterol) as an indicator of all MGR

Technologist
Page No: 7 of 15

Tanu
DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

BIOCHEMISTRY

LIVER PROFILE WITH GGT

SERUM BILIRUBIN (TOTAL) Method:- DMSO/Diazo	0.63	mg/dL	Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Method:- DMSO/Diazo	0.24	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.39	mg/dl	0.30-0.70
SGOT Method:- IFCC	24.9	U/L	Men- Up to - 37.0 Female - Up to - 31.0
SGPT Method:- IFCC	39.6	U/L	Men- Up to - 40.0 Female- Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- DGKC - SCE	124.00	U/L	80.00 - 306.00

InstrumentName:MISPA PLUS **Interpretation:**Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobiliary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

SERUM GAMMA GT Method:- Szasz methodology Instrument Name Randox Rx Imola	17.20	U/L	10.00 - 45.00
---	-------	-----	---------------

Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

SERUM TOTAL PROTEIN Method:- Direct Biuret Reagent	5.60	g/dl	5.10 - 8.00
SERUM ALBUMIN Method:- Bromocresol Green	3.85	g/dl	3.50 - 5.50
SERUM GLOBULIN Method:- CALCULATION	1.75 L	gm/dl	2.20 - 3.50
A/G RATIO	2.20		1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note :- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B ,C ,paracetamol toxicity etc.Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.

MGR

Technologist
Page No: 9 of 15

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA 17.10 mg/dl 10.00 - 50.00
Method:- Urease/GLDH

InstrumentName: HORIBA CA 60 **Interpretation :** Urea measurements are used in the diagnosis and treatment of certain renal and metabolic diseases.

SERUM CREATININE 1.37 mg/dl Males : 0.6-1.50 mg/dl
Females : 0.6 -1.40 mg/dl
Method:- Jaffe's Method

Interpretation : Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not clinically significant.

SERUM URIC ACID 5.76 mg/dl 2.40 - 7.00

InstrumentName:HORIBA YUMIZEN CA60 Daytona plus **Interpretation: Elevated Urate:**High purine diet,Alcohol• Renal insufficiency,Drugs , Polycythaemia vera, Malignancies,Hypothyroidism,Rare enzyme defects ,Downs syndrome,Metabolic syndrome, Pregnancy,Gout.

SODIUM 132.5 L mmol/L 135.0 - 150.0
Method:- ISE

Interpretation: Decreased sodium - Hyponatraemia Causes include: fluid or electrolyte loss,Drugs,Oedematous states,Legionnaire's disease and other chest infections,pseudonatremia, Hyperlipidaemias and paraproteinaemias,endocrine diseases ,SIADH.

POTASSIUM 4.30 mmol/L 3.50 - 5.10
Method:- Ion-Selective Electrode with Serum

Interpretation: A. Elevated potassium (hyperkalaemia)• Artefactual,Physiological activation,Drugs, Pathological states,Renal failure Adrenocortical insufficiency, metabolic acidoses, very high platelet or white cell counts B. Decreased potassium (hypokalaemia)Drugs, Liqueuric,Diarrhoea and vomiting,Metabolic alkalosis,Corticosteroid excess, Oedematous state,Anorexia nervosa/bulimia

CHLORIDE 102.8 mmol/L 98.0 - 107.0
Method:- Ion-Selective Electrode with Serum

Interpretation: Used for Electrolyte monitoring.

SERUM CALCIUM 11.00 mg/dl 8.10 - 11.50
Method:- Colorimetric method

InstrumentName:Rx Daytona plus **Interpretation:** Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia .Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN 5.60 g/dl 5.10 - 8.00
Method:- Direct Biuret Reagent

SERUM ALBUMIN 3.85 g/dl

Method:- Bromocresol Green

Technologist

Page No: 10 of 15

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 28/01/2023 16:24:42

BIOCHEMISTRY


SERUM GLOBULIN Method:- CALCULATION	1.75 L	gm/dl	2.20 - 3.50
A/G RATIO	2.20		1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR .in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.



MGR

Technologist

Page No: 11 of 15

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

TOTAL THYROID PROFILE

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

THYROID-TRIODOXYRONINE T3 Method:- ECLIA	1.17	ng/mL	0.70 - 2.04
--	------	-------	-------------

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level. 2.Low TSH, high FT4 and TSH receptor antibody (TRAb) +ve seen in patients with Graves disease 3.Low TSH, high FT4 and TSH receptor antibody (TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.HighTSH, Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis 5.HighTSH, Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency 6.Low TSH, Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism 7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels 8.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis 9.Normal or ↑ T3 & ↑ T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism . 11.Normal T3 & ↑ T4 along with ↓ TSH is seen in Hypothyroidism . 12.Normal T3 & T4 levels with ↓ TSH indicate Mild / Subclinical Hypo

DURING PREGNANCY - REFERENCE RANGE for TSH in uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL 2nd Trimester : 0.20-3.00 uIU/mL 3rd Trimester : 0.30-3.00 uIU/mL The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly. *** 5.10 - 14.10

THYROID THYRONINE (T4) Method:- ECLIA	8.09	μIU/mL	5.10 - 14.10
---	------	--------	--------------

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level. 2.Low TSH, high FT4 and TSH receptor antibody (TRAb) +ve seen in patients with Graves disease 3.Low TSH, high FT4 and TSH receptor antibody (TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.HighTSH, Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis 5.HighTSH, Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency 6.Low TSH, Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism 7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels 8.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis 9.Normal or ↑ T3 & ↑ T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism . 11.Normal T3 & ↑ T4 along with ↓ TSH is seen in Hypothyroidism . 12.Normal T3 & T4 levels with ↓ TSH indicate Mild / Subclinical Hypo

DURING PREGNANCY - REFERENCE RANGE for TSH in uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL 2nd Trimester : 0.20-3.00 uIU/mL 3rd Trimester : 0.30-3.00 uIU/mL The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

TSH Method:- ECLIA	4.200	μIU/mL	0.350 - 5.500
------------------------------	-------	--------	---------------

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level.

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
Page No: 14 of 15



B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 28/01/2023 16:24:42

IMMUNOASSAY


- 2.Low TSH,high FT4 and TSH receptor antibody(TRAb) +ve seen in patients with Graves disease
- 3.Low TSH,high FT4 and TSH receptor antibody(TRAb) -ve seen in patients with Toxic adenoma/Toxic Multinodular goiter
- 4.HighTSH,Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimotos thyroiditis
- 5.HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 synthesis deficiency
- 6.Low TSH,Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism
- 7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
- 8.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 9.Normal or ↓ T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 10.Normal T3 & T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism .
- 11.Normal T3 & ↓ T4 along with ↑ TSH is seen in Hypothyroidism .
- 12.Normal T3 & T4 levels with ↑ TSH indicate Mild / Subclinical Hypothyroidism .
- 13.Slightly ↑ T3 levels may be found in pregnancy and in estrogen therapy while ↓ levels may be encountered in severe illness , malnutrition , renal failure and during therapy with drugs like propranolol.
- 14.Although ↑ TSH levels are nearly always indicative of Primary Hypothyroidism ,rarely they can result from TSH secreting pituitary tumours.

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association)

- 1st Trimester : 0.10-2.50 uIU/mL
- 2nd Trimester : 0.20-3.00 uIU/mL
- 3rd Trimester : 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved.TSH is an important marker for the diagnosis of thyroid dysfunction.Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age ,and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.



MGR

Technologist

Page No: 15 of 15

Tanu

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com




NAME :- Mr. LALIT KUMAR	Patient ID :-12222951	Date :- 28/01/2023	10:13:17
Age :- 30 Yrs 5 Mon	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 28/01/2023 16:24:42

CLINICAL PATHOLOGY



Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
<u>PHYSICAL EXAMINATION</u>			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
<u>CHEMICAL EXAMINATION</u>			
REACTION(PH)	5.5		5.0 - 7.5
SPECIFIC GRAVITY	1.010		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIVE		NEGATIVE
UROBILINOGEN	NORMAL		NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
<u>MICROSCOPY EXAMINATION</u>			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT



MGR

Technologist
Page No: 12 of 15

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226





MR. LALIT KUMAR	30 Y/Male
Registration Date: 28/01/2023	Ref. by: BANK OF BARODA

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (14.9 cm). **Echotexture is increased obscuring periportal echogenicity.** No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape (11.3 cm). Echotexture is normal. No focal lesion is seen.

Right kidney is noted in right pelvic region, adjacent to right lateral wall of urinary bladder, measuring approx. 11.7 x 4.7 cm with hilum facing anteromedially – ***suggestive of ectopic right kidney.*** Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Left kidney is measuring approx. 12.0 x 5.2 cm. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Urinary bladder does not show any calculus or mass lesion.

Prevoid: 261 mL Postvoid: 22-23 mL (not significant)

Prostate is normal in size with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified.

No significant free fluid is seen in pelvis.

IMPRESSION:

- **Grade 2 fatty liver.**
- **Ectopic right kidney as described above.**

DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave - II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- ☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME:	MR. LALIT KUMAR	AGE	30 YRS/M
REF.BY	BANK OF BARODA	DATE	29/01/2023

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.


Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected.

DR.SHALINI GOEL
M.B.B.S, D.N.B (Radiodiagnosis)
RMC No.: 21954




PR Interval: 164 ms

QRS Duration: 76 ms

QT/QTc: 372/362ms

P-QRS-T Axis: 35 - 41 - 24 (Deg)



FINDINGS: Abnormal ECG with Indication of Sinus Bradycardia

Vent Rate : 56 bpm; PR Interval : 164 ms; QRS Duration: 76 ms; QT/QTc Int : 372/362 ms

P-QRS-T axis: 35 • 41 • 24 • (Deg)

Comments :

Baseline sinus bradycardia

(Signature)

Dr. Naresh Kumar Mohanka

RMC No.: 35703

IBBS, DIP, CARDIO (ESCORTS)

D.E.M. (RCGP-UK)



'S HEALTH SOLUTIONS LLP
B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur
 1322381/MR LALIT KUMAR 30 Yrs/Male 0 Kg/0 Cms
 Date: 28-Jan-2023 11:32:15 AM
 Ref. By : BANK OF BARODA

Protocol : BRUCE

Medication :
 Objective :

History :

Stage	StageTime (Min:Sec)	PhaseTime (Min:Sec)	Speed (mph)	Grade (%)	METS	H.R. (bpm)	B.P. (mmHg)	R.P.P. x100	PVC	Comments
Supine					1.0	59	120/80	70	-	
Standing					1.0	85	120/80	102	-	
HV					1.0	74	120/80	88	-	
HV					1.0	78	120/80	93	-	
EXStart					1.0	89	120/80	106	-	
Stage 1	3:01	3:02	1.7	10.0	4.7	124	130/80	161	-	
Stage 2	3:01	6:02	2.5	12.0	7.1	156	140/80	218	-	
PeakEx	1:16	7:17	3.4	14.0	8.4	171	150/85	256	-	
Recovery	1:00		0.0	0.0	1.2	135	150/85	202	-	
Recovery	2:00		0.0	0.0	1.0	104	160/85	166	-	
Recovery	3:00		0.0	0.0	1.0	97	150/85	145	-	
Recovery	4:00		0.0	0.0	1.0	91	140/80	127	-	

Findings :

Exercise Time :07:16


Max HR Attained :171 bpm 90% of Max Predictable HR 190

Max BP : 160/85(mmHg)

Max Workload attained :8.4(Fair Effort Tolerance)

TMT is Negative for RVT.

Advice/Comments:



Lalit



322381/MR LALIT KUMAR
 0 Yrs/Male

Kg/0 Cms
 Date: 28-Jan-2023 11:32:15 AM
 4X 103 ms Post-J


HR: 59 bpm
 METS: 1.0
 BP: 120/80

MPHR: 31% of 190
 Speed: 0.0 mph
 Grade: 0.0%

Raw ECG
 BRUCE
 (1.0-35)HZ

Ex Time 00:31
 BLC: On
 Notch: On

Supine
 5.0 mm/mV
 25 mm/Sec



'S HEALTH SOLUTIONS LLP
B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur

322381/MR LALIT KUMAR
0 Yrs/Male

Kg/0 Cms

Date: 28-Jan-2023 11:32:15 AM
4X 103 ms Post J


HR: 72 bpm
METS: 1.0
BP: 120/80

APHR: 37% of 190
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(1.0-35)Hz

Ex Time 01:10
BLC : On
Notch : On

HV
5.0 mm/mV
25 mm/Sec.



322381/MR LALIT KUMAR
 0 Yrs/Male

Kg/0 Cms
 Date: 28-Jan-2023 11:32:15 AM
 4X 103 ms Post-J


HR: 91 bpm
 METS: 1.0
 BP: 120/80

MPHR: 47% of 190
 Speed: 0.0 mph
 Grade: 0.0%

Raw ECG
 BRUCE
 (1.0-35)HZ

Ex Time 01:01
 BLC : On
 Notch : On

Standing
 5.0 mm/mV
 25 mm/Sec.



322381/MR LALIT KUMAR
 0 Yrs/Male
 Kg/0 Cms

Date: 28-Jan-2023 11:32:15 AM
 4X 103 ms Post J

HR: 125 bpm
 METS: 4.7
 BP: 130/80



MPHR: 65% of 190
 Speed: 1.7 mph
 Grade: 10.0%

Raw ECG
 BRUCE
 (1.0-35)Hz

Ex Time 02:59
 BLC: On
 Notch: On

BRUCE: Stage 1 (3:00)
 5.0 mm/mV
 25 mm/Sec

1.2 Lead + Median



'S HEALTH SOLUTIONS LLP
B-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur
322381/MR LALIT KUMAR
0 Yrs/Male
Kg/0 Cms

Date: 28-Jan-2023 11:32:15 AM
4X 103 ms Post J



HR: 156 bpm
METs: 7.1
BP: 140/80

MPHR: 82% of 190
Speed: 2.5 mph
Grade: 12.0%

Raw ECG
BRUCE
(1.0-35)Hz

Ex Time 05:59
BLC: On
Notch: On

BRUCE: Stage 2(3:00)
5.0 mm/mV
25 mm/Sec



I 1.2 1.2 1.3
II 1.9 2.5 1.2
III 0.8 1.2 0.8
aVR -1.5 -1.9
aVL 0.2 0.0
aVF 1.4 1.8
V1 -0.8 -1.6
V2 1.0 1.2
V3 0.8 2.4
V4 0.5 2.1
V5 0.5 2.2
V6 0.3 2.0

322381/MR LALIT KUMAR
 0 Yrs/Male
 Kg/0 Cms
 Date: 28-Jan-2023 11:32:15 AM
 4X 103 ms Post J


HR: 171 bpm
 METS: 8.4
 BP: 150/85

MPHR: 90% of 190
 Speed: 3.4 mph
 Grade: 14.0%

Raw ECG
 BRUCE
 (1.0-35)Hz

Ex Time 07:14
 BLC : On
 Notch : On

BRUCE: PeakEx(1:14)
 5.0 mm/mV
 25 mm/Sec.



322381/MR LALIT KUMAR
 0 Yrs/Male
 Kg/0 Cms
 Date: 28-Jan-2023 11:32:15 AM
 4X 103 ms Post J



HR: 136 bpm
 METS: 1.3
 BP: 150/85

MPHR: 71% of 190
 Speed: 0.0 mph
 Grade: 0.0%

Raw ECG
 BRUCE
 (1.0-35)Hz

Ex Time 07:16
 BLC: On
 Notch: On

Recovery(1:00)
 5.0 mm/mV
 25 mm/Sec



322381/MR LALIT KUMAR
 0 Yrs/Male
 Kg/0 Cms
 Date: 28-Jan-2023 11:32:15 AM
 4X 103 ms Post-J


HR: 101 bpm
 METS: 1.0
 Bp: 160/85

MPHR: 53% of 190
 Speed: 0.0 mph
 Grade: 0.0%

Raw ECG
 BRUCE
 (1.0-35)Hz

Ex Time 07:16
 BLC: On
 Notch: On

Recovery(2:00)
 5.0 mm/mV
 25 mm/Sec



'S HEALTH SOLUTIONS LLP
3-14, Vidhyadhar Nagar Enclave, Phase -2, Jaipur
322381/MR LALIT KUMAR
0 Yrs/Male
Kg/0 Cms

Date: 28-Jan-2023 11:32:15 AM
4X 103 ms Post J




HR: 97 bpm
METs: 1.0
BP: 150/85







MpHR: 51% of 190
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(1.0-35)Hz

Ex Time 07:16
BLC: On
Notch: On

Recovery(3:00)
5.0 mm/mV
25 mm/Sec








HR: 90 bpm
 METS: 1.0
 BP: 140/80







MPHR: 47% of 190
 Speed: 0.0 mph
 Grade: 0.0%


Raw ECG
 BRUCE
 (1.0-35)Hz

Ex Time 07:16
 BLC: On
 Notch: On

Recovery(4:00)
 5.0 mm/mV
 25 mm/Sec.







12222951 MR LALIT KUMAR 30 YRS BANK OF BARODA M
28.JAN.2023
MAXCARE DIAGNOSTIC (ASSOCIATES OF P3 HEALTH SOLUTIONS LLP)

