Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



General Physical Examination

Date of Examination: 26-11-22	
Name: RAJAT MATHUR.	Age:Sex:
DOB: 91 11 1989.	
Referred By: BOB-	
Photo ID: Nadhar . ID #: attach	ec
Ht: 170 (cm)	Wt: Kg)
Chest (Expiration): (cm)	Abdomen Circumference: 92. (cm)
Blood Pressure: 150/ 90 mm Hg PR: 18/	min RR: 15/min Temp: Alebane
	*
вмі 26.6	
Eye Examination: Dis Misron, 66	Dormal Color Vision.
M6 BIC eyes.	Dormal Color VIEron.
Other: Not Significan	2-
On examination he/she appears physically and me	ntally fit: Yes / No
On examination he/she appears physically and me	
Signature Of Examine :	Name of Examinee:
Signature of Examine .	h Goyal
Simple Madical Evaniant	Name Medical Examiner -047996
Signature Medical Examiner	M.B. Reg No .01
RAJAT MATHUR 09 @ CON	Name of Examinee:
9285599554,	



For for Good deky



Or PIVIL D.M.R.D. 8



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male

33 Yrs

Company :- MediWheel

Patient ID: -122228279 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 26/11/2022 10:59:04

BOB PACKAGE BELOW 40MALE

X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression :- Normal Study

(Please correlate clinically and with relevant further investigations)

DR ABHISHEK JAIN MBBS. DNB. (RADIO DIAGNOSIS)

RMC NO. 21687

*** End of Report ***

Page No: 1 of 1


AHSAN

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male 33 Yrs

Company :- MediWheel

Sample Type :- EDTA

Patient ID :-122228279

Ref. By Dr:- BOB

Lab/Hosp:-

Final Authentication: 26/11/2022 14:41:26

HAEMATOLOGY

Sample Collected Time 26/11/2022 09:13:20

Test Name	Value	Unit	Biological Ref Interval
BOB PACKAGE BELOW 40MALE		4	
HAEMOGARAM			
HAEMOGLOBIN (Hb)	16.0	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	8.94	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	55.6	%	40.0 - 80.0
LYMPHOCYTE	37.7	%	20.0 - 40.0
EOSINOPHIL	3.4	%	1.0 - 6.0
MONOCYTE	3.0	%	2.0 - 10.0
BASOPHIL	0.3	%	0.0 - 2.0
NEUT#	4.98	10^3/uL	1.50 - 7.00
LYMPH#	3.37	10^3/uL	1.00 - 3.70
EO#	0.30	10^3/uL	0.00 - 0.40
MONO#	0.26	10^3/uL	0.00 - 0.70
BASO#	0.03	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	5.44	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	45.70	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	84.1	fL '	83.0 - 101.0
MEAN CORP HB (MCH)	29.4	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	34.1	g/dL	31.5 - 34.5
PLATELET COUNT	251	x10^3/uL	150 - 410
RDW-CV	13.1	%	11.6 - 14.0
MENTZER INDEX	15.46		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

MUKESHSINGH Technologist

Page No: 1 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR

33 Yrs

Sex / Age :- Male Company :- MediWheel

Patient ID: -122228279

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 26/11/2022 14:41:26

Sample Collected Time 26/11/2022 09:13:20 HAEMATOLOGY

Test Name

Value

Biological Ref Interval

Erythrocyte Sedimentation Rate (ESR)

18 H

mm/hr.

00 - 13

(ESR) Methodology: Measurment of ESR by cells aggregation.

Sample Type :- EDTA

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation

: ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia of Connective disease and MCH, MCV, MCHC, MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L, Japan

MUKESHSINGH **Technologist**

Page No: 2 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male Company :- MediWheel

Test Name

33 Yrs

Patient ID: -122228279

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type: - EDTA, KOx/Na FLUORIDE-F, KSWINGETCHORIDETTRE DECIME 022 09:13:20 HAEMATOLOGY

Value

Biological Ref Interval

Final Authentication: 26/11/2022 14:41:26

BLOOD GROUP ABO

" B" NEGATIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone).

FASTING BLOOD SUGAR (Plasma)

Method:- GOD PAP

109.3

mg/dl

75.0 - 115.0

Impaired glucose tolerance (IGT) 111 - 125 mg/dL Diabetes Mellitus (DM) > 126 mg/dL

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases

BLOOD SUGAR PP (Plasma)

Method:- GOD PAP

120.3

mg/dl

70.0 - 140.0

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases

URINE SUGAR (FASTING) Collected Sample Received

Nil

Nil

KAUSHAL, MKSHARMA, MUKESHSINGH, POOJABOHRA

Technologist DR.HANSA Page No: 3 of 11



Dr. Piyush Goyal (D.M.R.D.) Dr. Chandrika Gupta

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male

33 Yrs

Company :- MediWheel
Sample Type :- STOOL

Patient ID :-122228279

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 26/11/2022 11:51:09

Sample Collected Time 26/11/2022 09:13:20

CLINICAL PATHOLOGY

Test Name

Value

Unit

Biological Ref Interval

STOOLANALYSIS

PHYSICAL EXAMINATION

COLOUR

CONSISTENCY

MUCUS

BLOOD

MICROSCOPIC EXAMINATION

RBC's

WBC/HPF

MACROPHAGES

OVA

CYSTS

TROPHOZOITES

CHARCOT LEYDEN CRYSTALS

OTHERS

Collected Sample Received

YELLOW BROWN

SEMI SOLID

ABSENT

ABSENT

NII.

/HPF

/HPF

0 - 1

ABSENT

ABSENT

ABSENT

ABSENT

ABSENT

NORMAL BACTERIA FLORA PRESENT

POOJABOHRA Technologist DR.HANSA Page No: 4 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgovalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male

33 Yrs

Company :- MediWheel

Patient ID: -122228279

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type :- PLAIN/SERUM

Sample Collected Time 26/11/2022 09:13:20

Final Authentication: 26/11/2022 11:46:37

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	190.94	mg/dl	Desirable <200 Borderline 200-239 High> 240
TRIGLYCERIDES Method:- GPO-PAP	195.37 H	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	34.15	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:-Direct clearance Method	124.23	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	39.07	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	5.59 H		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	3.64 H		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola	646.25	mg/dl	400.00 - 1000.00

EROL InstrumentName: Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obst

DIRECT HDLCHOLESTERO InstrumentName: Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROLInstrumentName: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its prog TOTAL LIPID AND VLDL ARE CALCULATED ress and to avoid plaque rupture

MKSHARMA

Page No: 5 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

Sample Type - PLAIN/SERUM

Sex / Age :- Male

NAME :- Mr. RAJAT MATHUR 33 Yrs

Company :- MediWheel

Patient ID :-122228279

Ref. By Dr:- BOB

Lab/Hosp:-

Final Authentication: 26/11/2022 11:46:37

Sample Collected Time 26/11/2022 09:13:20 RIOCHEMISTRY

nterval

BIOCHEMISTRY					
Test Name	Value	Unit	Biological Ref In		
LIVER PROFILE WITH GGT					
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.58	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12 1month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2 Ref-(ACCP 2020)		
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.14	mg/dL ▲	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL		
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.44	mg/dl	0.30-0.70		
SGOT Method:- IFCC	21.4	U/L	Men- Up to - 37.0 Women - Up to - 31.0		
SGPT Method:- IFCC	25.3	U/L	Men- Up to - 40.0 Women - Up to - 31.0		
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	116.60	IU/L	30.00 - 120.00		
SERUM GAMMA GT Method:- IFCC	48.10	U/L	11.00 - 50.00		
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.20	g/dl	6.40 - 8.30		
SERUM ALBUMIN Method:- Bromocresol Green	4.74	g/dl	3.80 - 5.00		
SERUM GLOBULIN Method:-CALCULATION	2.46	gm/dl	2.20 - 3.50		
A/G RATIO	1.93		1.30 - 2.50		

Total BilirubinMethodology Colorimetric method InstrumentName;Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duet and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving

the haemoglobin it is receiving.

All operate Aminotransferase Methodology IFCC InstrumentName Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans. ALT Alanine Aminotransferase Methodology: IFCCInstrumentName.Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular

At Machine Phosphatase Methodology AMP Buffer InstrumentName Randox Rx Imola Interpretation Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of licepatobility disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology Burret Reagent InstrumentName Randox Rx Imola Interpretation. Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromoeresol Green InstrumentName Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving e liver or kidneys. Globulin & A/G ratio is calculated

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal)

MKSHARMA

Page No: 6 of 11

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Sex / Age :- Male 33 Yrs

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Patient ID :-122228279

Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 26/11/2022 11:46:37

BIOCHEMISTRY

Sample Collected Time 26/11/2022 09:13:20

	DIOCHEN	HSTRY	
Test Name	Value	Unit	Biological Ref Interval
SERUM CREATININE Method:- Colorimetric Method	0.92	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20
SERUM URIC ACID Method:- Enzymatic colorimetric	5.50	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7

MKSHARMA

Page No: 7 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR Sex / Age :- Male 33 Yrs

Company :- MediWheel Sample Type :- PLAIN/SERUM Patient ID: -122228279

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 26/11/2022 09:13:20

Final Authentication: 26/11/2022 11:46:37

0.0 - 23.0

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
BLOOD UREA NITROGEN (BUN)	11.7	mg/dl	0.0 - 23.0

MKSHARMA

Page No: 8 of 11



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR Sex / Age :- Male

33 Yrs

Company :-MediWheel Sample Type - EDTA

Patient ID :-122228279

Ref. By Dr:- BOB

Lab/Hosp :-

HAEMATOLOGY

Final Authentication: 26/11/2022 14:41:26

Sample Collected Time 26/11/2022 09:13:20

Test Name

Value

Unit

Biological Ref Interval

GLYCOSYLATED HEMOGLOBIN (HbA1C) Method:- HPLC

6.4 H

Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4

Diabetics: = 6.5 or higher

ADA Target: 7.0 Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base.It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1cmeasurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

Method:- Calculated Parameter

137 H

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

MUKESHSINGH Technologist

Page No: 9 of 11



Dr. Chandrika Gupta MBBS.MD (Path) RMC NO. 21021/008037

"CONDITIONS OF REPORTING SEE OVER LEAF"

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

33 Yrs



:- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male Company :- MediWheel

Sample Type :- URINE

Patient ID :-122228279 Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication . 26/11/2022 11.51 09

Sample Collected Time 26/11/2022 09:13:20 **CLINICAL PATHOLOGY**

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
COLOUR	PALE YE	LLOW	PALE YELLOW
APPEARANCE	Clear	000 11	Clear
CHEMICAL EXAMINATION			Clear
REACTION(PH)	6.0		5.0 - 7.5
SPECIFIC GRAVITY	1.025		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIV	E ,	NEGATIVE
UROBILINOGEN	NORMAL		NORMAL.
KETONES	NEGATIV	Е	NEGATIVE
NITRITE	NEGATIV	E	NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		

POOJABOHRA **Technologist** DR.HANSA Page No: 10 of 11



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 26/11/2022 08:49:52

NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male 33 Yrs

Company :- MediWheel
Sample Type :- PLAIN/SERUM

Patient ID :-122228279

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 26/11/2022 09:13:20

Final Authentication: 26/11/2022 12:32:37

IMMUNOASSAY

	ALJEST A. A.	
Value	Unit	Biological Ref Interval
1.040	ng/ml	0.970 - 1.690
6.980	ug/dl	5.530 - 11.000
2.750	μIU/mL	0.400 - 4.649
	1.040 6.980	1.040 ng/ml 6.980 ug/dl

Interpretation: Trilodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT41) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overthypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

*** End of Report ***

NARENDRAKUMAR Technologist

Page No: 11 of 11



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52 Date

NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male 33 Yrs

Company :- MediWheel

Patient ID: -122228279

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Type :-

..........

Sample Collected Time

Final Authentication: 26/11/2022 13:17:14

BOB PACKAGE BELOW 40MALE

2D ECHO OPTION TMT (ADULT/CHILD)

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	4	NOR	MAL	TRICUS	SPID VALVE		NORMAL	
AORTIC VALVE		NOR	MAL	PULMO	DNARY VALVE		NORMAL	0
		M.MODE E	XAMITATION:		, E4			
AO .	21	mm	LA	29	Mm	IVS-D	10	mm
IVS-S	17	mm	LVID	44	Mm	LVSD	27	mm
LVPW-D	10	mm	LVPW-S	16	Mm	RV		mm
RVWT		mm	EDV		МІ	LVVS		ml
LVEF	69%		-	RWMA	1	ABSENT		
				CHAI	MBERS:			

LA	NORMAL	RA	NORMAL	
LV	NORMAL	RV	NORMAL	
PERICARDIUM		NORMAL		

COLOUR DOPPLER:

	MI	TRAL VALVE	E				
E VELOCITY	0.68	m/sec	PEAK GRADIENT			Mn	n/hg
A VELOCITY	0.48	m/sec	MEAN	GRADIEN	т	Mm/hg	
MVA BY PHT		Cm2	MVA	BY PLANIM	ETRY	Cm	2
MITRAL REGURGITAT	ION				ABSENT		
	AO	RTIC VALVE					
PEAK VELOCITY	1.41	m/	sec	PEAK G	RADIENT	m	m/hg
AR VMAX		m/	sec MEAN GRADIENT		RADIENT	m	m/hg
AORTIC REGURGITATION				ABSENT			
	TRIC	USPID VAL	VE		G.		
PEAK VELOCITY	0.55		m/sec	PEAK G	PEAK GRADIENT		mm/hg
MEAN VELOCITY			m/sec	MEAN GRADIENT			mm/hg
VMax VELOCITY		1			-		
TRICUSPID REGURGIT	TATION			ABSENT			
*	PUI	MONARY \	VALVE				
PEAK VELOCITY		1.2		M/sec. PEAK GRADIENT			Mm/hg
MEAN VALOCITY	122.50	,			MEAN GRADIENT		Mm/hg
PULMONARY REGUR	GITATION			-	ABSENT		

VIKAS

Page No: 1 of 2



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Final Authentication: 26/11/2022 13:17:14

Date :- 26/11/2022 08:49:52 NAME :- Mr. RAJAT MATHUR

Sex / Age :- Male 33 Yrs

Company :- MediWheel

Patient ID: -122228279 Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type :-Sample Collected Time

Impression--

- 1. Normal LV size & contractility.
- 2. No RWMA, LVEF 69%.
- 3. Normal cardiac chamber.
- 4. Normal valve.
- 5. No clot, no vegetation, no pericardial effusion.

(Cardiologist)

*** End of Report ***

VIKAS

Page No: 2 of 2





B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 26/11/2022 08:49:52

NAME

:- Mr. RAJAT MATHUR

Sex / Age :- Male

33 Yrs

Company :- MediWheel

Patient ID :-122228279 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 26/11/2022 12:08:12

BOB PACKAGE BELOW 40MALE

USG WHOLE ABDOMEN

Liver is of normal size. Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Urinary bladder is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Prostate is normal in size (11.9 gms) with normal echo-texture and outline.

No enlarged nodes are visualised. No retro-peritoneal lesion is identified No significant free fluid is seen in peritoneal cavity.

IMPRESSION:

Normal study

Needs clinical correlation for further evaluation

*** End of Report ***

C- Abhishek Jain Pagano to Bra DNB, (Radio Diagnosis) KMC No. 21 687

AHSAN

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant

Dr. Rathod Hetali Amrutlal MBBS, M.D. (Radio-Diagnosis) RMC No. 17163

Transcript by


Name: RAJAT MATHUR / M





Dr. Goyal's Path Lab


Name rajat mathur Patient ld RAJAT91_91389 Date 11/26/2022 Diagnosis Dr.














Allengers ECG (Pisces)(PIS218210312) DR. GOYALS PATH LAB & IMAGING CENTER

102220941 / MR RAJAT MATHUR / 33 Yrs / M/ Non Smoker

Heart Rate: 74 bpm / Tested On: 26-Nov-22 13:02:39 / HF 0.05 Hz - LF 100 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s / Refd By.: BOB MBBS, DIP, CARDIO (ESCORTS)
D.E.M (RCGP-UK) Dr. Nados Akumar Mollanken RMC Reported By: ECG