



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

**MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE****BLOOD COUNTS,EDTA WHOLE BLOOD**

HEMOGLOBIN (HB)	12.9	12.0 - 15.0	g/dL
METHOD : SPECTROPHOTOMETRY			
RED BLOOD CELL (RBC) COUNT	4.52	3.8 - 4.8	mil/ $\mu$ L
METHOD : ELECTRICAL IMPEDANCE			
WHITE BLOOD CELL (WBC) COUNT	6.40	4.0 - 10.0	thou/ $\mu$ L
METHOD : ELECTRICAL IMPEDANCE			
PLATELET COUNT	354	150 - 410	thou/ $\mu$ L
METHOD : ELECTRICAL IMPEDANCE			

**RBC AND PLATELET INDICES**

HEMATOCRIT (PCV)	39.8	36 - 46	%
METHOD : CALCULATED			
MEAN CORPUSCULAR VOLUME (MCV)	88.0	83 - 101	fL
METHOD : CALCULATED			
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	28.5	27.0 - 32.0	pg
METHOD : CALCULATED			
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	32.3	31.5 - 34.5	g/dL
METHOD : CALCULATED			
RED CELL DISTRIBUTION WIDTH (RDW)	12.3	11.6 - 14.0	%
METHOD : CALCULATED			
MENTZER INDEX	19.5		
MEAN PLATELET VOLUME (MPV)	7.8	6.8 - 10.9	fL
METHOD : CALCULATED			

**WBC DIFFERENTIAL COUNT**

NEUTROPHILS	56	40 - 80	%
METHOD : IMPEDANCE / MICROSCOPY			
LYMPHOCYTES	31	20 - 40	%
METHOD : IMPEDANCE / MICROSCOPY			
MONOCYTES	09	2 - 10	%
METHOD : IMPEDANCE / MICROSCOPY			
EOSINOPHILS	04	1 - 6	%
METHOD : IMPEDANCE / MICROSCOPY			
BASOPHILS	00	0 - 2	%
METHOD : IMPEDANCE / MICROSCOPY			
ABSOLUTE NEUTROPHIL COUNT	3.58	2.0 - 7.0	thou/ $\mu$ L



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
METHOD : CALCULATED				
ABSOLUTE LYMPHOCYTE COUNT		1.98	1.0 - 3.0	thou/ $\mu$ L
METHOD : CALCULATED				
ABSOLUTE MONOCYTE COUNT		0.58	0.2 - 1.0	thou/ $\mu$ L
METHOD : CALCULATED				
ABSOLUTE EOSINOPHIL COUNT		0.26	0.02 - 0.50	thou/ $\mu$ L
METHOD : CALCULATED				
<b>ERYTHROCYTE SEDIMENTATION RATE (ESR),WHOLE BLOOD</b>				
E.S.R		<b>29</b>	<b>High</b> 0 - 20	mm at 1 hr
METHOD : MODIFIED WESTERGREN				
<b>GLUCOSE FASTING,FLUORIDE PLASMA</b>				
FBS (FASTING BLOOD SUGAR)		89	74 - 99	mg/dL
METHOD : HEXOKINASE				
<b>GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD</b>				
HBA1C		5.2	Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 Therapeutic goals: < 7.0 Action suggested : > 8.0 (ADA Guideline 2021)	%
METHOD : HPLC TECHNOLOGY				
ESTIMATED AVERAGE GLUCOSE(EAG)		102.5	< 116.0	mg/dL
<b>GLUCOSE, POST-PRANDIAL, PLASMA</b>				
PPBS(POST PRANDIAL BLOOD SUGAR)		96	Normal: < 140, Impaired Glucose Tolerance:140-199 Diabetic > or = 200	mg/dL
METHOD : HEXOKINASE				
<b>LIPID PROFILE, SERUM</b>				
CHOLESTEROL, TOTAL		132	Desirable: <200 BorderlineHigh : 200-239 High : > or = 240	mg/dL
METHOD : OXIDASE, ESTERASE, PEROXIDASE				
TRIGLYCERIDES		110	Desirable: < 150 Borderline High: 150 - 199 High: 200 - 499 Very High : > or = 500	mg/dL
METHOD : ENZYMATIC ASSAY				
HDL CHOLESTEROL		<b>34</b>	<b>Low</b> < 40 Low > or = 60 High	mg/dL
METHOD : DIRECT- NON IMMUNOLOGICAL				



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002692211

**CLIENT CODE :** C000138355

**CLIENT'S NAME AND ADDRESS :**

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
 F-703, LADO SARAI, MEHRAULI  
 SOUTH WEST DELHI  
 NEW DELHI 110030  
 DELHI INDIA  
 8800465156

SRL LTD  
 Gate no 2, Residency Area, OPP. ST. Raphaels School,  
 INDORE, 452001  
 Madhya Pradesh, India  
 Tel : 0731 2490008

**PATIENT NAME :** MEGHNA CHAUREY (PKG10000292)

**PATIENT ID :** MEGHF210986290

**ACCESSION NO :** 0290WC005060 **AGE :** 36 Years **SEX :** Female

**ABHA NO :**

**DRAWN :** **RECEIVED :** 23/03/2023 09:16

**REPORTED :** 24/03/2023 11:40

**REFERRING DOCTOR :** DR. BANK OF BARODA

**CLIENT PATIENT ID :**

Test Report Status	Final	Results	Biological Reference Interval	Units
CHOLESTEROL LDL		76	Adult levels: Optimal < 100 Near optimal/above optimal: 100-129 Borderline high : 130-159 High : 160-189 Very high : = 190	mg/dL
NON HDL CHOLESTEROL		98	Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
METHOD : CALCULATED				
VERY LOW DENSITY LIPOPROTEIN		22.0		mg/dL
METHOD : CALCULATED				
CHOL/HDL RATIO		3.9		
LDL/HDL RATIO		2.2	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	





Patient Ref. No. 77500002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

## Interpretation(s)

- Cholesterol levels help assess the patient risk status and to follow the progress of patient under treatment to lower serum cholesterol concentrations.
- Serum Triglyceride (TG) are a type of fat and a major source of energy for the body. Both quantity and composition of the diet impact on plasma triglyceride concentrations. Elevations in TG levels are the result of overproduction and impaired clearance. High TG are associated with increased risk for CAD (Coronary artery disease) in patients with other risk factors, such as low HDL-C, some patient groups with elevated apolipoprotein B concentrations, and patients with forms of LDL that may be particularly atherogenic.
- HDL-C plays a crucial role in the initial step of reverse cholesterol transport, this considered to be the primary atheroprotective function of HDL
- LDL -C plays a key role in causing and influencing the progression of atherosclerosis and, in particular, coronary sclerosis. The majority of cholesterol stored in atherosclerotic plaques originates from LDL, thus LDL-C value is the most powerful clinical predictor.
- Non HDL cholesterol: Non-HDL-C measures the cholesterol content of all atherogenic lipoproteins, including LDL hence it is a better marker of risk in both primary and secondary prevention studies. Non-HDL-C also covers, to some extent, the excess ASCVD risk imparted by the sdLDL, which is significantly more atherogenic than the normal large buoyant particles, an elevated non-HDL-C indirectly suggests greater proportion of the small, dense variety of LDL particles

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

## Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category	
Extreme risk group	A. CAD with > 1 feature of high risk group B. CAD with > 1 feature of Very high risk group or recurrent ACS (within 1 year) despite LDL-C < or = 50 mg/dl or polyvascular disease
Very High Risk	1. Established ASCVD 2. Diabetes with 2 major risk factors or evidence of end organ damage 3. Familial Homozygous Hypercholesterolemia
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary Artery Calcium - CAC >300 AU. 7. Lipoprotein a >= 50mg/dl 8. Non stenotic carotid plaque
Moderate Risk	2 major ASCVD risk factors
Low Risk	0-1 major ASCVD risk factors
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors	
1. Age > or = 45 years in males and > or = 55 years in females	3. Current Cigarette smoking or tobacco use
2. Family history of premature ASCVD	4. High blood pressure
5. Low HDL	

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal <OR = 30 )	< 80 (Optional goal <OR = 60)	>OR = 50	>OR = 80



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Extreme Risk Group Category B	<OR = 30	<OR = 60	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

\*After an adequate non-pharmacological intervention for at least 3 months.

**References:** Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

## LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	0.44	0.0 - 1.2	mg/dL
METHOD : JENDRASSIK AND GROFF			
BILIRUBIN, DIRECT	0.18	0.0 - 0.2	mg/dL
METHOD : DIAZOTIZATION			
BILIRUBIN, INDIRECT	0.26	0.00 - 1.00	mg/dL
METHOD : CALCULATED			
TOTAL PROTEIN	7.6	6.4 - 8.3	g/dL
METHOD : BIURET			
ALBUMIN	4.4	3.50 - 5.20	g/dL
METHOD : BROMOCRESOL GREEN			
GLOBULIN	3.2	2.0 - 4.1	g/dL
METHOD : CALCULATED			
ALBUMIN/GLOBULIN RATIO	1.4	1.0 - 2.0	RATIO
METHOD : CALCULATED			
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	13	UPTO 32	U/L
METHOD : UV WITH P5P			
ALANINE AMINOTRANSFERASE (ALT/SGPT)	16	UPTO 34	U/L
METHOD : UV WITH P5P			
ALKALINE PHOSPHATASE	95	35 - 104	U/L
METHOD : PNPP			
GAMMA GLUTAMYL TRANSFERASE (GGT)	13	5 - 36	U/L
METHOD : G-GLUTAMYL-CARBOXY-NITROANILIDE			
LACTATE DEHYDROGENASE	188	135 - 214	U/L
METHOD : ENZYMATIC LACTATE - PYRUVATE(IFCC)			

## BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN	12	6 - 20	mg/dL
METHOD : UREASE KINETIC			

## CREATININE, SERUM



Scan to View Details



Scan to View Report

**DIAGNOSTIC REPORT**



**CLIENT CODE :** C000138355

**CLIENT'S NAME AND ADDRESS :**  
 ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
 F-703, LADO SARAI, MEHRAULI  
 SOUTH WEST DELHI  
 NEW DELHI 110030  
 DELHI INDIA  
 8800465156

SRL LTD  
 Gate no 2, Residency Area, OPP. ST. Raphaels School,  
 INDORE, 452001  
 Madhya Pradesh, India  
 Tel : 0731 2490008

**PATIENT NAME : MEGHNA CHAUREY (PKG10000292)** **PATIENT ID : MEGHF210986290**

ACCESSION NO : **0290WC005060** AGE : 36 Years SEX : Female ABHA NO :

DRAWN : RECEIVED : 23/03/2023 09:16 REPORTED : 24/03/2023 11:40

**REFERRING DOCTOR :** DR. BANK OF BARODA CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
CREATININE		0.59	0.50 - 0.90	mg/dL
METHOD : ALKALINE PICRATE KINETIC JAFFES				
<b>BUN/CREAT RATIO</b>				
BUN/CREAT RATIO		<b>20.34</b>	<b>High</b> 5.0 - 15.0	
METHOD : CALCULATED				
<b>URIC ACID, SERUM</b>				
URIC ACID		5.1	2.6 - 6.0	mg/dL
METHOD : URICASE/CATALASE UV				
<b>TOTAL PROTEIN, SERUM</b>				
TOTAL PROTEIN		7.6	6.4 - 8.3	g/dL
METHOD : BIURET				
<b>ALBUMIN, SERUM</b>				
ALBUMIN		4.4	3.5 - 5.2	g/dL
METHOD : BROMOCRESOL GREEN				
<b>GLOBULIN</b>				
GLOBULIN		3.2	2.0 - 4.1	g/dL
<b>ELECTROLYTES (NA/K/CL), SERUM</b>				
SODIUM, SERUM		143.8	136.0 - 146.0	mmol/L
METHOD : DIRECT ION SELECTIVE ELECTRODE				
POTASSIUM, SERUM		4.43	3.50 - 5.10	mmol/L
METHOD : DIRECT ION SELECTIVE ELECTRODE				
CHLORIDE, SERUM		<b>106.1</b>	<b>High</b> 98.0 - 106.0	mmol/L
METHOD : DIRECT ION SELECTIVE ELECTRODE				



Scan to View Details



Scan to View Report



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

## Interpretation(s)

Sodium	Potassium	Chloride
<b>Decreased in:</b> CCF,cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy,adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide,carbamazepine,anti depressants (SSRI), antipsychotics.	<b>Decreased in:</b> Low potassium intake,prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome,osmotic diuresis (e.g., hyperglycemia),alkalosis, familial periodic paralysis,trauma (transient).Drugs: Adrenergic agents, diuretics.	<b>Decreased in:</b> Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism,metabolic alkalosis. Drugs: chronic laxative,corticosteroids, diuretics.
<b>Increased in:</b> Dehydration (excessivesweating, severe vomiting or diarrhea),diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice,oral contraceptives.	<b>Increased in:</b> Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration,renal failure, Addison' s disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics,NSAIDs, beta-blockers, ACE inhibitors, high-dose trimethoprim-sulfamethoxazole.	<b>Increased in:</b> Renal failure, nephrotic syndrome, RTA,dehydration, overtreatment with saline,hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO <sub>3</sub> -), respiratory alkalosis,hyperadrenocorticism. Drugs: acetazolamide,androgens, hydrochlorothiazide,salicylates.
<b>Interferences:</b> Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	<b>Interferences:</b> Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	<b>Interferences:</b> Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

## PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW

APPEARANCE CLEAR

## CHEMICAL EXAMINATION, URINE

PH	6.0	4.7 - 7.5
SPECIFIC GRAVITY	1.010	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	<b>DETECTED (TRACE)</b>	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED

## MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS 2 - 3 NOT DETECTED /HPF



Scan to View Details



Scan to View Report





Patient Ref. No. 775000002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
PUS CELL (WBC'S)		2-3	0-5	/HPF
EPITHELIAL CELLS		3-5	0-5	/HPF
CASTS		NOT DETECTED		
CRYSTALS		NOT DETECTED		
BACTERIA		NOT DETECTED	NOT DETECTED	
YEAST		NOT DETECTED	NOT DETECTED	
REMARKS		Please note that all the urinary findings are confirmed manually as well.		

**Interpretation(s)**

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infection when present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

**THYROID PANEL, SERUM**

Scan to View Details



Scan to View Report





Patient Ref. No. 775000002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
T3		150.40	Non-Pregnant Women 80.0 - 200.0 Pregnant Women 1st Trimester:105.0 - 230.0 2nd Trimester:129.0 - 262.0 3rd Trimester:135.0 - 262.0	ng/dL
METHOD : CHEMILUMINESCENCE TECHNOLOGY				
T4		11.05	Non-Pregnant Women 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70	µg/dL
METHOD : CHEMILUMINESCENCE TECHNOLOGY				
TSH (ULTRASENSITIVE)		<b>0.115</b>	<b>Low</b> Non Pregnant Women 0.27 - 4.20 Pregnant Women 1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15	µIU/mL
METHOD : CHEMILUMINESCENCE TECHNOLOGY				





Patient Ref. No. 775000002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

## Interpretation(s)

**Triiodothyronine T3**, **Thyroxine T4**, and **Thyroid Stimulating Hormone TSH** are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low.

Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3) Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1) Subclinical Hypothyroidism (2) Patient with insufficient thyroid hormone replacement therapy (3) In cases of Autoimmune/Hashimoto thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical inflammation, drugs like amphetamines, Iodine containing drug and dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre (3) Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4 replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidelines of the American Thyroid association during pregnancy and Postpartum, 2011.

**NOTE: It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.** TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

## PAPANICOLAOU SMEAR

TEST METHOD

CONVENTIONAL GYNEC CYTOLOGY

SPECIMEN TYPE

TWO UNSTAINED CERVICAL SMEARS RECEIVED

REPORTING SYSTEM

2014 BETHESDA SYSTEM FOR REPORTING CERVICAL CYTOLOGY

SPECIMEN ADEQUACY

SATISFACTORY FOR EVALUATION WITH PRESENCE OF ENDOCERVICAL TRANSFORMATION ZONE COMPONENT.



Scan to View Details



Scan to View Report



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

## MICROSCOPY

- SQUAMOUS CELL ABNORMALITY

SMEARS SHOW CLUSTERS OF CELLS SHOWING NUCLEAR ENLARGEMENT, NUCLEAR OVERCROWDING AND IRREGULAR NUCLEAR MEMBRANE. FEW BINUCLEATED CELLS ARE ALSO SEEN.  
LOW GRADE SQUAMOUS INTRAEPITHELIAL LESION (LSIL).

## Comments

ADVISED HPV TESTING AND BIOPSY FOR CONFIRMATION.

\* THE REPORT RELATES ONLY TO THE SAMPLE SUBMITTED\*.

- PLEASE NOTE PAPANICOLAOU SMEAR STUDY IS A SCREENING PROCEDURE FOR CERVICAL CANCER WITH INHERENT FALSE NEGATIVE RESULTS, HENCE SHOULD BE INTERPRETED WITH CAUTION.
- NO CYTOLOGIC EVIDENCE OF HPV INFECTION IN THE SMEARS STUDIED.
- PRIMARY SCREENING AND REPORTING OF PAPANICOLAOU SMEARS IS CARRIED OUT BY SURGICAL PATHOLOGIST IN 100% OF CASES.

## PHYSICAL EXAMINATION,STOOL

COLOUR	BROWN	
CONSISTENCY	WELL FORMED	
MUCUS	ABSENT	NOT DETECTED
VISIBLE BLOOD	ABSENT	ABSENT
ADULT PARASITE	NOT DETECTED	

## CHEMICAL EXAMINATION,STOOL

OCCULT BLOOD	NOT DETECTED	NOT DETECTED
--------------	--------------	--------------

## MICROSCOPIC EXAMINATION,STOOL

PUS CELLS	2-3		/hpf
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
CYSTS	NOT DETECTED	NOT DETECTED	
OVA	NOT DETECTED		
LARVAE	NOT DETECTED	NOT DETECTED	
TROPHOZOITES	NOT DETECTED	NOT DETECTED	



Scan to View Details



Scan to View Report



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

## Interpretation(s)

Stool routine analysis is only a screening test for disorders of gastrointestinal tract like infection, malabsorption, etc. The following table describes the probable conditions, in which the analytes are present in stool.

PRESENCE OF	CONDITION
Pus cells	Pus in the stool is an indication of infection
Red Blood cells	Parasitic or bacterial infection or an inflammatory bowel condition such as ulcerative colitis
Parasites	Infection of the digestive system. Stool examination for ova and parasite detects presence of parasitic infestation of gastrointestinal tract. Various forms of parasite that can be detected include cyst, trophozoite and larvae. One negative result does not rule out the possibility of parasitic infestation. Intermittent shedding of parasites warrants examinations of multiple specimens tested on consecutive days. Stool specimens for parasitic examination should be collected before initiation of anti-diarrheal therapy or antiparasitic therapy. This test does not detect presence of opportunistic parasites like Cyclospora, Cryptosporidia and Isospora species. Examination of Ova and Parasite has been carried out by direct and concentration techniques.
Mucus	Mucus is a protective layer that lubricates, protects & reduces damage due to bacteria or viruses.
Charcot-Leyden crystal	Parasitic diseases.
Ova & cyst	Ova & cyst indicate parasitic infestation of intestine.
Frank blood	Bleeding in the rectum or colon.
Occult blood	Occult blood indicates upper GI bleeding.
Macrophages	Macrophages in stool are an indication of infection as they are protective cells.
Epithelial cells	Epithelial cells that normally line the body surface and internal organs show up in stool when there is inflammation or infection.
Fat	Increased fat in stool maybe seen in conditions like diarrhoea or malabsorption.
pH	Normal stool pH is slightly acidic to neutral. Breast-fed babies generally have an acidic stool.

## ADDITIONAL STOOL TESTS :

- Stool Culture**: - This test is done to find cause of GI infection, make decision about best treatment for GI infection & to find out if treatment for GI infection worked.
- Fecal Calprotectin**: It is a marker of intestinal inflammation. This test is done to differentiate Inflammatory Bowel Disease (IBD) from Irritable Bowel Syndrome (IBS).
- Fecal Occult Blood Test (FOBT)**: This test is done to screen for colon cancer & to evaluate possible cause of unexplained anaemia.
- Clostridium Difficile Toxin Assay**: This test is strongly recommended in healthcare associated bloody or watery diarrhoea, due to overuse of broad spectrum antibiotics which alter the normal GI flora.
- Biofire (Film Array) GI PANEL**: In patients of Diarrhoea, Dysentery, Rice watery Stool, FDA approved, Biofire Film Array Test, (Real Time Multiplex PCR) is strongly recommended as it identifies organisms, bacteria, fungi, virus, parasite and other opportunistic pathogens, Vibrio cholera infections only in 3 hours. Sensitivity 96% & Specificity 99%.
- Rota Virus Immunoassay**: This test is recommended in severe gastroenteritis in infants & children associated with watery diarrhoea, vomiting & abdominal cramps. Adults are also affected. It is highly contagious in nature.



Scan to View Details



Scan to View Report



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

## ABO GROUP &amp; RH TYPE, EDTA WHOLE BLOOD

ABO GROUP

TYPE B

METHOD : TUBE AGGLUTINATION

RH TYPE

POSITIVE

METHOD : TUBE AGGLUTINATION

## XRAY-CHEST

&gt;&gt;&gt;

BOTH THE LUNG FIELDS ARE CLEAR

&gt;&gt;&gt;

BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGLES ARE CLEAR

&gt;&gt;&gt;

BOTH THE HILA ARE NORMAL

&gt;&gt;&gt;

CARDIAC AND AORTIC SHADOWS APPEAR NORMAL

&gt;&gt;&gt;

BOTH THE DOMES OF THE DIAPHRAM ARE NORMAL

&gt;&gt;&gt;

VISUALIZED BONY THORAX IS NORMAL

IMPRESSION

NO ABNORMALITY DETECTED

Dr G S Saluja MBBS , DMRD  
(Consultant Radiologist)X

## TMT OR ECHO

TMT OR ECHO

2D ECHO DONE

IMPRESSION :

-MILD CONCENTRIC LVH

-TRIVILA AR

-NO RWMA AT REST

LVEF 62%

NORMAL PAP

## ECG

ECG

SINUS RHYTHM

NORMAL ECG

## MEDICAL HISTORY

RELEVANT PRESENT HISTORY

NOT SIGNIFICANT

RELEVANT PAST HISTORY

HYPOTHYROID - 6-7 YEARS

HTN - 7-8 MONTHS

LSC - 2016

RELEVANT PERSONAL HISTORY

NOT SIGNIFICANT

RELEVANT FAMILY HISTORY

MOTHER :- DM , HYPOTHYROID

FATHER :- HTN

OCCUPATIONAL HISTORY

NOT SIGNIFICANT

HISTORY OF MEDICATIONS

NOT SIGNIFICANT

## ANTHROPOMETRIC DATA &amp; BMI

HEIGHT IN METERS

1.54

mts



Scan to View Details



Scan to View Report



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

WEIGHT IN KGS.	97			Kgs
BMI	41			
			BMI & Weight Status as follows: kg/sqmts Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese	

## GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE	NORMAL
PHYSICAL ATTITUDE	NORMAL
GENERAL APPEARANCE / NUTRITIONAL STATUS	OBESE
BUILT / SKELETAL FRAMEWORK	AVERAGE
FACIAL APPEARANCE	NORMAL
SKIN	NORMAL
UPPER LIMB	NORMAL
LOWER LIMB	NORMAL
NECK	NORMAL
NECK LYMPHATICS / SALIVARY GLANDS	NOT ENLARGED OR TENDER
THYROID GLAND	NOT ENLARGED
CAROTID PULSATION	NORMAL
TEMPERATURE	AFEBRILE
PULSE	78/MIN, REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID BRUIT
RESPIRATORY RATE	NORMAL

## CARDIOVASCULAR SYSTEM

BP	150/100 MM HG (SUPINE)	mm/Hg
PERICARDIUM	NORMAL	
APEX BEAT	NORMAL	
HEART SOUNDS	NORMAL	
MURMURS	ABSENT	

## RESPIRATORY SYSTEM

SIZE AND SHAPE OF CHEST	NORMAL
MOVEMENTS OF CHEST	SYMMETRICAL
BREATH SOUNDS INTENSITY	NORMAL
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)
ADDED SOUNDS	ABSENT

## PER ABDOMEN



Scan to View Details



Scan to View Report



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

APPEARANCE	NORMAL
VENOUS PROMINENCE	ABSENT
LIVER	NOT PALPABLE
SPLEEN	NOT PALPABLE
HERNIA	ABSENT
<b>CENTRAL NERVOUS SYSTEM</b>	
HIGHER FUNCTIONS	NORMAL
CRANIAL NERVES	NORMAL
CEREBELLAR FUNCTIONS	NORMAL
SENSORY SYSTEM	NORMAL
MOTOR SYSTEM	NORMAL
REFLEXES	NORMAL
<b>MUSCULOSKELETAL SYSTEM</b>	
SPINE	NORMAL
JOINTS	NORMAL
<b>BASIC EYE EXAMINATION</b>	
CONJUNCTIVA	NORMAL
EYELIDS	NORMAL
EYE MOVEMENTS	NORMAL
CORNEA	NORMAL
DISTANT VISION RIGHT EYE WITHOUT GLASSES	6/6, WITHIN NORMAL LIMIT
DISTANT VISION LEFT EYE WITHOUT GLASSES	6/6, WITHIN NORMAL LIMIT
NEAR VISION RIGHT EYE WITHOUT GLASSES	N6, WITHIN NORMAL LIMIT
NEAR VISION LEFT EYE WITHOUT GLASSES	N6, WITHIN NORMAL LIMIT
COLOUR VISION	NORMAL
<b>BASIC ENT EXAMINATION</b>	
EXTERNAL EAR CANAL	NORMAL
TYMPANIC MEMBRANE	NORMAL
NOSE	NO ABNORMALITY DETECTED
SINUSES	NORMAL
THROAT	NORMAL
TONSILS	NOT ENLARGED
<b>SUMMARY</b>	
RELEVANT HISTORY	NOT SIGNIFICANT



Scan to View Details



Scan to View Report





Patient Ref. No. 775000002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN : RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

RELEVANT GP EXAMINATION FINDINGS

OBESE

REMARKS / RECOMMENDATIONS

NONE

## FITNESS STATUS

FITNESS STATUS

FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

## Comments

CLINICAL FINDINGS :-

OBESE WEIGHT STATUS.

FITNESS STATUS :-

FITNESS STATUS : FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

ADVICE : WEIGHT REDUCTION, LOW FAT&amp; CARBOHYDRATE DIET AND REGULAR PHYSICAL EXERCISE FOR OBESE WEIGHT STATUS

NEED PHYSICIAN CONSULTATION FOR LIFE STYLE MODIFICATION.

## Interpretation(s)

BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

ERYTHROCYTE SEDIMENTATION RATE (ESR),WHOLE BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

## TEST INTERPRETATION

**Increase** in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm/hr(95 if anemic). ESR returns to normal 4th week post partum.

**Decreased** in: Polycythemia vera, Sickle cell anemia

## LIMITATIONS

**False elevated** ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

**False Decreased** : Poikilocytosis,(SickleCells,spherocytes),Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine,

salicylates)

REFERENCE :



Scan to View Details



Scan to View Report



CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN : RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

## GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and so that no glucose is excreted in the urine.

**Increased in:** Diabetes mellitus, Cushing's syndrome (10 - 15%), chronic pancreatitis (30%). Drugs: corticosteroids, phenytoin, estrogen, thiazides.

**Decreased in:** Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical, stomach, fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases (e.g. galactosemia), Drugs-insulin, ethanol, propranolol, sulfonyleureas, tolbutamide, and other oral hypoglycemic agents.

**NOTE:** While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin (HbA1c) levels are favored to monitor glycaemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLYCOSYLATED HEMOGLOBIN (HbA1C), EDTA WHOLE BLOOD-Used For:

- Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patient's metabolic control has remained continuously within the target range.

- eAG (Estimated average glucose) converts percentage HbA1c to mg/dl, to compare blood glucose levels.
- eAG gives an evaluation of blood glucose levels for the last couple of months.
- eAG is calculated as  $eAG (mg/dl) = 28.7 * HbA1c - 46.7$

## HbA1c Estimation can get affected due to :

- Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.
- Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin).
- Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.
- Interference of hemoglobinopathies in HbA1c estimation is seen in

a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.

b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c) HbF > 25% on alternate platform (Boronate affinity chromatography) is recommended for testing of HbA1c. Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

**Bilirubin** is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. **Elevated levels** results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease. Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

**AST** is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

**ALP** is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease.

**GGT** is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

**Total Protein** also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

**Albumin** is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-**Causes of Increased** levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol,



Scan to View Details



Scan to View Report



**CLIENT CODE :** C000138355

**CLIENT'S NAME AND ADDRESS :**  
 ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
 F-703, LADO SARAI, MEHRAULI  
 SOUTH WEST DELHI  
 NEW DELHI 110030  
 DELHI INDIA  
 8800465156

SRL LTD  
 Gate no 2, Residency Area, OPP. ST. Raphaels School,  
 INDORE, 452001  
 Madhya Pradesh, India  
 Tel : 0731 2490008

**PATIENT NAME : MEGHNA CHAUREY (PKG10000292)** PATIENT ID : **MEGHF210986290**

ACCESSION NO : **0290WC005060** AGE : 36 Years SEX : Female ABHA NO :

DRAWN : RECEIVED : 23/03/2023 09:16 REPORTED : 24/03/2023 11:40

**REFERRING DOCTOR : DR. BANK OF BARODA** CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

**Causes of decreased level include Liver disease, SIADH.**

CREATININE, SERUM- **Higher than normal level may be due to:**

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

**Lower than normal level may be due to:**

• Myasthenia Gravis, Muscuophy

URIC ACID, SERUM- **Causes of Increased levels:-** Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome **Causes of decreased levels-** Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM- is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.

**Higher-than-normal levels may be due to:** Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.

**Lower-than-normal levels may be due to:** Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. **Low blood albumin levels (hypoalbuminemia) can be caused by:** Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD- Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A, B, O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.

MEDICAL

HISTORY-\*\*\*\*\* THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

\*\*\*\*\*

**FITNESS STATUS-** Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history as well as the comprehensiveness of the diagnostic panel which has been requested for. These are then further correlated with details of the job under consideration to eventually fit the right man to the right job.

Basis the above, SRL classifies a candidate's Fitness Status into one of the following categories:

- Fit (As per requested panel of tests) - SRL Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.
- Fit (with medical advice) (As per requested panel of tests) - This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician's consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.
- Fitness on Hold (Temporary Unfit) (As per requested panel of tests) - Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.
- Unfit (As per requested panel of tests) - An unfit report by SRL Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.





CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD  
Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Units
--------------------	-------	---------	-------

**MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE****ULTRASOUND ABDOMEN****ULTRASOUND ABDOMEN**

..

**Comments**

U.S.G OF WHOLE ABDOMEN

Liver is normal in size, shape with with smooth outline. Parenchymal echotexture is homogeneous. Intra & Extra hepatic biliary radicals are normal. Portal vein and C.B.D are normal in caliber.

Gall Bladder is normal, thin walled & its lumen is echo free.

Spleen is normal in size, shape & echotexture.

Pancreas is normal in size, shape & echotexture.

Both Kidneys are normal in size, shape and echotexture. Central pelvicalyceal system is normal. Corticomedullary differentiation is maintained.

IVC and AO is normal in caliber.No lymphadenopathy.

Urinary Bladder is normal thin walled,there is no calculus.

Uterus is anteverted and normal in size. Myometrial echotexture is homogeneous Endometrial echo reflection is normal. Cervix and endocervical canal appears normal.

Bilateral Ovaries are appears normal.

IMPRESSION- No Significant abnormality seen in USG of Whole Abdomen.

Dr G S Saluja MBBS, DMRD  
(Consultant Radiologist)

**\*\*End Of Report\*\***

Please visit [www.srlworld.com](http://www.srlworld.com) for related Test Information for this accession

**Dr.Arpita Pasari, MD**  
Consultant Pathologist

**Dr.Meena Jinwah ,MBBS . MD**  
Consultant Microbiologist



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002692211

CLIENT CODE : C000138355

## CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )  
F-703, LADO SARAI, MEHRAULI  
SOUTH WEST DELHI  
NEW DELHI 110030  
DELHI INDIA  
8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,  
INDORE, 452001  
Madhya Pradesh, India  
Tel : 0731 2490008

PATIENT NAME : MEGHNA CHAUREY (PKG10000292)

PATIENT ID : MEGHF210986290

ACCESSION NO : 0290WC005060 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 23/03/2023 09:16

REPORTED : 24/03/2023 11:40

REFERRING DOCTOR : DR. BANK OF BARODA

CLIENT PATIENT ID :

Test Report Status	Final	Results	Units
--------------------	-------	---------	-------

## CONDITIONS OF LABORATORY TESTING &amp; REPORTING

1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
4. A requested test might not be performed if:
  - i. Specimen received is insufficient or inappropriate
  - ii. Specimen quality is unsatisfactory
  - iii. Incorrect specimen type
  - iv. Discrepancy between identification on specimen container label and test requisition form
5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
8. Test results cannot be used for Medico legal purposes.
9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

## SRL Limited

Fortis Hospital, Sector 62, Phase VIII,  
Mohali 160062



Scan to View Details



Scan to View Report