

Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 28, 2023, 08:39 a.m. **Reporting Time :** Jan 29, 2023, 09:25 p.m.

Sample ID:

			000700010	
Test Description	Value(s)	Reference Range	Unit	
CBC; Complete Blood Count				
Hemoglobin (Hb)* Method : Cynmeth Photometric Measurement	16.7	13.5 - 18.0	gm/dL	
Erythrocyte (RBC) Count* Method : Electrical Impedence	6.44	4.7 - 6.0	mil/cu.mm	
Packed Cell Volume (PCV)* Method : Calculated	59.2	42 - 52	%	
Mean Cell Volume (MCV)* Method : Electrical Impedence	92	78 - 100	fL	
Mean Cell Haemoglobin (MCH)* Method : Calculated	25.9	27 - 31	pg	
Mean Corpuscular Hb Concn. (MCHC)* Method : Calculated	28.2	32 - 36	gm/dL	
Red Cell Distribution Width (RDW)* Method : Electrical Impedence	12.5	11.5 - 14.0	%	
Total Leucocytes (WBC) Count* Method : Electrical Impedence	8500	4000-10000	cell/cu.mm	
Neutrophils* Method : VCSn Technology	73	40 - 80	%	
Lymphocytes* Method : VCSn Technology	18	20 - 40	%	
Monocytes* Method : VCSn Technology	8	2 - 10	%	
Eosinophils* Method : VCSn Technology	1	1 - 6	%	
Basophils	0	0 - 1		
Platelet Count* Method : Electrical Impedence	2.7	1.5 - 4.5	10^3/ul	
Mean Platelet Volume (MPV)* Method : Electrical Impedence	8.3	7.2 - 11.7	fL	

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Odepter

Approved by





Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 28, 2023, 08:39 a.m. **Reporting Time :** Jan 29, 2023, 09:25 p.m.

Sample ID :

668766943

Test Description	Value(s)	Reference Range	Unit
PCT*	0.225	0.2 - 0.5	%
Method : Calculated			
PDW*	16.2	9.0 - 17.0	%
Method : Calculated			

Tests done on Automated Three Part Cell Counter. (WBC, RBC, Platelet count by impedance method, colorimetric method for Hemoglobin, WBC differential by flow cytometry using laser technology other parameters are calculated). All Abnormal Haemograms are reviewed confirmed microscopically.

Esr, Erythrocyte Sedimentation Rate

Esr, Erythrocyte Sedimentation Rate (Westergren)

20

0-10

mm/hr

Interpretation:

- It indicates presence and intensity of an inflammatory process. It does not diagnose a specific disease. Changes in the ESR are more significant than the abnormal results of a single test.
- It is a prognostic test and used to monitor the course or response to treatment of diseases like tuberculosis, bacterial endocarditis, acute rheumatic fever, rheumatoid arthritis, SLE, Hodgkins disease, temporal arteritis and polymyalgia rheumatica.
- It is also increased in pregnancy, multiple myeloma, menstruation, and hypothyroidism.

Urine Routine

Colour*	Pale Yellow		
Transparency (Appearance)*	Clear	Clear	
Reaction (pH)*	5.0	4.5 - 8	
Specific Gravity*	1.020	1.010 - 1.030	

Chemical Examination (Automated Dipstick Method) Urine

Urine Glucose*	Negative	Negative
Urine Protein*	Negative	Negative
Urine Ketone*	Negative	Negative
Blood*	Negative	Negative

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by

Scan to Validate



— Flower us —



Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time: Jan 28, 2023, 08:39 a.m. **Reporting Time:** Jan 29, 2023, 09:25 p.m.

Sample ID:

668766943

Test Description	Value(s)	Reference Range	Unit
Bilirubin*	Negative	Negative	
Nitrite*	Negative	Negative	
_eucocytes*	Negative	Negative	
Jrobilinogen*	Normal	Normal	
Ilicroscopic Examination Urine			
Pus Cells (WBCs)*	1-2	0 - 5	/hpf
pithelial Cells*	0-1	0 - 4	/hpf
Red blood Cells*	Absent	Absent	/hpf
Crystals*	Absent	Absent	
Cast*	Absent	Absent	
Bacteria*	Absent	Absent	

Blood Group & Rh Type

Blood Grouping & Rh Typing

Method : Forward and Reverse By Tube Method

"AB" POSITIVE (+VE)

Methodology

This is done by forward and reverse grouping by tube Agglutination method.

Interpretation

Newborn baby does not produce ABO antibodies until 3 to 6 months of age. So the blood group of the Newborn baby is done by ABO antigen grouping (forward grouping) only, antibody grouping (reverse grouping) is not required. Confirmation of the New-born's blood group is indicated when the A and B antigen expression and the isoagglutinins are fully developed (2–4 years).

Fasting - Glucose

Dr.CH.Deepthi Chandrika M.D. Pathology

Reg.No.APCM/FMR/77174

Approved by

Scan to Validate



— Flower us — ⑤ f ⑥ ⑥



Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time: Jan 28, 2023, 08:39 a.m. **Reporting Time:** Jan 29, 2023, 09:25 p.m.

Sample ID :

Test Description	Value(s)	Reference Range	Unit
Glucose Fasting*	116	Normal: 70-100	mg/dL
Method : Plasma, Hexokinase		Impaired Fasting Glucose (IFG):	9, 5.=
		100-125	
		Diabetes Mellitus: >= 126	
		(On more than one occasion)	
		(American Diabetes Association	
		guidelines 2017)	
Post Prandial Blood Sugar			
Blood Glucose-Post Prandial*	174	80-140	mg/dL
Method : Plasma - P, Hexokinase			Ü
Fasting Urine Sugar			
Fasting Urine Sugar	116	NEGATIVE -	
Post Prandial Urine Sugar			
Post Prandial Urine Sugar	174		
HBA1C (Glycosylated Haemoglobin)			
Glyco Hb (HbA1C)	6.1	Non-Diabetic: <=5.9	%
Method : EDTA Whole blood,HPLC		Pre Diabetic:6.0-6.4	
		Diabetic: >=6.5	
Estimated Average Glucose :	128		mg/dL
Interpretations			

- 1. HbA1C has been endorsed by clinical groups and American Diabetes Association guidelines 2017 for diagnosing diabetes using a cut off point of 6.5%
- 2. Low glycated haemoglobin in a non diabetic individual are often associated with systemic inflammatory diseases, chronic anaemia (especially severe iron deficiency and haemolytic), chronic renal failure and liver diseases. Clinical correlation suggested.

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Reg.No.APCIVI/FIVIK///I

Approved by

Scan to Validate



— Flower us —



Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time: Jan 28, 2023, 08:39 a.m. **Reporting Time:** Jan 29, 2023, 09:25 p.m.

Sample ID:

668766943

Test Description Value(s) Reference Range Unit

3. In known diabetic patients, following values can be considered as a tool for monitoring the glycemic control.

Excellent control-6-7 %
Fair to Good control – 7-8 %
Unsatisfactory control – 8 to 10 %

Poor Control – More than 10 %

Thyroid Function Test (TFT)			
TRI-IODO THYRONINE (T3) Method : CLIA	105	60 - 181	ng/dL
TOTAL THYROXINE (T4) Method : CLIA	11.4	4.2 - 12.0	ug/dL
THYROID STIMULATING HORMONE (TSH) Method: CLIA	1.10	0.46 – 8.10 : 1 Yrs – 5 Yrs 0.36 – 5.80 : 6 Yrs – 18 Yrs 0.35 – 5.50 : >18 Yrs Pregnancy Ranges 1st Trimester :0.1 - 2.5 2nd Trimester :0.2 - 3.0 3rd Trimester:0.3 - 3.0	uIU/mL

Comments:

IF NOT ON DRUGS SUGGESTED FT3 & FT4 ESTIMATION

Please correlate with clinical conditions.

Note: Serum T3, T4 and TSH form the three components of thyroid screening panel, useful in diagnosing various disorders of the thyroid gland. Primary Hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels. Although elevated TSH levels are nearly always indicative of Primary Hypothyroidism, rarely they can from TSH secreting pituitary tumors (Secondary hyperthyroidism)To confirm diagnosis - evaluate FT3 and FT4.

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by





Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 28, 2023, 08:39 a.m. **Reporting Time :** Jan 29, 2023, 09:25 p.m.

Sample ID:

668766943

Test Description	Value(s)	Reference Range	Unit
<u>Lipid Profile</u>			
Cholesterol-Total Method : Serum, Cholesterol oxidase esterase, peroxidase	166	Desirable: <= 200 Borderline High: 201-239 High: > 239 Ref: The National Cholesterol Education Program (NCEP) Adult Treatment Panel III Report.	mg/dL
Triglycerides Method : Serum, Enzymatic, endpoint	164	Normal: < 150 Borderline High: 150-199 High: 200-499 Very High: >= 500	mg/dL
Cholesterol-HDL Direct Method : Serum, Direct measure-PEG	40	<40: Low 40 - 60: Optimal > 60: Desirable	mg/dL
LDL Cholesterol Method : Serum	93.2	Optimal: < 100 Near optimal/above optimal: 100-129 Borderline high: 130-159 High: 160-189 Very High: >= 190	mg/dL
Non - HDL Cholesterol, Serum Method : calculated	126	Desirable: < 130 mg/dL Borderline High: 130-159mg/dL High: 160-189 mg/dL Very High: > or = 190 mg/dL	mg/dL
VLDL Cholesterol Method : calculated	32.8	6 - 38	mg/dL
CHOL/HDL RATIO Method : calculated	4.1	3.5 - 5.0	ratio

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by





Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time : Jan 28, 2023, 08:39 a.m. **Reporting Time :** Jan 29, 2023, 09:25 p.m.

Sample ID:

668766943

		000700943	
Test Description	Value(s)	Reference Range	Unit
LDL/HDL RATIO	2.3	Desirable / low risk - 0.5 -3.0	ratio
Method : calculated		Low/ Moderate risk - 3.0- 6.0	
		Elevated / High risk - > 6.0	
Note: 8-10 hours fasting sample is required.			
KIDNEY FUNCTION TEST			
Urea *	18	15- 50	mg/dL
Method : Serum			
Blood Urea Nitrogen-BUN*	8.41	7 - 24	mg/dL
Method : Serum, Urease			
Uric Acid*	5.9	3.5 - 7.2	mg/dL
Method : Serum, Uricase/POD			
Creatinine*	0.7	0.7 - 1.3	mg/dL
Method : Serum, Jaffe IDMS			
Liver Funtion Test (LFT) with GGT			
Bilirubin - Total	1.4	0.3 - 1.2	mg/dL
Method : Serum, Jendrassik Grof			
Bilirubin - Direct	0.2	Adults and Children: < 0.2	mg/dL
Method : Serum, Diazotization			
Bilirubin - Indirect	1.20	0.1 - 1.0	mg/dL
Method : Serum, Calculated			
SGOT	46	< 50	U/L
Method : Serum, UV with P5P, IFCC 37 degree			
SGPT	36	< 50	U/L
Method : Serum, UV with P5P, IFCC 37 degree	4.00	0 = 4.4	
SGOT/SGPT	1.28	0.7 - 1.4	ratio
Method : calculated		F.F.	11/1
GGT-Gamma Glutamyl Transpeptidase	-	< 55	U/L
Method : Serum, G-glutamyl-carboxy-nitoanilide			

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Obsepter

Approved by





Age / Gender: 49 years / Male

Patient ID: 16198

Source: MEDI WHEEL

Referral: SELF

Collection Time: Jan 28, 2023, 08:39 a.m. **Reporting Time:** Jan 29, 2023, 09:25 p.m.

Sample ID:

668766943

Test Description	Value(s)	Reference Range	Unit
Alkaline Phosphatase-ALPI	160	30-120	U/L
Method : Serum, PNPP, AMP Buffer, IFCC 37 degree			
Total Protein	6.9	6.6 - 8.3	g/dL
Method : Serum, Biuret, reagent blank end point			
Albumin	3.8	Adults: 3.5 - 5.2	g/dL
Method : Serum, Bromcresol purple			
Globulin	3.10	1.8 - 3.6	g/dL
Method : Calculated			
A/G Ratio	1.23	1.2 - 2.2	ratio
Method : Calculated			

END OF REPORT

Dr.CH.Deepthi Chandrika M.D. Pathology Reg.No.APCM/FMR/77174

Approved by

