



Units

CLIENT CODE: C000138361
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

**Test Report Status** 

SRL Ltd

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

**Biological Reference Interval** 

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email : wellness.eastdelhi@srl.in

PATIENT NAME: ANKIT SINGH PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

<u>Final</u>

DRAWN: RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48

**Results** 

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

MEDI WHEEL FULL BODY HEALTH CHECK L	IP BELOW 40 MALE		
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN	15.6	13.0 - 17.0	g/dL
METHOD: SPECTROPHOTOMETRY			-
RED BLOOD CELL COUNT	5.11	4.5 - 5.5	mil/µL
METHOD: ELECTRICAL IMPEDANCE			
WHITE BLOOD CELL COUNT	6.30	4.0 - 10.0	thou/µL
METHOD: ELECTRICAL IMPEDANCE			
PLATELET COUNT	223	150 - 410	thou/µL
METHOD: ELECTRICAL IMPEDANCE			
RBC AND PLATELET INDICES			
HEMATOCRIT	46.2	40.0 - 50.0	%
METHOD: CALCULATED PARAMETER			
MEAN CORPUSCULAR VOL	90.4	83.0 - 101.0	fL
METHOD : DERIVED/COULTER PRINCIPLE			
MEAN CORPUSCULAR HGB.	30.5	27.0 - 32.0	pg
METHOD: CALCULATED PARAMETER			
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION METHOD: CALCULATED PARAMETER	33.8	31.5 - 34.5	g/dL
MENTZER INDEX	17.7		
RED CELL DISTRIBUTION WIDTH	14.0	11.6 - 14.0	%
METHOD : DERIVED/COULTER PRINCIPLE			
MEAN PLATELET VOLUME	10.2	6.8 - 10.9	fL
METHOD : DERIVED/COULTER PRINCIPLE			
WBC DIFFERENTIAL COUNT - NLR			
SEGMENTED NEUTROPHILS	57	40 - 80	%
METHOD: VCS TECHNOLOGY/ MICROSCOPY			
ABSOLUTE NEUTROPHIL COUNT	3.60	2.0 - 7.0	thou/µL
METHOD: CALCULATED PARAMETER			
LYMPHOCYTES	31	20 - 40	%
METHOD: VCS TECHNOLOGY/ MICROSCOPY			
ABSOLUTE LYMPHOCYTE COUNT METHOD: CALCULATED PARAMETER	1.90	1.0 - 3.0	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	1.9		
EOSINOPHILS	4	1.0 - 6.0	%
METHOD: VCS TECHNOLOGY/ MICROSCOPY	•	1.0 0.0	70









**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030 DELHI INDIA 8800465156

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

**PATIENT NAME: ANKIT SINGH** PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results		Biological Reference Inter	val Units
ABSOLUTE EOSINOPHI	L COUNT	0.25		0.02 - 0.50	thou/µL
METHOD : CALCULATED PAR		0.20		0.02	εο α, μ=
MONOCYTES		8		2.0 - 10.0	%
METHOD: VCS TECHNOLOG					
ABSOLUTE MONOCYTE  METHOD: CALCULATED PAR		0.50		0.2 - 1.0	thou/μL
BASOPHILS		0		0 - 1	%
METHOD: VCS TECHNOLOG	SY/ MICROSCOPY				
ABSOLUTE BASOPHIL (	COUNT	0.00	Low	0.02 - 0.10	thou/µL
METHOD : CALCULATED PAR	RAMETER				
DIFFERENTIAL COUNT	PERFORMED ON:	EDTA SMEAR			
<b>ERYTHRO SEDIMENT</b>	ATION RATE, BLOOD				
SEDIMENTATION RATE	(ESR)	14		0 - 14	mm at 1 hr
METHOD: WESTERGREN ME	THOD				
GLUCOSE, FASTING,	PLASMA				
GLUCOSE, FASTING, PI	LASMA	93		74 - 106	mg/dL
METHOD: HEXOKINASE					
<b>GLYCOSYLATED HEM</b>	OGLOBIN, EDTA WHO	LE BLOOD			
GLYCOSYLATED HEMO	GLOBIN (HBA1C)	5.6		Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 ADA Target: 7.0 Action suggested: > 8.0	%
MEAN PLASMA GLUCOS METHOD : CALCULATED PAR		114.0		< 116.0	mg/dL
GLUCOSE, POST-PRA	NDIAL, PLASMA				
GLUCOSE, POST-PRANI	DIAL, PLASMA	101		70 - 140	mg/dL
METHOD: HEXOKINASE					
CORONARY RISK PRO	OFILE (LIPID PROFILE	E), SERUM			
CHOLESTEROL	VIDAGE FOR THE STATE OF THE STA	254	High	< 200 Desirable 200 - 239 Borderline High >/= 240 High	mg/dL
	XIDASE, ESTERASE,PEROXIDASE			. 450 N	,
TRIGLYCERIDES		296	High	< 150 Normal 150 - 199 Borderline High 200 - 499 High >/= 500 Very High	mg/dL
METHOD : ENZYMATIC, END	POINT				,
HDL CHOLESTEROL		40		< 40 Low >/=60 High	mg/dL





Scan to View Report





**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030

DELHI INDIA 8800465156

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

**PATIENT NAME: ANKIT SINGH** PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Biological Reference Interv	al Units
METHOD : DIRECT MEASURE POLYMER-POLYANION				
DIRECT LDL CHOLESTEROL	183	Hiah	< 100 Optimal	mg/dL
DIRECT EDE CHOLESTEROL	103		100 - 129 Near or above optim 130 - 160 Borderline High 161 - 189 High >/= 190 Very High	
METHOD : DIRECT MEASURE			, , ,	
NON HDL CHOLESTEROL	214	High	Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
METHOD: CALCULATED PARAMETER				
CHOL/HDL RATIO	6.4	High	3.3-4.4 Low Risk 4.5-7.0 Average Risk 7.1-11.0 Moderate Risk > 11.0 High Risk	
METHOD : CALCULATED PARAMETER				
LDL/HDL RATIO	4.6	High	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	
METHOD : CALCULATED PARAMETER				
VERY LOW DENSITY LIPOPROTEIN	59.2	High	Desirable value : 10 - 35	mg/dL
METHOD : CALCULATED PARAMETER				
LIVER FUNCTION PROFILE, SERUM				
BILIRUBIN, TOTAL	1.01		UPTO 1.2	mg/dL
METHOD : DIAZONIUM ION, BLANKED (ROCHE)				
BILIRUBIN, DIRECT	0.26		0.00 - 0.30	mg/dL
METHOD: DIAZOTIZATION	0.75	Himb	0.00	
BILIRUBIN, INDIRECT	0.75	High	0.00 - 0.60	mg/dL
METHOD : CALCULATED PARAMETER TOTAL PROTEIN	7.8		6.6 - 8.7	م اطا
METHOD: BIURET, SERUM BLANK, ENDPOINT	7.0		0.0 - 8.7	g/dL
ALBUMIN	4.9		3.97 - 4.94	g/dL
METHOD : BROMOCRESOL GREEN	4.5		3.37 4.34	g/ uL
GLOBULIN	2.9		2.0 - 4.0 Neonates - Pre Mature: 0.29 - 1.04	g/dL
METHOD: CALCULATED PARAMETER	4.7		10.20	DATIO
ALBUMIN/GLOBULIN RATIO  METHOD: CALCULATED PARAMETER	1.7		1.0 - 2.0	RATIO





Scan to View Report





**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030 DELHI INDIA 8800465156

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

**PATIENT NAME: ANKIT SINGH** PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Biological Reference Into	erval Units
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	38		0 - 40	U/L
METHOD: UV WITHOUT P5P				
ALANINE AMINOTRANSFERASE (ALT/SGPT)	79	High	0 - 41	U/L
METHOD: UV WITHOUT P5P				
ALKALINE PHOSPHATASE	71		40 - 129	U/L
METHOD: PNPP, AMP BUFFER-IFCC				
GAMMA GLUTAMYL TRANSFERASE (GGT)	93	High	8 - 61	U/L
METHOD : G-GLUTAMYL-CARBOXY-NITROANILIDE-IFCC				
LACTATE DEHYDROGENASE	225		135 - 225	U/L
METHOD : L TO P, IFCC				
SERUM BLOOD UREA NITROGEN				
BLOOD UREA NITROGEN	14		6 - 20	mg/dL
METHOD : UREASE - UV				
CREATININE, SERUM				
CREATININE	0.88		0.70 - 1.20	mg/dL
METHOD : ALKALINE PICRATE-KINETIC				
BUN/CREAT RATIO				
BUN/CREAT RATIO	15.91	High	5.00 - 15.00	
METHOD : CALCULATED PARAMETER				
URIC ACID, SERUM				
URIC ACID	5.4		3.4 - 7.0	mg/dL
METHOD : URICASE, COLORIMETRIC				
TOTAL PROTEIN, SERUM				
TOTAL PROTEIN	7.8		6.6 - 8.7	g/dL
METHOD: BIURET, SERUM BLANK, ENDPOINT				
ALBUMIN, SERUM				
ALBUMIN	4.9		3.97 - 4.94	g/dL
METHOD: BROMOCRESOL GREEN				
GLOBULIN				
GLOBULIN	2.9		2.0 - 4.0 Neonates - Pre Mature: 0.29 - 1.04	g/dL
METHOD: CALCULATED PARAMETER				
ELECTROLYTES (NA/K/CL), SERUM				
SODIUM	138		136 - 145	mmol/L
METHOD: ISE INDIRECT				
POTASSIUM	4.20		3.5 - 5.1	mmol/L









**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030

DELHI INDIA 8800465156

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

**PATIENT NAME: ANKIT SINGH** PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
METHOD: ISE INDIRECT			
CHLORIDE	101	98 - 107	mmol/L
METHOD: ISE INDIRECT			
URINALYSIS			
COLOR	PALE YELLOW		
METHOD : VISUAL			
APPEARANCE	CLEAR		
METHOD : VISUAL			
PH	6.0	4.7 - 7.5	
METHOD: DOUBLE INDICATOR PRINCIPLE			
SPECIFIC GRAVITY	1.025	1.003 - 1.035	
METHOD: PKA CHANGE OF PRETREATED POLYELECTROLYTES			
GLUCOSE	NOT DETECTED	NOT DETECTED	
METHOD: OXIDASE-PEROXIDASE REACTION			
PROTEIN	NOT DETECTED	NOT DETECTED	
METHOD: PROTEIN- ERROR INDICATOR			
KETONES	NOT DETECTED	NOT DETECTED	
METHOD: ACETOACETIC REACTION WITH NITROPRUSSIDE			
BLOOD	NOT DETECTED	NOT DETECTED	
METHOD: PEROXIDASE-LIKE ACTIVITY OF HEMOGLOBIN			
BILIRUBIN	NOT DETECTED	NOT DETECTED	
METHOD: DIAZOTIZATION			
UROBILINOGEN	NORMAL	NORMAL	
METHOD: MODIFIED EHRLICH REACTION			
NITRITE	NOT DETECTED	NOT DETECTED	
METHOD: CONVERTION OF NITRATE TO NITRITE			
PUS CELL (WBC'S)	1-2	0-5	/HPF
METHOD: MICROSCOPIC EXAMINATION			
EPITHELIAL CELLS	0-1	0-5	/HPF
METHOD: MICROSCOPIC EXAMINATION			
ERYTHROCYTES (RBC'S)	NOT DETECTED	NOT DETECTED	/HPF
METHOD: MICROSCOPIC EXAMINATION			
CASTS	NOT DETECTED		
METHOD: MICROSCOPIC EXAMINATION			
CRYSTALS	NOT DETECTED		
METHOD: MICROSCOPIC EXAMINATION			
BACTERIA	NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAMINATION			









CLIENT CODE: C000138361
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL Ltd

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email : wellness.eastdelhi@srl.in

PATIENT NAME: ANKIT SINGH PATIENT ID: ANKIMO2079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

DRAWN: RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Biological Reference Interval Units

REMARKS

MICROSCOPIC EXAMINATION DONE ON CENTRIFUGED URINE PLEASE NOTE THAT GRADING OF BACTERIA NEEDS TO BE CORELATED WITH THE CULTURE IN CASE FOUND SIGNIFICANT CLINICALLY. THE BACTERIA CAN ALSO BE A PART OF SURROUNDING SKIN FLORA.

METHOD: MANUAL

THYROID PANEL, SERUM

T3 144.0 80.00 - 200.00 ng/dL

METHOD : ECLIA

T4 9.64 5.10 - 14.10 μg/dL

METHOD : ECLIA

TSH 3RD GENERATION 1.970 0.270 - 4.200 μIU/mL

METHOD: ECLIA

#### Comments

Note: TSH

Please note that TSH values can show a diurnal variation (up to 50%). Subclinical thyroid/ gut diseases/ intake of calcium, multivitamin and several other drugs, resistance to thyroid hormones, noncompliance to medication can lead to discrepant thyroid results, the levels of thyroid can also be affected by weather, high fiber diet, estrogen surge, stress, alcohol, pregnancy, obesity/weight loss. The results can vary on different instruments. Serum TSH changes significantly in response to even minor changes in thyroid hormones. For the diagnosis of hypo and hyper thyroids, sole dependence on TSH should not be done and testing also needs to be performed for T3, T4 and other metabolic parameters as well as reasons elicited above.

#### **ABO GROUP & RH TYPE, EDTA WHOLE BLOOD**

ABO GROUP TYPE B

METHOD: COLUMN AGGLUTINATION TECHNOLOGY/AGGLUTINATION TECHNOLOGY
RH TYPE POSITIVE

METHOD: COLUMN AGGLUTINATION TECHNOLOGY/AGGLUTINATION TECHNOLOGY

## **XRAY-CHEST**

»» BOTH THE LUNG FIELDS ARE CLEAR

»» BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGELS ARE CLEAR

»» BOTH THE HILA ARE NORMAL

»» CARDIAC AND AORTIC SHADOWS APPEAR NORMAL»» BOTH THE DOMES OF THE DIAPHRAM ARE NORMAL

»» VISUALIZED BONY THORAX IS NORMAL

IMPRESSION NO ABNORMALITY DETECTED

**TMT OR ECHO** 

TMT OR ECHO 2D ECHO DONE-NORMAL STUDY

ECG

ECG WITHIN NORMAL LIMITS



Page 6 Of 13

o View Details Scan to View Report





CLIENT CODE: C000138361
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030

DELHI INDIA 8800465156 SRL Ltd

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email : wellness.eastdelhi@srl.in

PATIENT NAME: ANKIT SINGH PATIENT ID: ANKIMO2079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

DRAWN: RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status Final Results Biological Reference Interval Units

**MEDICAL HISTORY** 

RELEVANT PRESENT HISTORY NOT SIGNIFICANT RELEVANT PAST HISTORY NOT SIGNIFICANT

RELEVANT PERSONAL HISTORY UNMARRIED , NON-VEGETARIAN

RELEVANT FAMILY HISTORY MOTHER - HTN

OCCUPATIONAL HISTORY SENIOR MANAGER

HISTORY OF MEDICATIONS NOT SIGNIFICANT

**ANTHROPOMETRIC DATA & BMI** 

HEIGHT IN METERS1.72mtsWEIGHT IN KGS.79.5Kgs

BMI & Weight Status as follows: kg/sqmts

Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese

**GENERAL EXAMINATION** 

MENTAL / EMOTIONAL STATE NORMAL PHYSICAL ATTITUDE **NORMAL** GENERAL APPEARANCE / NUTRITIONAL STATUS **HFAITHY BUILT / SKELETAL FRAMEWORK AVERAGE** FACIAL APPEARANCE NORMAL SKIN **NORMAL** UPPER LIMB NORMAL LOWER LIMB **NORMAL NECK** NORMAL

NECK LYMPHATICS / SALIVARY GLANDS NOT ENLARGED OR TENDER

THYROID GLAND NOT ENLARGED

CAROTID PULSATION NORMAL TEMPERATURE NORMAL

PULSE 88/MIN REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID

**BRUIT** 

RESPIRATORY RATE NORMAL

**CARDIOVASCULAR SYSTEM** 

BP 124/80 mm/Hg

PERICARDIUM NORMAL APEX BEAT NORMAL





View Details Scan to View Report





CLIENT CODE: C000138361
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030

DELHI INDIA 8800465156 SRL Ltd

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email : wellness.eastdelhi@srl.in

PATIENT NAME: ANKIT SINGH PATIENT ID: ANKIM02079028

ACCESSION NO: **0028VC000973** AGE: 31 Years SEX: Male

DRAWN: RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48

REFERRING DOCTOR: DR. MEDIWHEEL		CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
HEART SOUNDS	S1, S2 HEARD NORMALL	_Y
MURMURS	ABSENT	
RESPIRATORY SYSTEM		
SIZE AND SHAPE OF CHEST	NORMAL	
MOVEMENTS OF CHEST	SYMMETRICAL	
BREATH SOUNDS INTENSITY	NORMAL	
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)	
ADDED SOUNDS	ABSENT	
PER ABDOMEN		
APPEARANCE	NORMAL	
VENOUS PROMINENCE	ABSENT	
LIVER	NOT PALPABLE	
SPLEEN	NOT PALPABLE	
CENTRAL NERVOUS SYSTEM		
HIGHER FUNCTIONS	NORMAL	
CRANIAL NERVES	NORMAL	
CEREBELLAR FUNCTIONS	NORMAL	

HIGHER FUNCTIONS NORMAL
CRANIAL NERVES NORMAL
CEREBELLAR FUNCTIONS NORMAL
SENSORY SYSTEM NORMAL
MOTOR SYSTEM NORMAL
REFLEXES NORMAL

# **MUSCULOSKELETAL SYSTEM**

SPINE NORMAL JOINTS NORMAL

# **BASIC EYE EXAMINATION**

**CONJUNCTIVA NORMAL EYELIDS NORMAL** EYE MOVEMENTS **NORMAL CORNEA NORMAL** DISTANT VISION RIGHT EYE WITH GLASSES **NORMAL** DISTANT VISION LEFT EYE WITH GLASSES **NORMAL** NEAR VISION RIGHT EYE WITH GLASSES **NORMAL** NEAR VISION LEFT EYE WITH GLASSES **NORMAL** COLOUR VISION **NORMAL** 

# **BASIC ENT EXAMINATION**



Page 8 Of 13





**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHT **NEW DELHI 110030** 

**DELHI INDIA** 8800465156

SRL Ltd

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

**PATIENT NAME: ANKIT SINGH** PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final** 

EXTERNAL EAR CANAL NORMAI TYMPANIC MEMBRANE NORMAL

NOSE NO ABNORMALITY DETECTED

**SINUSES** NORMAL

**THROAT** NO ABNORMALITY DETECTED

NOT ENLARGED TONSILS

**SUMMARY** 

RELEVANT HISTORY NOT SIGNIFICANT RELEVANT GP EXAMINATION FINDINGS NOT SIGNIFICANT

RELEVANT NON PATHOLOGY DIAGNOSTICS CHEST X RAY PA-NO ABNORMALITIES DETECTED

> USG ABDOMEN-HEPATOMEGALY WITH FATTY CHANGE LIVER FOLLOW UP TO MD PHYSICIAN WITH ALL REPORTS FOR FURTHER MANAGEMENT. GENERAL PHYSICAL EXAMINATION IS NORMAL.'

Interpretation(s)
BLOOD COUNTS, EDTA WHOLE BLOOD-

REMARKS / RECOMMENDATIONS

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-

Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT - NLR-

The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope. ERYTHRO SEDIMENTATION RATE, BLOOD-

Erythrocyte sedimentation rate (ÉSR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0 -1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

- 1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition
  2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin
  3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition" GLUCOSE, FASTING, PLASMA-

ADA 2021 guidelines for adults, after 8 hrs fasting is as follows: Pre-diabetics: 100 - 125 mg/dL
Diabetic: > or = 126 mg/dL
GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-

Glycosylated hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood, the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia





Page 9 Of 13





**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHT **NEW DELHI 110030** DELHI INDIA 8800465156

SRL Ltd E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

PATIENT ID: **PATIENT NAME: ANKIT SINGH** ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units **Final** 

or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells.

Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia, increased red cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of testing such as glycated serum protein (fructosamine) should be considered.

"Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations."

#### References

- 1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006, 879-884.
- 2. Forsham PH. Diabetes Mellitus: A rational plan for management. Postgrad Med 1982, 71,139-154.
- 3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184. GLUCOSE, POST-PRANDIAL, PLASMA-ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75grams of glucose in 300 ml water, over a period of 5 minutes.

CORONARY RISK PROFILE (LIPID PROFILE), SERUM.

Serum cholesterol is a blood test that can provide valuable information for the risk of coronary artery disease This test can help determine your risk of the build up of plaques in your arteries that can lead to narrowed or blocked arteries throughout your body (atherosclerosis). High cholesterol levels usually don't cause any signs or symptoms, so a cholesterol test is an important tool. High cholesterol levels often are a significant risk factor for heart disease and important for diagnosis of hyperlipoproteinemia, atherosclerosis, hepatic and thyroid diseases.

Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn't need into triglycerides, which are stored in fat cells. High triglyceride levels are associated with several factors, including being overweight, eating too many sweets or drinking too much alcohol, smoking, being sedentary, or having diabetes with elevated blood sugar levels. Analysis has proven useful in the diagnosis and treatment of patients with diabetes mellitus, nephrosis, liver obstruction, other diseases involving lipid metabolism, and various endocrine disorders. In conjunction with high density lipoprotein and total serum cholesterol, a triglyceride determination provides valuable information for the assessment of coronary heart disease risk.It is done in fasting state.

High-density lipoprotein (HDL) cholesterol. This is sometimes called the ""good"" cholesterol because it helps carry away LDL cholesterol, thus keeping arteries open and blood flowing more freely. HDL cholesterol is inversely related to the risk for cardiovascular disease. It increases following regular exercise, moderate alcohol consumption and with oral estrogen therapy. Decreased levels are associated with obesity, stress, cigarette smoking and diabetes mellitus.

SERUM LDL The small dense LDL test can be used to determine cardiovascular risk in individuals with metabolic syndrome or established/progressing coronary artery disease, individuals with triglyceride levels between 70 and 140 mg/dL, as well as individuals with a diet high in trans-fat or carbohydrates. Elevated sdLDL levels are associated with metabolic syndrome and an 'atherogenic lipoprotein profile', and are a strong, independent predictor of cardiovascular disease. Elevated levels of LDL arise from multiple sources. A major factor is sedentary lifestyle with a diet high in saturated fat. Insulin-resistance and pre-diabetes have also been implicated, as has genetic predisposition. Measurement of sdLDL allows the clinician to get a more comprehensive picture of lipid risk factors and tailor treatment accordingly. Reducing LDL levels will reduce the risk of CVD and MI.

Non HDL Cholesterol - Adult treatment panel ATP III suggested the addition of Non-HDL Cholesterol as an indicator of all atherogenic lipoproteins (mainly LDL and VLDL). NICE guidelines recommend Non-HDL Cholesterol measurement before initiating lipid lowering therapy. It has also been shown to be a better marker of risk in both primary and secondary prevention studies.

#### Recommendations:

Results of Lipids should always be interpreted in conjunction with the patient's medical history, clinical presentation and other findings.

NON FASTING LIPID PROFILE includes Total Cholesterol, HDL Cholesterol and calculated non-HDL Cholesterol. It does not include triglycerides and may be best used in patients for whom fasting is difficult. LIVER FUNCTION PROFILE, SERUM-

LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, is chemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of





Scan to View Details





**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHT **NEW DELHI 110030** 

**DELHI INDIA** 8800465156

SRL Ltd

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

PATIENT ID: **PATIENT NAME: ANKIT SINGH** ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final** 

normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc SERUM BLOOD UREA NITROGEN-

Causes of Increased levels

Pre renal

- High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal
- Renal Failure

• Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

- Liver disease

CREATININE, SERUM-

Higher than normal level may be due to:

- Blockage in the urinary tract
- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis
- Muscular dystrophy URIC ACID, SERUM-

Causes of Increased levels Dietary

- High Protein Intake.
- Prolonged Fasting,Rapid weight loss.
- Gout

Lesch nyhan syndrome.

Type 2 DM.

Metabolic syndrome.

Causes of decreased levels

- · Low Zinc Intake
- Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels

- Drink plenty of fluidsLimit animal proteins
- High Fibre foodsVit C Intake
- Antioxidant rich foods

TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUMHuman serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc. ELECTROLYTES (NA/K/CL), SERUM-

Sodium levels are Increased in dehydration, cushing's syndrome, aldosteronism & decreased in Addison's disease, hypopituitarism, liver disease. Hypokalemia (low K) is common in vomiting, diarrhea, alcoholism, folic acid deficiency and primary aldosteronism. Hyperkalemia may be seen in end-stage renal failure, hemolysis, trauma,





Scan to View Details





**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHT **NEW DELHI 110030 DELHI INDIA** 8800465156

SRL Ltd

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

**PATIENT NAME: ANKIT SINGH** PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final** 

Addison's disease, metabolic acidosis, acute starvation, dehydration, and with rapid K infusion. Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bicarbonate, diabetes insipidus, adrenocortical hyperfuction, salicylate intoxication and with excessive infusion of isotonic saline or extremely high dietary intake of salt. Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and prolonged vomiting,
URINALYSIS-Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria,

dehydration, urinary tract infections and acute illness with fever Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection. Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and

proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.

Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia THYROID PANEL, SERUM-

Triiodothyronine T3, is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3

Levels in TOTAL T4 TSH3G TOTAL T3 (μg/dL) 6.6 - 12.4 6.6 - 15.5 6.6 - 15.5 (μIU/mL) 0.1 - 2.5 0.2 - 3.0 0.3 - 3.0 (ng/dL) 81 - 190 100 - 260 100 - 260 Pregnancy First Trimester 2nd Trimester 3rd Trimester

Below mentioned are the guidelines for age related reference ranges for T3 and T4.

T3 T4 (μg/dL) 1-3 day: 8.2 - 19.9 1 Week: 6.0 - 15.9 (ng/dL) New Born: 75 - 260

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group.

Kindly note: Method specific reference ranges are appearing on the report under biological reference range.

- 1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.
- 2. Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.
- 3. Behrman R.E. Kilegman R.M., Jenson H. B. Nelson Text Book of Pediatrics, 17th Edition ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-

Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same

The test is performed by both forward as well as reverse grouping methods.

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.





Scan to View Report





**CLIENT CODE:** C000138361 **CLIENT'S NAME AND ADDRESS:** 

ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHT

**NEW DELHI 110030 DELHI INDIA** 8800465156

E-368, LGF, Nirman Vihar, Near Nirman Vihar Metro

NEW DELHI, 110092 NEW DELHI, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956 Email: wellness.eastdelhi@srl.in

**PATIENT NAME: ANKIT SINGH** PATIENT ID: ANKIM02079028

ACCESSION NO: 0028VC000973 AGE: 31 Years SEX: Male

RECEIVED: 26/03/2022 09:44 REPORTED: 28/03/2022 13:48 DRAWN:

REFERRING DOCTOR: DR. MEDIWHEEL CLIENT PATIENT ID:

Test Report Status Results Units **Final** 

#### MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

**ULTRASOUND ABDOMEN ULTRASOUND ABDOMEN** 

HEPATOMEGALY WITH FATTY CHANGE LIVER.

\*\*End Of Report\*\*

Please visit www.srlworld.com for related Test Information for this accession

Dr. Neena Verma **Pathologist** 

Dr. Shyla Goel, M.B.B.S, DCP **Pathologist** 

## **CONDITIONS OF LABORATORY TESTING & REPORTING**

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All Tests are performed and reported as per the turnaround time stated in the SRL Directory of services (DOS).
- 3. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 4. A requested test might not be performed if:
- a. Specimen received is insufficient or inappropriate specimen quality is unsatisfactory
  - b. Incorrect specimen type
  - c. Request for testing is withdrawn by the ordering doctor
- d. There is a discrepancy between the label on the specimen container and the name on the test requisition form

- 5. The results of a laboratory test are dependent on the quality of the sample as well as the assay technology.
- 6. Result delays could be because of uncontrolled circumstances. e.g. assay run failure.
- 7. Tests parameters marked by asterisks are excluded from the "scope" of NABL accredited tests. (If laboratory is accredited).
- 8. Laboratory results should be correlated with clinical information to determine Final diagnosis.
- 9. Test results are not valid for Medico- legal purposes. 10. In case of gueries or unexpected test results please call at SRL customer care (Toll free: 1800-222-000). Post proper investigation repeat analysis may be carried out.

#### **SRL Limited**

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062



Page 13 Of 13