B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019
Tele: 0141-2293346, 4049787, 9887049787
Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



General Physical Examination

Date of Examination: 10.12.2022
Name: PARI WARAMIYA Age: 26 Sex: Ferrale
DOB: 11.10.1996
Referred By: BOB, (Mediuneel.)
Photo ID: ADHAR ID#: attached,
Ht: 168 (cm) Wt: 68 (Kg)
Chest (Expiration): 96 (cm) Abdomen Circumference: 19 (cm)
Blood Pressure: 122/82 mm Hg PR: 16/min RR: 16/min Temp: Alebrole
BMI. 24.1
Eye Examination: Dis rision Rice. 69.1.6.66, Negr Nisson MG Bleeyes. Normal Colorvision
Other: 1001 Significant
On examination he/she appears physically and mentally fit: Yes/No
Signature Of Examine: Name of Examinee:
Signature Medical Examiner: Name Medical Examiner Name Medical Examiner Name Medical Examiner



Pari




IN PIVUSH GOVAN

M.B.B.S. R.M.R.D. (ALTA)

M.B.B.S. D.E.M. (ACGP. UK)

M.B.B.S. D.E.M. (ACGP. UK)

Heart Rate : 61 bpm / Tested On : 10-Dec-22 12:18:01 / HF 0.05 Hz - LF 100 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s / Refd By.: BOB



Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

:- 10/12/2022 10:13:01

NAME :- Mrs. PARI NARANIYA

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

26 Yrs



Patient ID :-122228495

Ref. By Dr:- BOB

Lab/Hosp:-

Sex / Age :- Female 26
Company :- MediWheel
Sample Type :- EDTA

Date

Sample Collected Time 10/12/2022 10:24:13

Final Authentication: 10/12/2022 14:38:19

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BOD BACKACEEEMALE BELOW 40			
BOB PACKAGEFEMALE BELOW 40			
HAEMOGARAM			
HAEMOGLOBIN (Hb)	11.3 L	g/dL	12.0 - 15.0
TOTAL LEUCOCYTE, COUNT	7.43	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	70.5	%	40.0 - 80.0
LYMPHOCYTE	26.0	%	20.0 - 40.0
EOSINOPHIL	1.1	%	1.0 - 6.0
MONOCYTE	2.2	%	2.0 - 10.0
BASOPHIL	0.2	%	0.0 - 2.0
NEUT#	5.24	10^3/uL	1.50 - 7.00
LYMPH#	1.94	10^3/uL	1.00 - 3.70
EO#	0.04	10^3/uL	0.00 - 0.40
MONO#	0.20	¹ 10^3/uL	0.00 - 0.70
BASO#	0.01	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	4.26	x10^6/uL	3.80 - 4.80
HEMATOCRIT (HCT)	33.00 L	%	36.00 - 46.00
MEAN CORP VOLUME (MCV)	77.4 L	fL	83.0 - 101.0
MEAN CORP HB (MCH)	26.5 L	pg	27.0 - 32.0
MEAN CORP HB CONÇ (MČHC)	34.2	g/dL	31.5 - 34.5
PLATELET COUNT	291	x10^3/uL	150 - 410
RDW-CV	14.0	%	11.6 - 14.0
MENTZER INDEX	18.17		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

AJAYSINGH Technologist

Page No: 1 of 11



Dr. Rashmi Bakshi MBBS. MD (Path) RMC No. 17975/008828

Dr. Goyal

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jainur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgovalspathlab.com | E-mail: drgovalpivush@gmail.com



Date :- 10/12/2022 10:13:01

NAME :- Mrs. PARI NARANIYA

Sex / Age :- Female

Company:-MediWheel Patient ID: -122228495

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type :- EDTA

Sample Collected Time 10/12/2022 10:24:13

Final Authentication: 10/12/2022 14:38:19

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval

Erythrocyte Sedimentation Rate (ESR)

08

mm/hr.

00 - 20

(ESR) Methodology: Measurment of ESR by cells aggregation.

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

: ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia (FBC) Methodology disease. Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance. and MCH, MCV, MCHC, MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L, Japan

AJAYSINGH Technologist

Page No: 2 of 11



Dr. Rashmi Bakshi MBBS. MD (Path) RMC No. 17975/008828

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgovalspathlab.com | E-mail: drgovalpiyush@gmail.com

26 Yrs



Date :- 10/12/2022 10:13:01

NAME :- Mrs. PARI NARANIYA

Patient ID :-122228495

Ref. By Dr:- BOB

Lab/Hosp :-

Company :- MediWheel

Sex / Age :- Female

Final Authentication: 10/12/2022 17:28:21

Sample Type :- EDTA, KOx/Na FLUORIDE-F, KSaw/MadeFCb/Marketerfrine WRIN2F2022 10:24:13

HAEMATOLOGY

Test Name Value Unit Biological Ref Interval

BLOOD GROUP ABO

"B" POSITIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone).

FASTING BLOOD SUGAR (Plasma)
Method:- GOD PAP

104.4

mg/dl

75.0 - 115.0

Impaired glucose tolerance (IGT)

Diabetes Mellitus (DM)

111 - 125 mg/dL

> 126 mg/dL

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

BLOOD SUGAR PP (Plasma)

135.8

mg/dl

70.0 - 140.0

Method:- GOD PAP
Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

URINE SUGAR (FASTING) Collected Sample Received Nil

Nil

AJAYSINGH, KAUSHAL, VIJENDRAMEENA
Technologist
DR.HANSA

Page No: 3 of 11



Dr. Piyush Goyal (D.M.R.D.) Dr. Rashmi Bakshi Dr. Chandrika Gupta

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgovalspathlab.com | E-mail: drgovalpiyush@gmail.com



Date :- 10/12/2022 10:13:01

NAME :- Mrs. PARI NARANIYA

Patient ID :-122228495

Ref. By Dr:- BOB

Lab/Hosp :-

Company :- MediWheel

Sample Type :- STOOL

Sex / Age :- Female 26 Yrs

Sample Collected Time 10/12/2022 10:24:13

Final Authentication: 10/12/2022 12:30:46

CLINICAL PATHOLOGY

Test Name Value Unit **Biological Ref Interval**

STOOL ANALYSIS

PHYSICAL EXAMINATION

MUCUS

BLOOD

MICROSCOPIC EXAMINATION

RBC's

/HPF

WBC/HPF

/HPF

OVA

CYSTS

OTHERS Collected Sample Received

VIJENDRAMEENA **Technologist** DR.HANSA Page No: 4 of 11



Dr. Chandrika Gupta MBBS.MD (Path) RMC NO. 21021/008037

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Sample Type :- PLAIN/SERUM

Website: www.drgovalspathlab.com | E-mail: drgovalpiyush@gmail.com



:- 10/12/2022 10:13:01 Date NAME :- Mrs. PARI NARANIYA Patient ID: -122228495

Ref. By Dr:- BOB

Sex / Age :- Female 26 Yrs

Lab/Hosp :-

Company:- MediWheel

Sample Collected Time 10/12/2022 10:24:13

Final Authentication: 10/12/2022 14:03:42

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	148.56	mg/dl	Desirable <200 Borderline 200-239 High> 240
TRIGLYCERIDES Method:- GPO-PAP	67.59	" mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	36.62	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	100.68	mg/dl	Optimal <100 Near Optimal/above optimal 100-129
•			Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	13.52	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	4.06		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	2.75		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED	422.27	mg/dl	400.00 - 1000.00

TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDLCHOLESTERO InstrumentName: Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROLInstrumentName: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.

TOTAL LIPID AND VLDL ARE CALCULATED

KAUSHAL

Page No: 5 of 11



Dr. Pivush Goval (D.M.R.D.) Dr. Rashmi Bakshi

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/12/2022 10:13:01 Date

Patient ID: -122228495 Ref. By Dr:- BOB

NAME :- Mrs. PARI NARANIYA 26 Yrs

Sample Type :- PLAIN/SERUM

Lab/Hosp :-

Sex / Age :- Female

Company:- MediWheel

Final Authentication: 10/12/2022 17:28:21

Sample Collected Time 10/12/2022 10:24:13

BIOCHEMISTRY							
Test Name	Value	Unit	Biological Ref Interval				
LIVER PROFILE WITH GGT							
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.36	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12				
	*	٠	1month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2 Ref-(ACCP 2020)				
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.17	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL				
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.19	mg/dl	0.30-0.70				
SGOT Method:- IFCC	16.1	U/L	Men- Up to - 37.0 Women - Up to - 31.0				
SGPT Method:- IFCC	19.2	U/L	Men- Up to - 40.0 Women - Up to - 31.0				
SERUM ALKALINE PHOSPHATASE Method:-AMP Buffer	86.60	* IU/L	30.00 - 120.00				
SERUM GAMMA GT Method:- IFCC	12.10	U/L	7.00 - 32.00				
SERUM TOTAL PROTEIN Method:- Biuret Reagent	7.60	g/dl	6.40 - 8.30				
SERUM ALBUMIN Method:- Bromocresol Green	4.47	g/dl	3.80 - 5.00				
SERUM GLOBULIN Method:- CALCULATION	3.13	gm/đl	2.20 - 3.50				
A/G RATIO	1.43		1.30 - 2.50				

Total BilirubinMethodology: Colorimetric method InstrumentName:Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName: Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of huma ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular

dystrophy and organ damage. Alkaline Phosphatase Methodology: AMP Buffer InstrumentName: Randox Rx Imola Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease

TOTAL PROTEIN Methodology: Biuret Reagent InstrumentName: Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology. Bromocresol Green InstrumentName Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys, Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal)

KAUSHAL

Page No: 6 of 11,



Dr. Piyush Goyal (D.M.R.D.) Dr. Rashmi Bakshi

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/12/2022 10:13:01

Sample Type :- PLAIN/SERUM

NAME :- Mrs. PARI NARANIYA

Patient ID :-122228495

Ref. By Dr:- BOB

Lab/Hosp :-

Sex / Age :- Female 26 Yrs

Company :- MediWheel

Sample Collected Time 10/12/2022 10:24:13

Final Authentication: 10/12/2022 14:03:42

BIOCHEMISTRY

BIOCHEMISTRI					
Test Name	Value	Unit	Biological Ref Interval		
SERUM CREATININE Method:- Colorimetric Method	0.72	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20		
SERUM URIC ACID Method:- Enzymatic colorimetric	3.96	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7		

KAUSHAL

Page No: 7 of 11



Dr. Piyush Goyal (D.M.R.D.) Dr. Rashmi Bakshi

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Sample Type :- PLAIN/SERUM

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/12/2022 10:13:01

NAME :- Mrs. PARI NARANIYA

Ref. By Dr:- BOB

Patient ID :-122228495

Lab/Hosp :-

Sex / Age :- Female 26 Yrs

Company :- MediWheel

Sample Collected Time 10/12/2022 10:24:13

Final Authentication: 10/12/2022 14:03:42

BIOCHEMISTRY

	DIOCIAL	1220 2222	
Test Name	Value	Unit	Biological Ref Interval
BLOOD UREA NITROGEN (BUN)	19.7	mg/dl	0.0 - 23.0

KAUSHAL

Page No: 8 of 11



Dr. Piyush Goyal (D.M.R.D.) Dr. Rashmi Bakshi

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/12/2022 10:13:01 Date

NAME :- Mrs. PARI NARANIYA

Sex / Age :- Female 26 Yrs

Company:-MediWheel

Sample Type :- EDTA

Test Name

Method:- HPLC

Patient ID: -122228495

Ref. By Dr:- BOB

Lab/Hosp:-

Final Authentication: 10/12/2022 14:38:19

Sample Collected Time 10/12/2022 10:24:13

HAEMATOLOGY **Biological Ref Interval** Value Unit

GLYCOSYLATED HEMOGLOBIN (HbA1C) 6.0 %

Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0

Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb.High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measureof the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1cmeasurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

Method:- Calculated Parameter

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 mg/dL Diabetic 126 mg/dL or Higher

AJAYSINGH Technologist

Page No: 9 of 11



Dr. Rashmi Bakshi MBBS, MD (Path) RMC No. 17975/008828

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/12/2022 10:13:01 Date

NAME :- Mrs. PARI NARANIYA

Sex / Age :- Female 26 Yrs

Company :- MediWheel

Sample Type :- URINE

Sample Collected Time 10/12/2022 10:24:13

Final Authentication: 10/12/2022 12:30:46

CLINICAL PATHOLOGY

Patient ID: -122228495

Ref. By Dr:- BOB

Lab/Hosp :-

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
COLOUR	PALE YE	LLQW	PALE YELLOW
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH)	5.5		5.0 - 7.5
SPECIFIC GRAVITY	1.020		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR '	NIL		NIL
BILIRUBIN	NEGATIV	/E	NEGATIVE
UROBILINOGEN	NORMAI		NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
W.BC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT	1	ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		

VIJENDRAMEENA **Technologist** DR.HANSA Page No: 10 of 11



Dr. Chandrika Gupta MBBS.MD (Path) RMC NO. 21021/008037

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/12/2022 10:13:01

NAME :- Mrs. PARI NARANIYA

Sex / Age :- Female 26 Yrs

Company :- MediWheel
Sample Type :- PLAIN/SERUM

Patient ID :-122228495

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 10/12/2022 10:24:13 Final Authentication: 10/12/2022 14:15:26

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.125	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	8.589	ug/dl	5.500 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	1.380	$\mu IU/mL$	0.500 - 6.880

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy.that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation: TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

*** End of Report ***

KAUSHAL Technologist

Page No: 11 of 11



Dr. Chandrika Gupta MBBS.MD (Path) RMC NO. 21021/008037



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/12/2022 10:13:01

:- Mrs. PARI NARANIYA

Sex / Age :- Female 26 Yrs MediWheel Company :-

NAME

Patient ID: -122228495 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 10/12/2022 12:15:13

BOB PACKAGEFEMALE BELOW 40

X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P. angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression :- Normal Study

(Please correlate clinically and with relevant further investigations)

*** End of Report ***

Dr. Piyush Goyal (D.M.R.D.) BILAL

Page No: 1 of 1

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 82495

Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant

FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain MBBS, DNB, (Radio-Diagnosis) RMC No. 21687

Transcript by.



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 10/12/2022 10:13:01 NAME :- Mrs. PARI NARANIYA

26 Yrs Sex / Age :- Female Company :- MediWheel

Date

Patient ID: -122228495 Ref. By Doctor:-BOB

Lab/Hosp:-

Final Authentication: 10/12/2022 11:54:09

BOB PACKAGEFEMALE BELOW 40

ULTRA SOUND SCAN OF ABDOMEN

Liver is of normal size. Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Urinary Bladder: is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

Uterus is anteverted and normal in size and measures: 78x48x39 mm. Myometrium shows normal echo - pattern. No focal space occupying lesion is seen. Endometrial echo is normal. Endometrial thickness is 6.7 mm.

Both ovaries are visualised and are normal. No adnexal mass is seen.

No enlarged nodes are visualised. No retro-peritoneal lesion is identified. No significant free fluid is seen in pouch of douglas.

IMPRESSION:

Normal Study.

Needs clinical correlation & further evaluation

*** End of Report ***

AHSAN

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Ashish Choudhary MBBS, MD (Radio Diagnosis) Fetal Medicine Consultant FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain MBBS, DNB, (Radio-Diagnosis) RMC No. 21687

Transcript by.



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/12/2022 10:13:01

NAME :- Mrs. PARI NARANIYA

Sex / Age :- Female 26 Yrs Company :- MediWheel Patient ID:-122228495 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 10/12/2022 11:56:41

BOB PACKAGEFEMALE BELOW 40 2D ECHO OPTION TMT (ADULT/CHILD)

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

/E	NOR	MAL	TRICUSPID VALVE			NORMAL	
/E	NOR	ORMAL PULMONARY VALVE			NORMAL		
	M.MODE	EXAMITATION:					
24	mm	LA	30	Mm	IVS-D	8	mm
13	mm	LVID	47	Mm	LVSD	30	mm
7	mm	LVPW-S	13	Mm	RV		mm
	mm	EDV		MI	LVVS		ml
64%			RWMA		ABSENT		
	24 13 7	/E NOR M.MODE 24 mm 13 mm 7 mm	NORMAL M.MODE EXAMITATION: 24 mm	NORMAL PULMO	NORMAL	NORMAL	NORMAL

CHAMBERS:

LA	NORMAL	RA	NORMAL	
LV	NORMAL	RV	NORMAL	
PERICARDIUM		NORMAL		

COLOUR DOPPLER:

	M	TRAL VALVE						
E VELOCITY	0.74	m/sec	PEAK	GRADIENT		Mn	Mm/hg	
A VELOCITY	0.57	m/sec MEAN		GRADIENT		Mn	Mm/hg	
MVA BY PHT		Cm2	MVA	Y PLANIMETRY		Cm	Cm2	
MITRAL REGURGITAT	ION				ABSENT			
	AC	RTIC VALVE						
PEAK VELOCITY	1.0	m/s	sec	PEAK GRADIENT		m	mm/hg	
AR VMAX	m		sec MEAN GR		RADIENT	m	mm/hg	
AORTIC REGURGITATION				ABSENT				
	TRI	CUSPID VAL	VE					
PEAK VELOCITY	0.46		m/sec	PEAK GRADIENT			mm/hg	
MEAN VELOCITY		ı	m/sec	MEAN GRADIENT			mm/hg	
VMax VELOCITY								
TRICUSPID REGURGITATION				ABSENT				
	PL	ILMONARY \	VALVE					
PEAK VELOCITY 0.5		0.98		M/sec.	PEAK GRADIENT	PEAK GRADIENT		
MEAN VALOCITY					MEAN GRADIENT		Mm/hg	
PULMONARY REGURGITATION					ABSENT			

Page No: 1 of 2

AHSAN



B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 10/12/2022 10:13:01
NAME :- Mrs. PARI NARANIYA

Sex / Age :- Female 26 Yrs Company :- MediWheel Patient ID:-122228495 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 10/12/2022 11:56:41

Impression--

- 1. Normal LV size & contractility
- 2. No RWMA, LVEF 64 %.
- 3. Normal cardiac chamber.
- 4. Normal valve
- 5. No clot, no vegetation, no pericardial effusion. (Cardiologist)

*** End of Report ***

Name: PARI NARANIYA / F

