



for Medical Purposes

[Handwritten signature]

[Handwritten signature]



Dr. RIYUSH GOYAL
MBBS, DMRD (Radiologist)
RMC No.-037041



📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com

General Physical Examination

Date of Examination: 15/10/2023

Name: Karanveer Singh Bindra Age: 33 DOB: 29/01/1990 Sex: Male

Referred By: Bank of Baroda

Photo ID: Adhar Card ID #: 7119

Ht: 185 (cm)

Wt: 127 (Kg)

Chest (Expiration): 127 (cm)

Abdomen Circumference: 144 (cm)

Blood Pressure: 130/80 mm Hg PR: 78 / min RR: 18 / min Temp: Alebiale

BMI 37

Eye Examination: R/E, 6/6 N/G, NCB
L/E 6/6 N/G NCB

Other: no

On examination he/she appears physically and mentally fit: Yes / No

Signature Of Examinee:

Name of Examinee: Karanveer Singh Bindra

Signature Medical Examiner:
PIYUSH GOYAL
MBBS, DNB (Radiologist)
RMC No.-037041

Name Medical Examiner: Dr. Piyush Goyal



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)

Ⓞ B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
Ⓞ +91 141 4824885 Ⓞ maxcarediagnostics1@gmail.com



NAME :- Mr. KARANVEER SINGH BINDRA
Age :- 33 Yrs 8 Mon 16 Days
Sex :- Male

Patient ID :-42233749 Date :- 15/10/2023 08:40:35
Ref. By Doctor:-BANK OF BARODA
Lab/Hosp :-
Company :- Mr.MEDIWHEEL

Final Authentication : 15/10/2023 13:39:23

HAEMOGARAM

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
FULL BODY HEALTH CHECKUP BELOW 40 MALE			
HAEMOGLOBIN (Hb)	14.2	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	7.40	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	62.0	%	40.0 - 80.0
LYMPHOCYTE	31.0	%	20.0 - 40.0
EOSINOPHIL	3.0	%	1.0 - 6.0
MONOCYTE	4.0	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	4.97	$\times 10^6/\mu\text{L}$	4.50 - 5.50
HEMATOCRIT (HCT)	44.90	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	90.0	fL	83.0 - 101.0
MEAN CORP HB (MCH)	28.5	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	31.6	g/dL	31.5 - 34.5
PLATELET COUNT	293	$\times 10^3/\mu\text{L}$	150 - 410
RDW-CV	13.6	%	11.6 - 14.0

Technologist
VIKARANTSIJ
Page No. 1 of 16

Tanu
DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



Ⓞ B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
Ⓞ +91 141 4824885 Ⓞ maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

09

mm in 1st hr

00 - 15

Method :- Westergren

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases. ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
VIKARUNU J

Page No. 2 of 16



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-42233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance, and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 pari fully automatic analyzer XN-L,Japan





P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com




NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :42233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) <small>Method - GOD POB</small>	89.0	mg/dl	70.0 - 115.0
Impaired glucose tolerance (IGT)	111 - 125 mg/dL		
Diabetes Mellitus (DM)	> 126 mg/dL		

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.



Technologist
VIKAS RANJAN

Tanu Rungta
DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. KARANVEER SINGH BINDRA

Age :- 33 Yrs 8 Mon 16 Days

Sex :- Male

Patient ID :-12233749

Date :- 15/10/2023 08:40:35

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 15/10/2023 14:35:46

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

FULL BODY HEALTH CHECKUP BELOW 40 MALE

BLOOD SUGAR PP (Plasma)

112.3

mg/dl

70.0 - 140.0

Method - GOD PAP

Instrument Name: HORIBA **Interpretation:** Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases.

*** End of Report ***



Technologist
VIKASH KANTH
Page No. 1 of 1

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (HbA1C) Method - CAPILLARY with EDTA	5.1	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Method - Calculated Parameter	102	mg/dL	68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Reference Group HbA1c in %
Non diabetic adults >=18 years < 5.7
At risk (Prediabetes) 5.7 - 6.4
Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In-vitro quantitative determination of HbA1c in whole blood is valued in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings.
Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al.]

1. Erythropoiesis

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, sickle cell disease, chronic liver disease.

2. Altered hemoglobin-Genetic or chemical alterations in hemoglobin, hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intracellular pH
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH

4. Erythrocyte destruction

- Increased HbA1c: increased erythrocyte lifespan, Splenectomy
- Decreased HbA1c: decreased RBC life span, hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as aniline dyes, ribavirin & dapsone

5. Others

- Increased HbA1c: hyperlipidemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
- Decreased HbA1c: hyperglycemia, sickle cell disease, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Note:

1. Shortened RBC life span-HbA1c test will not be accurate when a person has a condition that affects the average lifespan of red blood cells (RBCs), such as hemolytic anemia or blood loss. When the lifespan of RBCs in circulation is shortened, the A1c result is falsely low and is an unreliable measurement of a person's average glucose over time.
2. Abnormal forms of hemoglobin - The presence of some hemoglobin variants, such as hemoglobin S in sickle cell anemia, may affect certain methods for measuring A1c. In these cases, fructosamine can be used to monitor glucose control.

Advised:

1. To follow patient for glycemic control test like HbA1c or glycolated albumin may be performed instead.
2. Hemoglobin HPLC screen to analyze abnormal hemoglobin variant.

Technologist
VIKARAN S
Page No. 5 of 16

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)



📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

HAEMATOLOGY

BLOOD GROUP ABO
Method :- Haemagglutination reaction

"B" POSITIVE



Technologist
VIKARAN JI
Page No. 6 of 16

Tanu
DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-42233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 15/10/2023 13:39:23

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIPID PROFILE			
TOTAL CHOLESTEROL Method - CHOD-PAP methodology	125.00	mg/dl	Desirable <200 Borderline 200-239 High > 240
<i>InstrumentName: MISPA PLUS Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.</i>			
TRIGLYCERIDES Method - GPO-PAP	101.00	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
<i>InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.</i>			
DIRECT HDL CHOLESTEROL Method - Direct clearance Method	46.80	mg/dl	MALE - 30-70 FEMALE - 30-85
<i>Instrument Name: Rx Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.</i>			
LDL CHOLESTEROL Method - Calculated Method	61.37	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method - Calculated	20.20	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method - Calculated	2.67		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method - Calculated	1.31		0.00 - 3.50
TOTAL LIPID Method - CALCULATED	402.19	mg/dl	400.00 - 1000.00
I. Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.			

Technologist
VIKARAN SINGH
Page No. 7 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-42233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

BIOCHEMISTRY

2. As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.
3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

Comments: 1- ATP III suggested the addition of Non HDL Cholesterol (Total Cholesterol – HDL Cholesterol) as an indicator of all atherogenic lipoproteins (mainly LDL & VLDL). The Non HDL Cholesterol is used as a secondary target of therapy in persons with triglycerides ≥ 200 mg/dL. The goal for Non HDL Cholesterol in those with increased triglyceride is 30 mg/dL, above that set for LDL Cholesterol.
2 -For calculation of CHD risk, history of smoking, any medication for hypertension & current B.P. levels are required.



DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
VIKARANI D

Page No: 6 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

BIOCHEMISTRY

LIVER PROFILE WITH GGT

SERUM BILIRUBIN (TOTAL) Method:- DMSO/Diaz	0.62	mg/dl.	Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Method:- DMSO/Diaz	0.23	mg/dl.	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.39	mg/dl	0.30-0.70
SGOT Method:- IFCC	76.9 H	U/L	0.0 - 40.0
SGPT Method:- IFCC	64.1 H	U/L	0.0 - 40.0
SERUM ALKALINE PHOSPHATASE Method:- DGKC - SCT	96.80	U/L	53.00 - 141.00
SERUM GAMMA GT Method:- S-zinc methodology Instrument: Nemo Random Ms India Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 50 times normal levels in cirrhosis or post- hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.	19.20	U/L	10.00 - 45.00
SERUM TOTAL PROTEIN Method:- Direct Biamer Reagent	6.59	g/dl	6.00 - 8.40
SERUM ALBUMIN Method:- Bromocresol Green	3.98	g/dl	3.50 - 5.50
SERUM GLOBULIN Method:- CALCULATION	2.61	gm/dl	2.20 - 3.50
A/G RATIO	1.52		1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note :- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B ,C ,paracetamol toxicity etc Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.

Technologist
Vikas Ranjan

Tanu Rungta
DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA

Age :- 33 Yrs 8 Mon 16 Days

Sex :- Male

Patient ID :-12233749

Date :- 15/10/2023 08:40:35

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 15/10/2023 13:39:23

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA 21.60 mg/dl 10.00 - 50.00
Method:- Urease/GLDH

InstrumentName: HORIBA CA 60 Interpretation : Urea measurements are used in the diagnosis and treatment of certain renal and metabolic diseases.

SERUM CREATININE 1.13 mg/dl Males : 0.6-1.50 mg/dl
Females : 0.6 -1.40 mg/dl
Method:- Jaffe's Method

Interpretation :
Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not clinically significant.

SERUM URIC ACID 5.96 mg/dl 2.40 - 7.00

InstrumentName: HORIBA YUMIZEN CA60 Daytona plus Interpretation: Elevated Urate: High purine diet, Alcohol, Renal insufficiency, Drugs, Polycythaemia vera, Malignancies, Hypothyroidism, Rare enzyme defects, Downs syndrome, Metabolic syndrome, Pregnancy, Gout.

SODIUM 140.7 mmol/L 135.0 - 150.0
Method:- ISE
Interpretation:

Electrolytes are minerals that are found in body tissues and blood in the form of dissolved salts. As electrically charged particles, electrolytes help move nutrients into and wastes out of the body's cells, maintain a healthy water balance, and help stabilize the body's acid/base (pH) level. The electrolyte panel measures the blood levels of the main electrolytes in the body. *

* **Sodium**—most of the body's sodium is found in the fluid outside of the body's cells, where it helps to regulate the amount of water in the body. *

POTASSIUM 4.70 mmol/L 3.50 - 5.50
Method:- ISE

* **Potassium**—this electrolyte is found mainly inside the body's cells. A small but vital amount of potassium is found in the plasma, the liquid portion of the blood. Potassium plays an important role in regulating muscle contraction. Monitoring potassium is important as small changes in the potassium level can affect the heart's rhythm and ability to contract.

CHLORIDE 100.8 mmol/L 94.0 - 110.0
Method:- ISE

Technologist
VIKARAN (9)
Page No: 10 of 18

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA

Age :- 33 Yrs 8 Mon 16 Days

Sex :- Male

Patient ID :-12233749

Date :- 15/10/2023 08:40:35

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 15/10/2023 13:39:23

BIOCHEMISTRY

* **Chloride**—this electrolyte moves in and out of the cells to help maintain electrical neutrality (concentrations of positively charged cations and negatively charged anions must be equal) and its level usually mirrors that of sodium. Due to its close association with sodium, chloride also helps to regulate the distribution of water in the body.

SERUM CALCIUM	9.89	mg/dL	8.80 - 10.20
Method - Arsenazo III Method			

InstrumentName: MISPA PLUS **Interpretation:** Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia. Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN	6.59	g/dl	6.00 - 8.40
Method - Direct Buret Reagent			

SERUM ALBUMIN	3.98	g/dl	3.50 - 5.50
Method - Bromocresol Green			

SERUM GLOBULIN	2.61	gm/dl	2.20 - 3.50
Method - CALCULATION			

A/G RATIO	1.52		1.30 - 2.50
-----------	------	--	-------------

Interpretation - Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR. In urine, it can remove the need for 24-hour collections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection. Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the blood increases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.

Technologist
VIKARANJJI
Page No: 11 of 18

Tanu

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

TOTAL THYROID PROFILE

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

THYROID-TRIiodothyronine T3
Method- ECLIA

1.25 ng/mL

0.70 - 2.04

NOTE-T3H levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 8-10 PM. The variation is the order of 50% hence time of the day has influence on the measured serum T3H concentration. Dose and time of drug intake also influence the test result. Transient increase in T3H levels or abnormal T3H levels can be seen in some non thyroidal conditions. Simultaneous measurement of T3H with free T4 is useful in evaluating differential diagnosis.

INTERPRETATION-Ultra Sensitive 4th generation assay. 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↑ TSH level. 2.Low TSH/High FT4 and TSH receptor antibody (TRAb) -we seen in patients with Graves disease. 3.Low TSH/High FT4 and TSH receptor antibody (TRAb) -we seen in patients with Toxic adenoma/Toxic Multinodular goiter. 4.High TSH/Low FT4 and Thyroid microsomal antibody (increased seen in patients with Hashimoto's thyroiditis). 5.High TSH/Low FT4 and Thyroid microsomal antibody normal seen in patients with iodine deficiency/ Congenital T4 synthesis deficiency. 6.Low TSH/Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism. 7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels. Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis. Normal or T3 & T4 Normal T3 & T4 along with ↑ TSH indicate mild / Subclinical Hyperthyroidism. 11 Normal T3 & T4 along with ↑ TSH is seen in Hypothyroidism. 12 Normal T3 & T4 levels with ↑ TSH indicate Mild / Subclinical Hypoth

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL, 2nd Trimester : 0.20-3.00 uIU/mL, 3rd Trimester : 0.30-3.00 uIU/mL. The production, circulation, and degradation of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radioactive scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of uncorrected thyroid disease in the elderly. ** 5.10 - 14.10

THYROID-THYRONINE (T4)

Method- ECLIA

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 8-10 PM. The variation is the order of 50% hence time of the day has influence on the measured serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions. Simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis.

INTERPRETATION-Ultra sensitive 4th generation assay. 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↑ TSH level. 2.Low TSH/High FT4 and TSH receptor antibody (TRAb) -we seen in patients with Graves disease. 3.Low TSH/High FT4 and TSH receptor antibody (TRAb) -we seen in patients with Toxic adenoma/Toxic Multinodular goiter. 4.High TSH/Low FT4 and Thyroid microsomal antibody (increased seen in patients with Hashimoto's thyroiditis). 5.High TSH/Low FT4 and Thyroid microsomal antibody normal seen in patients with iodine deficiency/ Congenital T4 synthesis deficiency. 6.Low TSH/Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism. 7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels. Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis. Normal or T3 & T4 Normal T3 & T4 along with ↑ TSH indicate mild / Subclinical Hyperthyroidism. 11 Normal T3 & T4 along with ↑ TSH is seen in Hypothyroidism. 12 Normal T3 & T4 levels with ↑ TSH indicate Mild / Subclinical Hypoth

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL, 2nd Trimester : 0.20-3.00 uIU/mL, 3rd Trimester : 0.30-3.00 uIU/mL. The production, circulation, and degradation of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radioactive scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of uncorrected thyroid disease in the elderly.

TSH

Method- ECLIA

1.430 uIU/mL

0.350 - 5.500

4th Generation Assay/Reference ranges vary between laboratories

Tanu Rungta

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
VIKARAN (P)
Page No: 14 of 16



📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

IMMUNOASSAY

PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association)

1st Trimester : 0.10-2.50 uIU/mL

2nd Trimester : 0.20-3.00 uIU/mL

3rd Trimester : 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

- 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level.
- 2.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
- 3.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4.Normal or ↓ T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism

COMMENTS: Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

Reference ranges are from Teitz fundamental of clinical chemistry 8th ed (2016)

Test performed by Instrument : Beckman coulter Dxi 600

Note - The result obtained relate only to the sample given/ received & tested. A single test result is not always indicative of a disease, it has to be correlated with

4th Generation Assay, Reference ranges vary between laboratories

PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association)

1st Trimester : 0.10-2.50 uIU/mL

2nd Trimester : 0.20-3.00 uIU/mL

3rd Trimester : 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

- 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level.
- 2.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
- 3.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4.Normal or ↓ T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism

Tanu Rungta

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
VIKARANTSI
Page No. 15 of 16



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 15/10/2023 13:39:23

. **COMMENTS:** Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

. **Disclaimer:** TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

. **Reference ranges are from Toltz fundamental of clinical chemistry 8th ed (2018)**
Test performed by instrument : Beckman coulter Dxi 800

. **Note:** The result obtained relate only to the sample given/ received & tested. A single test result is not always indicative of a disease. It has to be correlated with clinical data for interpretation.

*** End of Report ***



Technologist
VIKRAM LAL
Page No. 16 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)




B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com

NAME :- Mr. KARANVEER SINGH BINDRA	Patient ID :-12233749	Date :- 15/10/2023	08:40:35
Age :- 33 Yrs 8 Mon 16 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 15/10/2023 13:39:23

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH)	5.0		5.0 - 7.5
SPECIFIC GRAVITY	1.025		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIVE		NEGATIVE
UROBILINOGEN	NORMAL		NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT



Technologist
VIKARANTSI

Tanu Rungta

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- ☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME:	MR.KARANVEER SINGH BINDRA	AGE	33 YRS/M
REF.BY	BANK OF BARODA	DATE	15/10/2023

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected

DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



MR. KARANVEER SINGH BINDRA	Age:- 33 Y/Male
Registration date: 15/10/2023	Ref. By: BANK OF BARODA

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:
FAIR TRANSTHORACIC ECHOCARDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	NORMAL	TRICUSPID VALVE	NORMAL
AORTIC VALVE	NORMAL	PULMONARY VALVE	NORMAL

M.MODE EXAMINATION:

AO	2.8	cm	LA	3.2	cm	IVS-D	1.1	cm
IVS-S	1.3	cm	LVID	4.4	cm	LVSD	3.8	cm
LVPW-D	0.9	cm	LVPW-S	1.2	cm	RV		cm
RVWT		cm	EDV		ml	LVVS		ml
LVEF	60%		RWMA			ABSENT		

CHAMBERS:

LA	NORMAL	RA	NORMAL
LV	NORMAL	RV	NORMAL
PERICARDIUM	NORMAL		

COLOUR DOPPLER:

MITRAL VALVE				
E VELOCITY	0.86	m/sec	PEAK GRADIENT	Mm/hg
A VELOCITY	0.71	m/sec	MEAN GRADIENT	Mm/hg
MVA BY PHT		Cm2	MVA BY PLANIMETRY	Cm2
MITRAL REGURGITATION	ABSENT			
AORTIC VALVE				
PEAK VELOCITY	1.04	m/sec	PEAK GRADIENT	mm/hg
AR VMAX		m/sec	MEAN GRADIENT	mm/hg
AORTIC REGURGITATION	ABSENT			
TRICUSPID VALVE				
PEAK VELOCITY	1.11	m/sec	PEAK GRADIENT	mm/hg
MEAN VELOCITY		m/sec	MEAN GRADIENT	mm/hg
VMax VELOCITY				
TRICUSPID REGURGITATION	ABSENT			
PULMONARY VALVE				
PEAK VELOCITY	0.66	M/sec.	PEAK GRADIENT	Mm/hg
MEAN VELOCITY			MEAN GRADIENT	Mm/hg
PULMONARY REGURGITATION	ABSENT			

Impression—

- NORMAL LV SIZE & CONTRACTILITY
- NO RWMA, LVEF 60%.
- ALL CARDIAC VALVES ARE NORMAL.
- NO CLOT, NO VEGETATION, NO PERICARDIAL EFFUSION.
- NORMAL LV DIASTOLIC FUNCTION.

(Cardiologist)

Dr. JYOTI AGARWAL
M.B.B.S, PGDCC (Cardiologist)
RMC No.- 27255





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



MR. KARANVEER SINGH BINDRA	33 YEARS/Male
Registration Date: 15/10/2023	Ref. by: BANK OF BARODA

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (14.7 cm) with **bright parenchymal echotexture**. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape (11.3 cm). Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Right kidney is measuring approx. 12.2 x 5.4 cm.

Left kidney is measuring approx. 11.7 x 5.5 cm.

Urinary bladder is normally distended and shows normal wall thickness. No calculus or mass lesion.

Prostate is normal in size with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified.

No significant free fluid is seen in pelvis.


IMPRESSION:-

- **Grade I fatty liver.**
- **Rest no significant abnormality is detected.**



DR. SHALINI GOEL
M.B.B.S, D.N.B (Radiodiagnosis)
RMC no.: 21954

SHALINI GOEL
BBS, DNB (Radiologist)
RMC No. 21954
P-3 Health Solutions LLP



FINDINGS: Normal Sinus Rhythm

Vent Rate : 71 bpm; PR Interval : 162 ms; QRS Duration: 102 ms; QT/QTc Int : 362/396 ms

P-QRS-T axis: 10 • -10 • 39 • (Deg)

Comments :

WNL

BBS, D.EM. (FCGP,UK)



भारत सरकार

GOVERNMENT OF INDIA



करणवीर सिंह बिन्द्रा

Karanveer Singh Bindra

जन्म तिथि/ DOB: 29/01/1990


पुरुष / MALE



2326 4708 7119

मेरा आधार, मेरी पहचान



 **GPS Map Camera**

Jaipur, Rajasthan, India

G-22 Vidhadher Enclave 14, near Cine Star, Sector 2, Central Spine,
Vidyadhar Nagar, Jaipur, Rajasthan 302039, India


Lat 26.964676°

Long 75.782386°

15/10/23 09:43 AM GMT +05:30





 **GPS Map Camera**


Jaipur, Rajasthan, India

P. No. B - 14, G - 47, Vidhyadhar Enclave Iled, Central Spine Rd, Sector 2,
Central Spine, Vidyadhar Nagar, Jaipur, Rajasthan 302023, India

Lat 26.964511°

Long 75.782498°

15/10/23 09:41 AM GMT +05:30



Google



12233749 KARANVEER SINGH BHIRA BOB M
18 OCT 2023
MAXCARE DIAGNOSTIC (ASSOCIATES OF P3 HEALTH SOLUTIONS LLP)

