PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	13.9	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	42.5	%	42 - 52
RBC Count (EDTA Blood/Impedance Variation)	5.09	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	83.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	27.2	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	32.6	g/dL	32 - 36
RDW-CV (Derived from Impedance)	15.2	%	11.5 - 16.0
RDW-SD (Derived from Impedance)	44.16	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	6400	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	48.2	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	37.1	%	20 - 45
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	6.6	%	01 - 06





 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes	7.0	%	01 - 10
(Blood/Impedance Variation & Flow Cytometry)			
Basophils (Blood/Impedance Variation & Flow Cytometry)	1.1	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.08	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.37	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.42	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.45	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.07	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	321	10^3 / μl	150 - 450
MPV (Blood/Derived from Impedance)	8.1	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.26	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	7	mm/hr	< 20





 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.6	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.4	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	6.9	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.0	gm/dL	3.5 - 5.2
Globulin (Serum/Derived)	2.9	gm/dL	2.3 - 3.6
A: G Ratio (Serum/Derived)	1.4		1.1 - 2.2
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC Kinetic)	17	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	17	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/IFCC Kinetic)	76	U/L	56 - 119
GGT(Gamma Glutamyl Transpeptidase) (Serum/SZASZ standarised IFCC)	69	U/L	< 55





 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	220	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	124	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	52	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	143.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	24.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	168.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220





PID No. : MED120883125 Register On : 12/03/2022 8:46 AM : 522211325 SID No. Collection On : 12/03/2022 12:48 PM Age / Sex : 53 Year(s) / Male Report On : 13/03/2022 11:49 AM **Type** : OP **Printed On** : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation Observed **Unit Biological** Reference Interval **Value**

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio 4.2 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated)

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio Optimal: < 2.52.4

Mild to moderate risk: 2.5 - 5.0 (TG/HDL)

High Risk: > 5.0(Serum/Calculated)

LDL/HDL Cholesterol Ratio 2.8 Optimal: 0.5 - 3.0

Borderline: 3.1 - 6.0 (Serum/Calculated) High Risk: > 6.0



VERIFIED BY



 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	5.4	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 108.28 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.





 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.04 ng/mL 0.4 - 1.81

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

T4 (Thyroxine) - Total 6.90 μg/dL 4.2 - 12.0

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.09 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Microparticle

Îmmunoassay(CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'B' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



VERIFIED BY



 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
BIOCHEMISTRY			
BUN / Creatinine Ratio	10		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	85	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Postprandial (PPBS) 94 mg/dL 70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	10	mg/dL	7.0 - 21
(Serum/ <i>Urease-GLDH</i>)			
Creatinine	1.0	mg/dL	0.9 - 1.3
(0 / / 00 / 1)		· ·	

(Serum/Jaffe Kinetic)

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 5.3 mg/dL 3.5 - 7.2

(Serum/Uricase/Peroxidase)





 PID No.
 : MED120883125
 Register On
 : 12/03/2022 8:46 AM

 SID No.
 : 522211325
 Collection On
 : 12/03/2022 12:48 PM

 Age / Sex
 : 53 Year(s) / Male
 Report On
 : 13/03/2022 11:49 AM

 Type
 : OP
 Printed On
 : 17/03/2022 2:07 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>IMMUNOASSAY</u>			
Prostate specific antigen - Total(PSA) (Serum/Chemiluminescent Microparticle Immunoassay(CMIA))	1.37	ng/mL	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0



-- End of Report --

Name	MR.Y RENUKA PRASAD	ID	MED120883125
Age & Gender	53Y/MALE	Visit Date	12 Mar 2022
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is enlarged in size (16.0 cms) and shows mild diffusely increased echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER show normal shape and has clear contents. Wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN is upper limit of normal size, measures 12 cms. Normal in shape and echopattern. No focal lesions.

No demonstrable Para-aortic lymphadenopathy.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cm)	Parenchymal thickness (cm)
Right Kidney	10.5	1.9
Left Kidney	10.4	1.5

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

Prevoid: 193 cc Postvoid: 12cc

PROSTATE is enlarged in size. It measures 4.2 x 4.6 x 3.6 cm, volume: 36cc.

No evidence of ascites.

IMPRESSION:

- Mild hepatomegaly with grade I fatty infiltration.
- Prostatomegaly with insignificant postvoid residue.

DR. H.K. ANAND DR. C.R RAMACHANDRA DR. LOHITH H.P DR. VARSHA KALE CONSULTANT RADIOLOGISTS

Vk/Ss

Name	MR.Y RENUKA PRASAD	ID	MED120883125
Age & Gender	53Y/MALE	Visit Date	12 Mar 2022
Ref Doctor Name	MediWheel	-	

2D ECHOCARDIOGRAPHIC STUDY

M-mode measurement:

AORTA 3.23 cms. LEFT ATRIUM 3.38 cms. **AVS** 1.47 cms. LEFT VENTRICLE 5.02 (DIASTOLE) cms. (SYSTOLE) 2.60 cms. **VENTRICULAR SEPTUM** (DIASTOLE) 0.96 cms. (SYSTOLE) 1.45 cms. **POSTERIOR WALL** (DIASTOLE) 0.96 cms. (SYSTOLE) 1.40 cms. **EDV** 119 ml. **ESV** 24 ml. % FRACTIONAL SHORTENING 30 **EJECTION FRACTION** % 60 **EPSS** cms. **RVID** 1.80 cms.

DOPPLER MEASUREMENTS:

MITRAL VALVE: $E - 0.8 \text{ m/s} \quad A - 0.7 \text{ m/s}$ NO MR.

AORTIC VALVE: 1.0 m/s NO AR.

TRICUSPID VALVE: E - 0.4 m/s A - 0.3 m/s, MILD TR, PASP 10mmHg.

PULMONARY VALVE: 0.8 m/s NO PR.

Name	MR.Y RENUKA PRASAD	ID	MED120883125
Age & Gender	53Y/MALE	Visit Date	12 Mar 2022
Ref Doctor Name	MediWheel		

2D ECHOCARDIOGRAPHY FINDINGS:

Left Ventricle : Normal size, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal.Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- NORMAL SIZED CARDIAC CHAMBERS.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. ANAND KUMAR M MD DM
CONSULTANT INTERVENTIONAL CARDIOLOGIST

Name	MR.Y RENUKA PRASAD	ID	MED120883125
Age & Gender	53Y/MALE	Visit Date	12 Mar 2022
Ref Doctor Name	MediWheel	-	



Name	Y RENUKA PRASAD	ID	MED120883125
Age & Gender	53Y/M	Visit Date	Mar 12 2022 8:45AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

DR. H.K. ANAND

• No significant abnormality detected.

DR. C.R. RAMACHANDRA DR. VARSHA KALE
CONSULTANT RADIOLOGISTS

DR. LOHITH H.P