





PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO: 0022VL002012

AGE: 36 Years

SEX: Male

ABHA NO:

10/12/2022 14:42:03

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

Final

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status

Results

Biological Reference Interval

Units

SPECIALISED CHEMISTRY - HORMONE

THYROID PANEL, SERUM

T3

87.7

80 - 200

ng/dL

T4

METHOD: ELECTROCHEMILUMINESCENCE, COMPETITIVE IMMUNOASSAY

5.1 - 14.1

µg/dL

METHOD: ELECTROCHEMILUMINESCENCE, COMPETITIVE IMMUNOASSAY

TSH (ULTRASENSITIVE)

1.420

0.270 - 4.200

µIU/mL

METHOD: ELECTROCHEMILUMINESCENCE, COMPETITIVE IMMUNOASSAY

Interpretation(s)















PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002012

AGE: 36 Years

SEX: Male

ABHA NO:

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

10/12/2022 14:42:03

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status

Results

Biological Reference Interval

Final

Units

SPECIALISED CHEMISTRY - TUMOR MARKER

PROSTATE SPECIFIC ANTIGEN, SERUM

PROSTATE SPECIFIC ANTIGEN

0.544

< 1.4

ng/mL

METHOD: ELECTROCHEMILUMINESCENCE, SANDWICH IMMUNOASSAY

Interpretation(s)
PROSTATE SPECIFIC ANTIGEN, SERUM-- PSA is detected in the male patients with normal, benign hyperplastic and malignant prostate tissue and in patients with prostating PROSTATE SPECIFIC ANTIGEN, SERUM-- PSA is not detected (or detected at very low levels) in the patients without prostate tissue (because of radical prostatectomy or cystoprostatectomy) and also in the

- It a suitable marker for monitoring of patients with Prostate Cancer and it is better to be used in conjunction with other diagnostic procedures. female patient.

- It a suitable marker for monitoring or patients with prostate cancer and it is better to be used in conjunction with other diagnostic procedures.

- Serial PSA levels can help determine the success of prostatectomy and the need for further treatment, such as radiation, endocrine or chemotherapy and useful in

oetecong residual disease and early recurrence of tumor.

- Elevated levels of PSA can be also observed in the patients with non-malignant diseases like Prostatitis and Benign Prostatic Hyperplasia.

- Specimens for total PSA assay should be obtained before biopsy, prostatectomy or prostatic massage, since manipulation of the prostate gland may lead to elevated PS/
(false positive) levels persisting up to 3 weeks. (raise positive) levels persisting up to 3 weeks.

- As per American urological guidelines, PSA screening is recommended for early detection of Prostate cancer above the age of 40 years. Following Age specific reference range can be used as a guide lines-

Age of male Reference range (ng/ml)

40-49 years 0-2.5 50-59 years 0-3.5

0-3.5 60-69 years

70-79 years 0-6.5

(* conventional reference level (< 4 ng/ml) is already mentioned in report, which covers all agegroup with 95% prediction interval)

References- Teitz ,textbook of clinical chemiistry, 4th edition) 2.Wallach's Interpretation of Diagnostic Tests

End Of Report

Please visit www.srlworld.com for related Test Information for this accession

Dr. Swapnil Sirmukaddam

Consultant Pathologist

SRL Ltd BHOOMI TOWER, 1ST FLOOR, HALL NO.1, PLOT NO.28 SECTOR 4, KHARGHAR NAVI MUMBAI, 410210 MAHARASHTRA, INDIA Tel: 9111591115,



Scan to View Details



Scan to View Report



Page 2 Of 2







PATIENT NAME: MR. MR.SAMEER WANKHEDE

FH.2341958 PATIENT ID:

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002081

SEX: Male AGE: 36 Years

ABHA NO:

REPORTED :

10/12/2022 14:14:46

DRAWN: 10/12/2022 12:50:00

RECEIVED: 10/12/2022 12:56:10 REFERRING DOCTOR:

CLIENT NAME : FORTIS VASHI-CHC -SPLZD CLINICAL INFORMATION:

UID:2341958 REQNO-1342076

CORP-OPD BILLNO-1501220PCR062807 Units Biological Reference Interval BILLNO-150122OPCR062807 Results <u>Final</u> Test Report Status

BIO CHEMISTRY

GLUCOSE, POST-PRANDIAL, PLASMA


PPBS(POST PRANDIAL BLOOD SUGAR)

137

70 - 139

mg/dL

METHOD: HEXOKINASE



Interpretation(s)
GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin response & sensitivity etc. Additional test HbA1c treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c


Please visit www.srlworld.com for related Test Information for this accession

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA Tel: 022-39199222,022-49723322,



Scan to View Details











PATIENT NAME: MR. MR.SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002081

AGE: 36 Years

SEX: Male

ABHA NO:

REPORTED: 10/12/2022 14:14:46

DRAWN: 10/12/2022 12:50:00

RECEIVED: 10/12/2022 12:56:10

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR:

CLINICAL INFORMATION:

UID:2341958 REQNO-1342076 CORP-OPD BILLNO-150122OPCR062807 BILLNO-150122OPCR062807

Test Report Status

<u>Final</u>

Results

Biological Reference Interval

Units

Dr.Akta Dubey Counsultant Pathologist

SRL Ltd HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA Tel: 022-39199222,022-49723322,



Scan to View Details



Scan to View Report



Ęż FORTIS HIRANANDANIHOSPITAL VASHI 덩 PH100B F 50~ 0.50-100 HZ W 12/10/2022 11:37:08 AM Sinus rhythm.....grs area>0 in V2ST >0.10mV, V1-V4 Unconfirmed Diagnosis Chest: 10.0 mm/mV - BORDERLINE ECG -**V3** V2. Z Borderline T abnormalities, inferior leads..... Limb: 10 mm/mV Minimal ST elevation, anterior leads.. MR. SAMEER WANKHEDE Baseline wander in lead(s) V2,V4 navri ce: Speed: 25 mm/sec aVF aVR 12 Lead; Standard Placement 133 98 368 411 2341958 --AXIS--36 Years III PR QRSD QT QTC Rate QRS T н

Board Line: 022 - 39199222 | Fax: 022 - 39133220

Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199200 | Health Checkup: 022 - 39199300

www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN: 27AABCH5894D1ZG PAN NO: AABCH5894D







Date: 12/Dec/2022

DEPARTMENT OF NIC

UHID | Episode No: 2341958 | 62181/22/1501 Order No | Order Date: 1501/PN/OP/2212/132171 | 10-Dec-2022

Name: Mr. SAMEER WANKHEDE Admitted On | Reporting Date: 12-Dec-2022 10:47:03 Age | Sex: 36 YEAR(S) | Male

Order Doctor Name: Dr.SELF. Order Station: FO-OPD Bed Name:

ECHOCARDIOGRAPHY TRANSTHORACIC

FINDINGS:

- No left ventricle regional wall motion abnormality at rest.
- Normal left ventricle systolic function. LVEF = 60%.
- No left ventricle diastolic dysfunction.
- No left ventricle Hypertrophy. No left ventricle dilatation.
- Structurally normal valves.
- No mitral regurgitation.
- No aortic regurgitation. No aortic stenosis.
- No tricuspid regurgitation. No pulmonary hypertension.
- Intact IAS and IVS.
- No left ventricle clot/vegetation/pericardial effusion.
- Normal right atrium and right ventricle dimensions.
- Normal left atrium and left ventricle dimension.
- Normal right ventricle systolic function. No hepatic congestion

M-MODE MEASUREMENTS:

I-MODE MEASUREMENTS:	37	mm
LA	32	mm
AO Root	23	mm
AO CUSP SEP	30	mm
LVID (s)	44	mm
LVID (d)	10	mm
IVS (d)	10	mm
LVPW (d)	24	mm
RVID (d)	29	mm
RA	60	%

Hiranandani Healthcare Pvt. Ltd. Mini Sea Shore Road, Sector 10-A, Vashi, Navi Mumbai - 400703.

Board Line: 022 - 39199222 | Fax: 022 - 39133220

Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199200 | Health Checkup: 022 - 39199300

www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN: 27AABCH5894D1ZG PAN NO: AABCH5894D







Date: 12/Dec/2022

DEPARTMENT OF NIC

UHID | Episode No: 2341958 | 62181/22/1501 Order No | Order Date: 1501/PN/OP/2212/132171 | 10-Dec-2022 Name: Mr. SAMEER WANKHEDE Admitted On | Reporting Date: 12-Dec-2022 10:47:03

Age | Sex: 36 YEAR(S) | Male

Order Doctor Name: Dr.SELF. Order Station: FO-OPD Bed Name:

DOPPLER STUDY:

E WAVE VELOCITY: 0.9 m/sec. A WAVE VELOCITY:0.5 m/sec

A RATIO:1.4	PEAK (mmHg)	MEAN (mmHg)	V max (m/sec)	GRADE OF REGURGITATION Nil
MITRAL VALVE	N			Nil
AORTIC VALVE	05			Nil
TRICUSPID VALVE PULMONARY VALVE	N 2.0			Nil

Final Impression:

Normal 2 Dimensional and colour doppler echocardiography study.

DR. PRASHANT PAWAR DNB(MED), DNB (CARDIOLOGY)

Hiranandani Healthcare Pvt. Ltd. Mini Sea Shore Road, Sector 10-A, Vashi, Navi Mumbai - 400703.

Board Line: 022 - 39199222 | Fax: 022 - 39133220 Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199200 | Health Checkup: 022 - 39199300

www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN: 27AABCH5894D1ZG PAN NO: AABCH5894D





DEPARTMENT OF RADIOLOGY

Date: 10/Dec/2022

UHID | Episode No : 2341958 | 62181/22/1501 Order No | Order Date: 1501/PN/OP/2212/132171 | 10-Dec-2022 Name: Mr. SAMEER WANKHEDE

Admitted On | Reporting Date: 10-Dec-2022 12:41:31 Age | Sex: 36 YEAR(S) | Male Order Doctor Name : Dr.SELF .

Order Station : FO-OPD Bed Name:

X-RAY-CHEST- PA

Findings:

Both lung fields are clear.

The cardiac shadow appears within normal limits.

Trachea and major bronchi appears normal.

Both costophrenic angles are well maintained.

Bony thorax is unremarkable.

flehal DR. YOGINI SHAH DMRD., DNB. (Radiologist) Hiranandani Healthcare Pvt. Ltd.

Mini Sea Shore Road, Sector 10-A, Vashi, Navi Mumbai - 400703.

Board Line: 022 - 39199222 | Fax: 022 - 39133220 Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199200 | Health Checkup: 022 - 39199300

www.fortishealthcare.com | vashi@fortishealthcare.com

CIN: U85100MH2005PTC 154823 GST IN: 27AABCH5894D1ZG PAN NO: AABCH5894D





DEPARTMENT OF RADIOLOGY

Date: 10/Dec/2022

Name: Mr. SAMEER WANKHEDE

Age | Sex: 36 YEAR(S) | Male

Order Station: FO-OPD

Bed Name:

UHID | Episode No: 2341958 | 62181/22/1501

Order No | Order Date: 1501/PN/OP/2212/132171 | 10-Dec-2022

Admitted On | Reporting Date: 10-Dec-2022 13:06:51

Order Doctor Name: Dr.SELF.

US-WHOLE ABDOMEN

LIVER is normal in size (13.3 cm) and shows raised echogenicity. No IHBR dilatation. No focal lesion is seen in liver. Portal vein appears normal in caliber.

GALL BLADDER is physiologically distended. Gall bladder reveals normal wall thickness. No evidence of calculi in gall bladder. No evidence of pericholecystic collection. CBD appears normal in caliber.

SPLEEN is normal in size (12.2 cm) and echogenicity.

BOTH KIDNEYS are normal in size and echogenicity. The central sinus complex is normal. No evidence of calculi/hydronephrosis.

Right kidney measures 11.6 x 5.1 cm.

Left kidney measures 12.4 x 5.6 cm.

PANCREAS: Head and body of pancreas appears unremarkable. Rest of the pancreas is

URINARY BLADDER is normal in capacity and contour. Bladder wall is normal in thickness. No evidence of intravesical calculi.

PROSTATE is normal in size & echogenicity. It measures ~ 16 cc in volume.

No evidence of ascites.

IMPRESSION:

Grade I fatty infiltration of liver.

DR. VIVER MANE

MBBS., DMRE. (Radiologist)

Hiranandani Healtheare Pvt. Ltd.

Mini Sea Shore Road, Sector 10 -A, Vashi, Navi Mumbai - 400703

Board Line: 022 - 39199222 | Fax: 022 - 39199220 Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199222 | Health Checkup: 022 - 39199300

www.fortishealthcare.com |

CIN: U85100MH2005PTC154823

GST IN: 27AABCH5894D1ZG | PAN NO: AABCH5894D



		Date	10/12/20)22	
UHID	2341958	Sex	Male	Age	36
Name	Mr.Sameer Wankhede	100000000000000000000000000000000000000	1		
1141110		Healt	h Check I) p	
OPD	Opthal 14			_	

Drug allergy: -> Not kown;
Sys illness: -> Wo

Hiranandani Healthcare Pvt. Ltd.

Mini Sea Shore Road, Sector 10 -A, Vashi, Navi Mumbai - 400703

Board Line: 022 - 39199222 | Fax: 022 - 39199220 Emergency: 022 - 39199100 | Ambulance: 1255

For Appointment: 022 - 39199222 | Health Checkup: 022 - 39199300

www.fortishealtheare.com |

CIN: U85100MH2005PTC154823

GST IN: 27AABCH5894D1ZG | PAN NO: AABCH5894D



		Date	10/12/2	022	
UHID	2341958	Sex		Age	36
Name	Mr.Sameer Wankhede	Healtl			
OPD	Dental 12	Heam	II CHECK	-	eriest "' '

No significant findings Adv orel prophy bais.

Drug allergy: Sys illness:







PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

SEX: Male

ABHA NO:

ACCESSION NO: 0022VL002012 DRAWN: 10/12/2022 10:03:00

AGE: 36 Years RECEIVED: 10/12/2022 10:06:43

REPORTED:

10/12/2022 16:27:58

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807

BILLNO-1501220PCR06 BILLNO-1501220PCR06	62807		Biological Reference Interval	Units
	7-4 97 4	Results	Biological Referen	
Test Report Status	<u>Final</u>		*	

KIDNEY PANEL - 1

BLOOD UREA NITROGEN (BUN), SERUM


BLOOD UREA NITROGEN

8

6 - 20

mg/dL

METHOD: UREASE - UV



CREATININE EGFR- EPI

CREATININE

0.81

Low 0.90 - 1.30

mg/dL

AGE

METHOD: ALKALINE PICRATE KINETIC JAFFES

36

117.18

Refer Interpretation Below

years mL/min/1.7

GLOMERULAR FILTRATION RATE (MALE) METHOD: CALCULATED PARAMETER

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SRL Ltd SECTOR 10,


NAVI MUMBAI, 400703 MAHARASHTRA, INDIA



Scan to View Details



Scan to View Report









PATIENT NAME: MR. SAMEER WANKHEDE

FH.2341958 PATIENT ID:

CLIENT PATIENT ID: UID:2341958

DRAWN: 10/12/2022 10:03:00

ACCESSION NO: 0022VL002012

SEX: Male AGE: 36 Years RECEIVED: 10/12/2022 10:06:43

Results

ABHA NO:

10/12/2022 16:27:58 REPORTED:

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:


UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status Final **Biological Reference Interval**

Units



ੈ 0.46 – ਵਿੱ			
	SEP-2018	10-DEC-2022	
Biological Reference Interval: 0.90 - 1.30	mg/dl Date	>	
BUN/CREAT RATIO	9.88	5.00 - 15.00	
BUN/CREAT RATIO	5.00		
METHOD: CALCULATED PARAMETER			a / d l
URIC ACID, SERUM	5.8	3.5 - 7.2	mg/dL
URIC ACID			
METHOD : URICASE UV			
TOTAL PROTEIN, SERUM	7.3	6.4 - 8.2	g/dL
TOTAL PROTEIN	800		
METHOD: BIURET			X-31
ALBUMIN, SERUM	4.2	3.4 - 5.0	g/dL
ALBUMIN			
METHOD: BCP DYE BINDING			9-77-00
GLOBULIN	3.1	2.0 - 4.1	g/dL
GLOBULIN	3.1		
METHOD : CALCULATED PARAMETER			
ELECTROLYTES (NA/K/CL), SERUM	140	136 - 145	mmol/L
SODIUM, SERUM	110		
METHOD: ISE INDIRECT	4.52	3.50 - 5.10	mmol/L

4.52

METHOD: ISE INDIRECT POTASSIUM, SERUM

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD,

SECTOR 10, NAVI MUMBAI, 400703

MAHARASHTRA, INDIA Tel: 022-39199222,022-49723322,



Scan to View Details



Scan to View Report









PATIENT NAME: MR. SAMEER WANKHEDE

FH.2341958 PATIENT ID:

CLIENT PATIENT ID: UID:2341958

ACCESSION NO: 0022VL002012

SEX: Male AGE: 36 Years RECEIVED: 10/12/2022 10:06:43 ABHA NO: REPORTED:

10/12/2022 16:27:58

DRAWN: 10/12/2022 10:03:00

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807 Test Report Status Final	Results	Biological Reference Interval	Units
		m	mol/L

METHOD: ISE INDIRECT CHLORIDE, SERUM METHOD : ISE INDIRECT

Interpretation(s)

103

98 - 107

mmol/L

PHYSICAL EXAMINATION, URINE

PALE YELLOW

METHOD: PHYSICAL **APPEARANCE**

CLEAR

METHOD : VISUAL

CHEMICAL EXAMINATION, URINE

60

4.7 - 7.5

METHOD: REFLECTANCE SPECTROPHOTOMETRY- DOUBLE INDICATOR METHOD

1.003 - 1.035

<=1.005 METHOD: REFLECTANCE SPECTROPHOTOMETRY (APPARENT PKA CHANGE OF PRETREATED POLYELECTROLYTES IN RELATION TO IONIC CONCENTRATION) SPECIFIC GRAVITY NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY - PROTEIN-ERROR-OF-INDICATOR PRINCIPLE

GLUCOSE

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, DOUBLE SEQUENTIAL ENZYME REACTION-GOD/POD

NOT DETECTED

NOT DETECTED

KETONES METHOD: REFLECTANCE SPECTROPHOTOMETRY, ROTHERA'S PRINCIPLE

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, PEROXIDASE LIKE ACTIVITY OF HAEMOGLOBIN

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, DIAZOTIZATION- COUPLING OF BILIRUBIN WITH DIAZOTIZED SALT BILIRUBIN

UROBILINOGEN

NORMAL

NORMAL

METHOD: REFLECTANCE SPECTROPHOTOMETRY (MODIFIED EHRLICH REACTION)

NITRITE

NOT DETECTED

NOT DETECTED

METHOD: REFLECTANCE SPECTROPHOTOMETRY, CONVERSION OF NITRATE TO NITRITE

NOT DETECTED

NOT DETECTED

LEUKOCYTE ESTERASE METHOD: REFLECTANCE SPECTROPHOTOMETRY, ESTERASE HYDROLYSIS ACTIVITY

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS

NOT DETECTED

NOT DETECTED

/HPF

METHOD: MICROSCOPIC EXAMINATION

PUS CELL (WBC'S)

1-2

0-5

/HPF

METHOD: MICROSCOPIC EXAMINATION

SRL Ltd

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD,


SECTOR 10,

NAVI MUMBAI, 400703 MAHARASHTRA, INDIA

Scan to View Details



Scan to View Report









PATIENT NAME: MR. SAMEER WANKHEDE

CLIENT PATIENT ID: UID:2341958 FH.2341958 PATIENT ID:

SEX: Male

ABHA NO: 10/12/2022 16:27:58 36 Years AGE: 0022VL002012 REPORTED: RECEIVED: 10/12/2022 10:06:43 ACCESSION NO: DRAWN: 10/12/2022 10:03:00

REFERRING DOCTOR: SELF CLIENT NAME : FORTIS VASHI-CHC -SPLZD

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807

Results	Biological Reference I	nterval Uni	
Vezairo			
		/HPF	
0-1	0-5	\underset \under	
NOT DETECTED			
NOT DETECTED			
	NOT DETECTED		
	NOT DETECTED		
IIDINARY MICROSCO	PIC EXAMINATION DONE ON U	JRINARY	
CENTRIFUGED SEDI	MENT		
	NOT DETECTED NOT DETECTED NOT DETECTED NOT DETECTED	NOT DETECTED NOT DETECTED NOT DETECTED NOT DETECTED	

Interpretation(s)

Interpretation(s)
BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, BLOOD UREA NITROGEN (BUN), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)
Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)
Causes of decreased level include Liver disease, SIADH.
CREATININE EGFR- EPICREATININE EGFR- EPICR

CREATININE EGFR- EPIGFR— Glomerular filtration rate (GFR) is a measure of the function of the kidneys. The GFR is a calculation based on a serum creatinine test. Creatinine is a muscle waste product that is filtered from the blood by the kidneys and excreted into urine at a relatively steady rate. When kidney function decreases, less creatinine is excreted and concentrations increase in the blood. With the creatinine test, a reasonable estimate of the actual GFR can be determined.

A GFR of 60 or higher is in the normal range.

A GFR below 60 may mean kidney disease.

A GFR below 60 may mean kidney disease.

A GFR of 15 or lower may mean kidney failure.

Estimated GFR (eGFR) is the preferred method for identifying people with chronic kidney disease (CKD). In adults, eGFR calculated using the Modification of Diet in Renal Estimated GFR (eGFR) is the preferred method for identifying people with chronic kidney function than serum creatinine alone.

Disease (MDRD) Study equation provides a more clinically useful measure of kidney function than serum creatinine alone.

The CKD-EPI creatinine equation is based on the same four variables as the MDRD Study equation, but uses a 2-slope spline to model the relationship between estimated than CKD-EPI creatinine equation is based on the same four variables as the MDRD Study equation was reported to perform better and with less bias than the MDRD Study equation as recorded in the contract of CKD especially in patients with higher GFR. This results in reduced misclassification of CKD.

ESPECIALLY IN THE CKD-EPI creatinine equation has not been validated in children & will only be reported for patients = 18 years of age. For pediatric and childrens, Schwartz Pediatric EVCD-EPI creatinine equation has not been validated in children & will only be reported for patients and height.

Bedside eGFR (2009) formulae is used. This revised "bedside" pediatric eGFR requires only serum creatinine and height.

the CKD-EPI creatinine equation has not been valuated in children & will only be reported for patients — 16 years of age Bedside eGFR (2009) formulae is used. This revised "bedside" pediatric eGFR requires only serum creatinine and height.

Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciencesis
Cause of the Cause of

TOTAL PROTEIN, SERUM-Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Higner-trian-normal levels may be due to: Circuit inflammation of infection, including fits and negatives both c, prompte myeloma, wateristion, Malnutrition, Nephrotic Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic

ALBUMIN, SERUMHuman serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703


MAHARASHTRA, INDIA Tel: 022-39199222,022-49723322,



Scan to View Details



Scan to View Report



Page 4 Of 15

Patient Ref. No. 220000001







PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO: DRAWN: 10/12/2022 10:03:00

0022VL002012

SEX: Male 36 Years AGE: RECEIVED: 10/12/2022 10:06:43 ABHA NO: REPORTED:

10/12/2022 16:27:58

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status

Results

Biological Reference Interval

HAEMATOLOGY

CBC-5, EDTA WHOLE BLOOD

MORPHOLOGY

RBC

PREDOMINANTLY NORMOCYTIC NORMOCHROMIC

METHOD: MICROSCOPIC EXAMINATION

WBC

METHOD: MICROSCOPIC EXAMINATION

PLATELETS

METHOD: MICROSCOPIC EXAMINATION

ADEQUATE

NORMAL MORPHOLOGY

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD

E.S.R

METHOD: WESTERGREN METHOD

0 - 14

mm at 1 hr

CBC-5, EDTA WHOLE BLOOD

BLOOD COUNTS, EDTA WHOLE BLOOD

HEMOGLOBIN (HB) METHOD: SPECTROPHOTOMETRY

15.2 4.98

03

13.0 - 17.0

g/dL mil/µL

RED BLOOD CELL (RBC) COUNT METHOD: ELECTRICAL IMPEDANCE

5.16

4.5 - 5.54.0 - 10.0

thou/µL

WHITE BLOOD CELL (WBC) COUNT METHOD: DOUBLE HYDRODYNAMIC SEQUENTIAL SYSTEM(DHSS)CYTOMETRY

218

150 - 410

thou/µL

PLATELET COUNT METHOD: ELECTRICAL IMPEDANCE **RBC AND PLATELET INDICES**

HEMATOCRIT (PCV)

44.8

40 - 50

%

METHOD: CALCULATED PARAMETER

90.1

83 - 101

fL

MEAN CORPUSCULAR VOLUME (MCV) METHOD: CALCULATED PARAMETER MEAN CORPUSCULAR HEMOGLOBIN (MCH)

30.6

27.0 - 32.0

pg

METHOD: CALCULATED PARAMETER MEAN CORPUSCULAR HEMOGLOBIN

33.9

31.5 - 34.5

g/dL

CONCENTRATION(MCHC) METHOD: CALCULATED PARAMETER

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10,

NAVI MUMBAI, 400703 MAHARASHTRA, INDIA













PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002012

AGE: 36 Years

SEX: Male

ABHA NO:

REPORTED:

10/12/2022 16:27:58

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43 REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

BILLNO-1501220PCR062807		Biological Reference	Biological Reference Interval		
Test Report Status <u>Final</u>	Results				
	4.4.7	High 11.6 - 14.0	%		
RED CELL DISTRIBUTION WIDTH (RDW)	14.3				
METHOD: CALCULATED PARAMETER	18.1				
MENTZER INDEX	10.6	6.8 - 10.9	fL		
MEAN PLATELET VOLUME (MPV)	10.0				
METHOD: CALCULATED PARAMETER					
WBC DIFFERENTIAL COUNT	C 4	40 - 80	%		
NEUTROPHILS	64				
METHOD: FLOW CYTOMETRY	28	20 - 40	%		
LYMPHOCYTES	20		*		
METHOD: FLOW CYTOMETRY	5	2 - 10	%		
MONOCYTES	3				
METHOD: FLOW CYTOMETRY	03	1 - 6	%		
EOSINOPHILS	103				
METHOD: FLOW CYTOMETRY	00	0 - 2	%		
BASOPHILS	00				
METHOD : FLOW CYTOMETRY	3.30	2.0 - 7.0	thou/µL		
ABSOLUTE NEUTROPHIL COUNT	5.50				
METHOD : CALCULATED PARAMETER	1.44	1.0 - 3.0	thou/µL		
ABSOLUTE LYMPHOCYTE COUNT	*****				
METHOD : CALCULATED PARAMETER	0.26	0.2 - 1.0	thou/µL		
ABSOLUTE MONOCYTE COUNT	9.22		99990 BL W		
METHOD : CALCULATED PARAMETER	0.15	0.02 - 0.50	thou/µL		
ABSOLUTE EOSINOPHIL COUNT	ARCHES		yug nerel		
METHOD : CALCULATED PARAMETER	0	Low 0.02 - 0.10	thou/μL		
ABSOLUTE BASOPHIL COUNT					
METHOD: CALCULATED PARAMETER NEUTROPHIL LYMPHOCYTE RATIO (NLR)	2.3				
NEUTROPHIL LYMPHOCYTE RATIO (NEIX)					

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION:

Erythrocyte Sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA

Tel: 022-39199222,022-49723322,

METHOD: CALCULATED PARAMETER



Scan to View Details



Scan to View Report









PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002012

SEX: Male 36 Years AGE: RECEIVED: 10/12/2022 10:06:43 ABHA NO:

REPORTED:

10/12/2022 16:27:58

DRAWN: 10/12/2022 10:03:00 CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status

Final

Results

Biological Reference Interval

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, ringing a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS
False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia
False elevated ESR: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, False Decreased: Poikilocytosis, Counts,

REFERENCE:

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with elicited carealables and deficiency anaemia. This needs to be interpreted in line with elicited carealables and deficiency anaemia.

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical symptoms to change from mild to severe in COVID positive diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.3.

2.3, COVID-19 patients tend to show thind disease.
(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

IMMUNOHAEMATOLOGY

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP

TYPE O

METHOD: TUBE AGGLUTINATION

RH TYPE

POSITIVE

METHOD: TUBE AGGLUTINATION

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

BIO CHEMISTRY

LIPID PROFILE, SERUM

CHOLESTEROL, TOTAL

212

High < 200 Desirable

mg/dL

mg/dL

200 - 239 Borderline High

>/= 240 High

METHOD: ENZYMATIC/COLORIMETRIC, CHOLESTEROL OXIDASE, ESTERASE, PEROXIDASE 186

TRIGLYCERIDES

High < 150 Normal

150 - 199 Borderline High

200 - 499 High

>/=500 Very High

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10,

NAVI MUMBAI, 400703 MAHARASHTRA, INDIA



Scan to View Details



Scan to View Report









PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID : FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO: 0022VL002012 AGE: 36 Years

SEX: Male

ABHA NO:

10/12/2022 16:27:58

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status	Final	Results		Biological Reference Interva	d
Test Report Status	Liliai				
METHOD: ENZYMATIC ASSA	AY			40.1	mg/dL
HDL CHOLESTEROL		46		< 40 Low >/=60 High	mg/ac
METHOD: DIRECT MEASUR	E - PEG			31	/dl
LDL CHOLESTEROL, D	IRECT	132	High	< 100 Optimal 100 - 129 Near or above optim 130 - 159 Borderline High 160 - 189 High >/= 190 Very High	mg/dL al
METHOD: DIRECT MEASUR	E WITHOUT SAMPLE PRETRI	EATMENT			
NON HDL CHOLESTER	OL	166	High	Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
METHOD: CALCULATED PA	RAMETER			8	
CHOL/HDL RATIO		4.6	High	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk	
METHOD: CALCULATED PA	RAMETER				
LDL/HDL RATIO		2.9		0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate >6.0 High Risk	Risk
METHOD: CALCULATED PA	RAMETER				50 - SRH1
VERY LOW DENSITY L	IPOPROTEIN	37.2	High	= 30.0</td <td>mg/dL</td>	mg/dL

SRL LtdHIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703

MAHARASHTRA, INDIA

METHOD: CALCULATED PARAMETER







Scan to View Report







PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002012

36 Years AGE:

SEX: Male

ABHA NO:

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

10/12/2022 16:27:58

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD


BILLNO-1501220PCR062807 BILLNO-1501220PCR062807


Test Report Status

Final

Results

Biological Reference Interval






SRL Ltd HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA



Scan to View Details



Scan to View Report









PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID: FH.2341958 CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002012

36 Years AGE:

SEX: Male

ABHA NO:

10/12/2022 16:27:58

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

Final

CORP-OPD


BILLNO-1501220PCR062807


BILLNO-1501220PCR062807

Test Report Status

Results

Biological Reference Interval





LIVER FUNCTION PROFILE, SERUM


HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA



Scan to View Details



Scan to View Report









PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID : FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO: 0022VL002012

AGE: 36 Years

SEX: Male

ABHA NO:

10/12/2022 16:27:58

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status <u>Final</u>	Results		Biological Referen	ce Interval
BILIRUBIN, TOTAL	0.76		0.2 - 1.0	mg/dL
METHOD: JENDRASSIK AND GROFF				
BILIRUBIN, DIRECT	0.19		0.0 - 0.2	mg/dL
METHOD: JENDRASSIK AND GROFF	5			
BILIRUBIN, INDIRECT	0.57		0.1 - 1.0	mg/dL
METHOD: CALCULATED PARAMETER				2021
TOTAL PROTEIN	7.3		6.4 - 8.2	g/dL
METHOD: BIURET				
ALBUMIN	4.2		3.4 - 5.0	g/dL
METHOD: BCP DYE BINDING				
GLOBULIN	3.1		2.0 - 4.1	g/dL
METHOD: CALCULATED PARAMETER				
ALBUMIN/GLOBULIN RATIO	1.4		1.0 - 2.1	RATIO
METHOD: CALCULATED PARAMETER				
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	49	High	15 - 37	U/L
METHOD: UV WITH P5P	1272 21	***	- 0285-00-00-0	
ALANINE AMINOTRANSFERASE (ALT/SGPT)	114	High	< 45.0	U/L
METHOD : UV WITH P5P				2227
ALKALINE PHOSPHATASE	61		30 - 120	U/L
METHOD : PNPP-ANP	72		15 05	7770
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: GAMMA GLUTAMYLCARBOXY 4NITROANILIDE	73		15 - 85	U/L
LACTATE DEHYDROGENASE	164		100 100	1176
METHOD : LACTATE -PYRUVATE	104		100 - 190	U/L
DELINO - DELINE-PIROVAIE				
GLUCOSE FASTING, FLUORIDE PLASMA				
FBS (FASTING BLOOD SUGAR)	88		74 - 99	mg/dL
METHOD : HEXOKINASE				mg/ac


SRL Ltd HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA

Tel: 022-39199222,022-49723322,



Scan to View Details











PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID: FH.2341958 CLIENT PATIENT ID: UID:2341958

ACCESSION NO: 0022VL002012

AGE: 36 Years SEX: Male

ABHA NO:

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

10/12/2022 16:27:58

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD


BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status

Final

Results

Biological Reference Interval



GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD

HBA1C

5.0

Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4

Diabetics: > or = 6.5Therapeutic goals: < 7.0 Action suggested: > 8.0

(ADA Guideline 2021)

METHOD: HB VARIANT (HPLC)

METHOD: CALCULATED PARAMETER

ESTIMATED AVERAGE GLUCOSE(EAG)

96.8

< 116.0

mg/dL

%


SRL Ltd HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA

Tel: 022-39199222,022-49723322,



Scan to View Details











PATIENT NAME: MR. SAMEER WANKHEDE

FH.2341958 PATIENT ID:

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002012

36 Years AGE :

SEX: Male

ABHA NO:

10/12/2022 16:27:58

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD


BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status

Final

Results

Biological Reference Interval



NOTE: RESULTS OBTAINED ON REPEAT ANALYSIS (S -WINDOW WITH RETENTION TIME 1.66 AND AREA 37.4 %). THIS COULD BE PROBABLY DUE TO INTERFERENCE BY PRESENCE OF ABNORMAL HEMOGLOBIN VARIANTS. ADVISED HEMOGLOBIN VARIANT STUDY FOR THE SAME.

Interpretation(s)
LIPID PROFILE, SERUM-Serum cholesterol is a blood test that can provide valuable information for the risk of coronary artery disease This test can help determine your risk of the build up of plaques in your arteries that can lead to narrowed or blocked arteries throughout your body (atherosclerosis). High cholesterol levels usually don cause any signs or symptoms, so a cholesterol test is an important tool. High cholesterol levels often are a significant risk factor for heart disease and important for diagnosis of hyperlipoproteinemia, atherosclerosis, hepatic and thyroid diseases.

Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn""""t need into triglycerides, which are stored in fat cells. High triglyceride levels are associated with several factors, including being overweight, eating too many sweets or drinking too much alcohol, smoking, being sedentary, or having diabetes with elevated blood sugar levels. Analysis has proven useful in the diagnosis and treatment of patients with diabetes mellitus, nephrosis, liver obstruction, other diseases involving lipid metabolism, and various endocrine disorders. In conjunction with high density lipoprotein and total serum cholesterol, a triglyceride determination of the provides valuable information for the acceptance of conspany levels diseased involved. Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn' provides valuable information for the assessment of coronary heart disease risk. It is done in fasting state.

High-density lipoprotein (HDL) cholesterol. This is sometimes called the ""good"" cholesterol because it helps carry away LDL cholesterol, thus keeping arteries open and blood flowing more freely. HDL cholesterol is inversely related to the risk for cardiovascular disease. It increases following regular exercise, moderate alcohol consumption and with oral estrogen therapy. Decreased levels are associated with obesity, stress, cigarette smoking and diabetes mellitus.

SERUM LDL The small dense LDL test can be used to determine cardiovascular risk in individuals with metabolic syndrome or established/progressing coronary artery disease, individuals with triglyceride levels between 70 and 140 mg/dL, as well as individuals with a diet high in trans-fat or carbohydrates. Elevated sdLDL levels are associated with metabolic syndrome and an 'atherogenic lipoprotein profile', and are a strong, independent predictor of cardiovascular disease. Elevated levels of LDL arise from multiple sources. A major factor is sedentary lifestyle with a diet high in saturated fat. Insulin-resistance and pre-diabetes have also been implicated, as has genetic predisposition. Measurement of sdLDL allows the clinician to get a more comprehensive picture of lipid risk factors and tailor treatment accordingly. Reducing LDL levels will reduce the risk of CVD and MI.

Non HDL Cholesterol - Adult treatment panel ATP III suggested the addition of Non-HDL Cholesterol as an indicator of all atherogenic lipoproteins (mainly LDL and VLDL), NICE guidelines recommend Non-HDL Cholesterol measurement before initiating lipid lowering therapy. It has also been shown to be a better marker of risk in both primary and secondary prevention studies.

SRL Ltd

HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10,

NAVI MUMBAI, 400703 MAHARASHTRA, INDIA

Tel: 022-39199222,022-49723322,

Scan to View Details











PATIENT NAME: MR. SAMEER WANKHEDE

CLIENT PATIENT ID: UID:2341958 PATIENT ID: FH.2341958

0022VL002012 36 Years SEX: Male ABHA NO: ACCESSION NO: AGE :

10/12/2022 16:27:58 RECEIVED: 10/12/2022 10:06:43 REPORTED : DRAWN: 10/12/2022 10:03:00

REFERRING DOCTOR: SELF CLIENT NAME : FORTIS VASHI-CHC -SPLZD

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Results **Biological Reference Interval Test Report Status Final**

Recommendations:

Results of Lipids should always be interpreted in conjunction with the patient's medical history, clinical presentation and other findings.

NON FASTING LIPID PROFILE includes Total Cholesterol, HDL Cholesterol and calculated non-HDL Cholesterol. It does not include triglycerides and may be best used in patients for whom fasting is difficult.

LIVER FUNCTION PROFILE, SERUM-

LIVER FUNCTION PROFILE

ELIVER FONCTION PROFILE

Billirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Billirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg., hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg., obstruction and hepatitis), and abnormal bilirubin metabolism (eg., hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatifits, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Billary obstruction, ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin level

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

Increased in

Diabetes mellitus, Cushing' s syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides. Decreased in

Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency, hypopituitarism,diffuse liver disease, malignancy (adrenocortical, stomach,fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia),Drugs- insulin, ethanol, propranolol; sulfonylureas,tolbutamide, and other oral hypoglycemic agents.

NOTE:

While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus,

while random serum glucose levels correlate with nome glucose monitoring results (weekly mean capillary glucose values), there is wide nucleation within molviduals. However, and glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:

- Evaluating the long-term control of blood glucose concentrations in diabetic patients.

1.Evaluating the long-term control of blood glucose concentrations in diabetic patients.
2.Diagnosing diabetes.
3.Identifying patients at increased risk for diabetes (prediabetes).
The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.
1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
2. eAG gives an evaluation of blood glucose levels for the last couple of months.
3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c - 46.7

HbA1c Estimation can get affected due to :

I.Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II.Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

III.Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism,chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.


HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA

Tel: 022-39199222,022-49723322,



Scan to View Details











PATIENT NAME: MR. SAMEER WANKHEDE

PATIENT ID:

FH.2341958

CLIENT PATIENT ID: UID:2341958

ACCESSION NO:

0022VL002012

AGE: 36 Years

SEX: Male

ABHA NO:

DRAWN: 10/12/2022 10:03:00

RECEIVED: 10/12/2022 10:06:43

REPORTED:

10/12/2022 16:27:58

CLIENT NAME : FORTIS VASHI-CHC -SPLZD

REFERRING DOCTOR: SELF

CLINICAL INFORMATION:

UID:2341958 OLD UHID -FHL34.215432 REQNO-1342076

CORP-OPD

BILLNO-1501220PCR062807 BILLNO-1501220PCR062807

Test Report Status

Final

Results

Biological Reference Interval

IV.Interference of hemoglobinopathies in HbA1c estimation is seen in

1V.Interference or nemoglobinopathies in HDA1c estimation is seen in a.Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b.Heterozygous state detected (D10 is corrected for HbS & HbC trait.) c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

End Of Report

Please visit www.srlworld.com for related Test Information for this accession

Dr.Akta Dubey

Counsultant Pathologist

Dr. Rekha Nair, MD

Microbiologist

SRL Ltd HIRANANDANI HOSPITAL-VASHI, MINI SEASHORE ROAD, SECTOR 10, NAVI MUMBAI, 400703 MAHARASHTRA, INDIA

Tel: 022-39199222,022-49723322,



Scan to View Details



