PID No.
 : MED121894715
 Register On
 : 27/05/2023 9:04 AM

 SID No.
 : 522308443
 Collection On
 : 27/05/2023 10:55 AM

 Age / Sex
 : 23 Year(s) / Female
 Report On
 : 27/05/2023 5:42 PM

Type : OP Printed On : 29/05/2023 9:47 AM

Ref. Dr : MediWheel



- i Mediwheel			
Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination)	'B' 'Positive'		
INTERPRETATION: Note: Slide method is screen	ing method. Kindly con	firm with Tube me	ethod for transfusion.
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	12.5	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	38.0	%	37 - 47

(EDIA Blood Spectropnotometry)			
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	38.0	%	37 - 47
RBC Count (EDTA Blood)	4.91	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood)	77.5	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	25.5	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.0	g/dL	32 - 36
RDW-CV	14.1	%	11.5 - 16.0
RDW-SD	38.25	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	4800	cells/cu.m m	4000 - 11000
Neutrophils (Blood)	58.2	%	40 - 75
Lymphocytes (Blood)	35.7	%	20 - 45
Eosinophils (Blood)	0.3	%	01 - 06





 PID No.
 : MED121894715
 Register On
 : 27/05/2023 9:04 AM

 SID No.
 : 522308443
 Collection On
 : 27/05/2023 10:55 AM

 Age / Sex
 : 23 Year(s) / Female
 Report On
 : 27/05/2023 5:42 PM

Type : OP Printed On : 29/05/2023 9:47 AM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
Monocytes (Blood)	5.6	%	01 - 10
Basophils (Blood)	0.2	%	00 - 02
INTERPRETATION: Tests done on Automated Fi	ive Part cell counter. All	abnormal results are r	reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	2.79	$10^3 / \mu l$	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	1.71	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.01	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.27	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.01	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	173	$10^3 / \mu l$	150 - 450
MPV (Blood)	10.1	fL	8.0 - 13.3
PCT (Automated Blood cell Counter)	0.17	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	13	mm/hr	< 20
BUN / Creatinine Ratio	12.2		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	75.66	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.





 PID No.
 : MED121894715
 Register On
 : 27/05/2023 9:04 AM

 SID No.
 : 522308443
 Collection On
 : 27/05/2023 10:55 AM

Type : OP Printed On : 29/05/2023 9:47 AM

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	100.37	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	6.2	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.51	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/Enzymatic)	2.14	mg/dL	2.6 - 6.0
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.25	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.12	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.13	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	34.11	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	28.74	U/L	5 - 41





APPROVED BY

The results pertain to sample tested.

Page 3 of 8

Lab Address: BANGALORE REFERENCE LABORATORY-CLUMAX DIAGNOSTIC AND RESEARCH CENTRE PVT LTD. (A UNIT OF MEDALL HEALTHCARE PVT. LTD.),Old No 66 & New No 1, 2nd Main Road, Bashyam Circle, Sankey Cross Road Malleswaram, Gayathri Devi Park Extension, Kodandarampura, Sadashiva Nagar, Bengaluru, Karnataka-560003,.

PID No. Register On : MED121894715 : 27/05/2023 9:04 AM

: 522308443 SID No. Collection On : 27/05/2023 10:55 AM

Age / Sex : 23 Year(s) / Female Report On 27/05/2023 5:42 PM **Type**

Ref. Dr : MediWheel

: OP **Printed On** : 29/05/2023 9:47 AM



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	7.44	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	86.0	U/L	42 - 98
Total Protein (Serum/Biuret)	6.32	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.91	gm/dl	3.5 - 5.2
Globulin (Serum/ <i>Derived</i>)	2.41	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.62		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	138.58	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	84.34	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

29.36 Optimal(Negative Risk Factor): >= 60 **HDL Cholesterol** mg/dL Borderline: 50 - 59 (Serum/Immunoinhibition) High Risk: < 50





PID No. : MED121894715 Register On : 27/05/2023 9:04 AM SID No.

: 522308443 Collection On : 27/05/2023 10:55 AM

Age / Sex : 23 Year(s) / Female Report On : 27/05/2023 5:42 PM **Type** : OP **Printed On** : 29/05/2023 9:47 AM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
LDL Cholesterol (Serum/Calculated)	92.3	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	16.9	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	109.2	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	4.7	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	2.9	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	3.1	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

Glycosylated Haemoglobin (HbA1c)





 PID No.
 : MED121894715
 Register On
 : 27/05/2023 9:04 AM

 SID No.
 : 522308443
 Collection On
 : 27/05/2023 10:55 AM

Type : OP Printed On : 29/05/2023 9:47 AM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HbA1C	5.7	%	Normal: 4.5 - 5.6
(Whole Blood/HPLC)			Prediabetes: 5.7 - 6.4
			Diabetic: \geq 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 116.89 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.48 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 9.99 $\mu g/dl$ 4.2 - 12.0

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) $0.882 \, \mu IU/mL \, 0.35 - 5.50$

(Serum/ECLIA)





 PID No.
 : MED121894715
 Register On
 : 27/05/2023 9:04 AM

 SID No.
 : 522308443
 Collection On
 : 27/05/2023 10:55 AM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
_	<u>Value</u>		Reference Interval

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment

Colour

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

Pale yellow

- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

URINE ROUTINE

PHYSICAL EXAMINATION (URINE COMPLETE)

(Urine)		
Appearance	Clear	Clear
(Urine)		
Volume(CLU)	35	
(Urine)		
CHEMICAL EXAMINATION (URINE		
<u>COMPLETE)</u>		
pH	7	4.5 - 8.0
(Urine)		
Specific Gravity	1.007	1.002 - 1.035
(Urine)		
Ketone	Negative	Negative
(Urine)		
Urobilinogen	Normal	Normal
(Urine)		





Yellow to Amber

APPROVED BY

The results pertain to sample tested.

Page 7 of 8

Lab Address: BANGALORE REFERENCE LABORATORY-CLUMAX DIAGNOSTIC AND RESEARCH CENTRE PVT LTD. (A UNIT OF MEDALL HEALTHCARE PVT. LTD.),Old No 66 & New No 1, 2nd Main Road, Bashyam Circle, Sankey Cross Road Malleswaram, Gayathri Devi Park Extension, Kodandarampura, Sadashiva Nagar, Bengaluru, Karnataka-560003,.

 PID No.
 : MED121894715
 Register On : 27/05/2023 9:04 AM

 SID No.
 : 522308443
 Collection On : 27/05/2023 10:55 AM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Blood (Urine)	Negative		Negative
Nitrite (Urine)	Negative		Negative
Bilirubin (Urine)	Negative		Negative
Protein (Urine)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Leukocytes(CP) (Urine)	Negative		
<u>MICROSCOPIC EXAMINATION</u> (<u>URINE COMPLETE</u>)			
Pus Cells (Urine)	0-1	/hpf	NIL
Epithelial Cells (Urine)	0-1	/hpf	NIL
RBCs (Urine)	NIL	/HPF	NIL
Others (Urine)	NIL		
INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.			
Casts (Urine)	NIL	/hpf	NIL
Crystals	NIL	/hpf	NIL





APPROVED BY

-- End of Report --

The results pertain to sample tested.

(Urine)

Page 8 of 8

Lab Address: BANGALORE REFERENCE LABORATORY-CLUMAX DIAGNOSTIC AND RESEARCH CENTRE PVT LTD. (A UNIT OF MEDALL HEALTHCARE PVT. LTD.),Old No 66 & New No 1, 2nd Main Road, Bashyam Circle, Sankey Cross Road Malleswaram, Gayathri Devi Park Extension, Kodandarampura, Sadashiva Nagar, Bengaluru, Karnataka-560003,.