		TEST REPORT				
Reg. No	: 2107102624		Reg. Date : 24-Jul-2021			
Name	: Ratan Singh		Collected On : 24-Jul-2021 10:27			
Age/Sex	: 55 Years / Male		Approved On : 24-Jul-2021 11:22			
Ref. By	:		Printed On : 24-Jul-2021 11:46			
Client	: MEDIWHEEL WELLNESS					
Paramete	<u>er</u>	Result				
BLOOD GROUP & RH Specimen: EDTA and Serum; Method: Haemagglutination						
ABO		'A'				
Rh (D)		Positive				
End Of Report						

 Reg. No
 : 2107102624

 Name
 : Ratan Singh

 Age/Sex
 : 55 Years / Male

Collected On : 24-Jul-2021 10:27 **Approved On** : 24-Jul-2021 11:22

Reg. Date

Ref. By

Client : MEDIWHEEL WELLNESS

Printed On : 24-Jul-2021 11:46

: 24-Jul-2021

<u>Parameter</u>	Result	<u>Unit</u>	Reference Interval			
COMPLETE BLOOD COUNT (CBC)						
SPECIMEN: EDTA BLOOD						
Hemoglobin	14.3	g/dL	13.0 - 17.0			
RBC Count	5.04	million/cmm	4.5 - 5.5			
Hematrocrit (PCV)	43.0	%	40 - 54			
MCH	28.4	Pg	27 - 32			
MCV	85.3	fL	83 - 101			
MCHC	33.3	%	31.5 - 34.5			
RDW	12.7	%	11.5 - 14.5			
WBC Count	5230	/cmm	4000 - 11000			
DIFFERENTIAL WBC COUNT (Flow	cytometry)					
Neutrophils (%)	64	%	38 - 70			
Lymphocytes (%)	30	%	20 - 40			
Monocytes (%)	04	%	2 - 8			
Eosinophils (%)	02	%	0 - 6			
Basophils (%)	00	%	0 - 2			
Neutrophils	3347	/cmm				
Lymphocytes	1569	/cmm				
Monocytes	209	/cmm				
Eosinophils	105	/cmm				
Basophils	0	/cmm				
Platelet Count (Flow cytometry)	185000	/cmm	150000 - 450000			
MPV	11.0	fL	7.5 - 11.5			
ERYTHROCYTE SEDIMENTATION I	ERYTHROCYTE SEDIMENTATION RATE					
ESR (After 1 hour)	12	mm/hr	0 - 14			
Modified Westergren Method						

----- End Of Report -----



: 2107102624 Reg. No

Name : Ratan Singh Age/Sex : 55 Years / Male

Ref. By

Client : MEDIWHEEL WELLNESS Reg. Date : 24-Jul-2021

Collected On : 24-Jul-2021 10:27 Approved On : 24-Jul-2021 11:20

Printed On : 24-Jul-2021 11:46

<u>Unit</u> Reference Interval **Parameter** Result

PLASMA GLUCOSE

Fasting Blood Sugar (FBS)

93.2

mg/dL

70 - 110

Hexokinase Method

Criteria for the diagnosis of diabetes1. HbA1c >/= 6.5 *

Or 2. Fasting plasma glucose >126 gm/dL. Fasting is defined as no caloric intake at least for 8 hrs.

Or

3. Two hour plasma glucose >/= 200mg/dL during an oral glucose tolerence test by using a glucose load containing equivalent of 75 gm anhydrous glucose dissolved in water. Or

4. In a patient with classic symptoms of hyperglycemia or hyperglycemic crisis, a random plasma glucose >/= 200 mg/dL. *In the absence of unequivocal hyperglycemia, criteria 1-3 should be confirmed by repeat testing.

American diabetes association. Standards of medical care in diabetes 2011. Diabetes care 2011;34;S11.

----- End Of Report -----

Reg. No : 2107102624 Name : Ratan Singh : 55 Years / Male Age/Sex

Collected On : 24-Jul-2021 10:27 Approved On : 24-Jul-2021 11:20

: 24-Jul-2021

Reg. Date

Ref. By

Printed On : 24-Jul-2021 11:46

Client : MEDIWHEEL WELLNESS

<u>Parameter</u>	Result	<u>Unit</u>	Reference Interval		
LIPID PROFILE					
Cholesterol (Enzymatic colorimetric)	254.6	mg/dL	Desirable : < 200.0 Borderline High : 200-239 High : > 240.0		
Triglyceride (Enzymatic colorimetric)	158.2	mg/dL	Normal : < 150.0 Borderline : 150-199 High : 200-499 Very High : > 500.0		
VLDL	31.64	mg/dL	15 - 35		
Calculated					
LDL CHOLESTEROL	168.06	mg/dL	Optimal : < 100.0 Near / above optimal : 100-129 Borderline High : 130-159 High : 160-189 Very High : >190.0		
HDL Cholesterol	54.9	mg/dL	30 - 70		
Homogeneous enzymatic colorime	etric				
Cholesterol /HDL Ratio Calculated	4.64		0 - 5.0		
LDL / HDL RATIO Calculated	3.06		0 - 3.5		



 Reg. No
 : 2107102624

 Name
 : Ratan Singh

 Age/Sex
 : 55 Years / Male

Collected On : 24-Jul-2021 10:27 **Approved On** : 24-Jul-2021 11:20

Reg. Date

Ref. By

Parameter

Printed On : 24-Jul-2021 11:46

: 24-Jul-2021

Client : MEDIWHEEL WELLNESS

Result Unit Reference Interval

NEW ATP III GUIDELINES (MAY 2001), MODIFICATION OF NCEP<?xml:namespace prefix = "o" ns = "urn:schemas-microsoft-com:office:office" />

LDL CHOLESTEROL
CHOLESTEROL
HDL CHOLESTEROL
TRIGLYCERIDES
Optimal<100
Desirable<200
Low<40
Normal<150
Near Optimal 100-129
Border Line 200-239
High >60
Border High 150-199

Borderline 130-159 High >240 -High 200-499

High 160-189

-

- LDL Cholesterol level is primary goal for treatment and varies with risk category and assesment
- For LDL Cholesterol level Please consider direct LDL value

Risk assessment from HDL and Triglyceride has been revised. Also LDL goals have changed.

- Detail test interpreation available from the lab
- All tests are done according to NCEP guidelines and with FDA approved kits.
- · LDL Cholesterol level is primary goal for treatment and varies with risk category and assesment

For test performed on specimen's received or collected from non-KSHIPRA locations, it is presumed that the specimen belongs to the patient named or identified as labeled on the container/test request and such verification has been carried out at the point generation of the said specimen by the sender.

KSHIPRA will be responsible Only for the analytical part of test carried out. All other responsibility will be of referring Laboratory.

. All other responsibility will be of referring Laboratory.

----- End Of Report ------

Page 5 of 12

Approved by: DR PS RAO

TEST REPORT

: 2107102624 Reg. No Name : Ratan Singh Age/Sex

55 Years / Male

Ref. By

Client : MEDIWHEEL WELLNESS Reg. Date : 24-Jul-2021

Collected On : 24-Jul-2021 10:27 **Approved On** : 24-Jul-2021 11:20

Printed On : 24-Jul-2021 11:46

Parameter Result Unit Reference Interval

HEMOGLOBIN A1 C ESTIMATION

Specimen: Blood EDTA

Hb A1C

6.0

% of Total Hb

Poor Control: > 7.0 % Good Control: 6.2-7.0 % Non-diabetic Level: 4.3-6.2 %

Boronate Affinity with Fluorescent Quenching

136.30

mg/dL

Mean Blood Glucose Calculated

Degree of Glucose Control Normal Range:

Poor Control >7.0% *

Good Control 6.0 - 7.0 %**Non-diabetic level < 6.0 %

- * High risk of developing long term complication such as retinopathy, nephropathy, neuropathy, cardiopathy, etc.
- * Some danger of hypoglycemic reaction in Type I diabetics.
- * Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1c levels in this area.

EXPLANATION:-

Total haemoglobin A1 c is continuously symthesised in the red blood cell throught its 120 days life span. The concentration of HBA1c in the cell reflects the average blood glucose concentration it encounters.

*The level of HBA1c increases proportionately in patients with uncontrolled diabetes. It reflects the average blood glucose oncentration over an extended time period and remains unaffected by short-term fluctuations in blood glucose levels.

*The measurement of HbA1c can serve as a convenient test for evaluating the adequacy of diabetic control and in preventing various diabetic complications. Because the average half life of a red blood cell is sixty days. HbA1c has been accepted as a measurnment which eflects the mean daily blood glucose concentration, better than fasting blood glucose determination, and the degree of carbohydrate imbalance over the preceding two months.

*It may also provide a better index of control of the diabetic patient without resorting to glucose loading procedures.

HbA1c assay Interferences:

*Errneous values might be obtained from samples with abnormally elevated quantities of other Haemoglobins as a result of either their simultaneous elution with HbA1c(HbF) or differences in their glycation from that of HbA(HbS)

----- End Of Report -----

Page 6 of 12

DR PS RAO Approved by:

: 2107102624 Reg. No Name : Ratan Singh Age/Sex : 55 Years / Male Reg. Date : 24-Jul-2021 Collected On : 24-Jul-2021 10:27

Approved On : 24-Jul-2021 11:20

Ref. By

Printed On : 24-Jul-2021 11:46

Client : MEDIWHEEL WELLNESS

<u>Parameter</u>	Result	<u>Unit</u>	Reference Interval		
LIVER FUNCTION TEST					
Total Bilirubin	1.69	mg/dL	0.10 - 1.0		
Colorimetric diazo method					
Conjugated Bilirubin	0.73	mg/dL	0.0 - 0.3		
Sulph acid dpl/caff-benz					
Unconjugated Bilirubin	0.96	mg/dL	0.0 - 1.1		
Sulph acid dpl/caff-benz					
SGOT	27.7	U/L	0 - 37		
(Enzymatic)					
SGPT	36.4	U/L	0 - 40		
(Enzymatic)					
Alakaline Phosphatase	104.8	U/L	53 - 130		
(Colorimetric standardized method)					
Protien with ratio					
Total Protein	7.8	g/dL	6.5 - 8.7		
(Colorimetric standardized method)					
Albumin	4.8	mg/dL	3.5 - 5.3		
(Colorimetric standardized method)					
Globulin	3.00	g/dL	2.3 - 3.5		
Calculated					
A/G Ratio	1.60		0.8 - 2.0		
Calculated					

DR PS RAO MD Pathologist

TEST REPORT

: 2107102624 Reg. No Name : Ratan Singh Age/Sex

: 55 Years / Male

Ref. By

<u>Parameter</u>

Client : MEDIWHEEL WELLNESS Reg. Date

: 24-Jul-2021

Collected On

: 24-Jul-2021 10:27

Approved On : 24-Jul-2021 11:20

Printed On : 24-Jul-2021 11:46

Result <u>Unit</u> **Reference Interval**

SERUM PROTEIN WITH A/G RATIO

Protien with ratio						
Total Protein (Colorimetric standardized method)	7.8	g/dL	6.5 - 8.7			
Albumin (Colorimetric standardized method)	4.8	mg/dL	3.5 - 5.3			
Globulin Calculated	3.00	g/dL	2.3 - 3.5			
A/G Ratio Calculated	1.60		0.8 - 2.0			
BIO - CHEMISTRY						
Creatinine	1.35	mg/dL	0.5 - 1.4			
(Jaffe method)						
(Jaffe method) Alakaline Phosphatase (Colorimetric standardized method)	104.8	U/L	53 - 130			
Alakaline Phosphatase	104.8 15.2					
Alakaline Phosphatase (Colorimetric standardized method)		U/L	53 - 130			

----- End Of Report -----

Page 8 of 12

This is an electronically authenticated report.

Approved by: MD Pathologist

 Reg. No
 : 2107102624

 Name
 : Ratan Singh

 Age/Sex
 : 55 Years / Male

Collected On : 24-Jul-2021 10:27 **Approved On** : 24-Jul-2021 11:46

Reg. Date

Ref. By

Printed On : 24-Jul-2021 11:46

: 24-Jul-2021

Client: MEDIWHEEL WELLNESS

<u>Parameter</u>	Result	<u>Unit</u>	Reference Interval		
THYROID FUNCTION TEST					
T3 (Triiodothyronine)	1.40	ng/mL	0.87 - 1.81		
Chemiluminescence					
T4 (Thyroxine)	8.29	μg/dL	5.89 - 14.9		
Chemiluminescence					
TSH (ultra sensitive)	3.742	μIU/ml	0.34 - 5.6		
Chemiluminescence					

SUMMARY The hypophyseal release of TSH (thyrotropic hormone) is the central regulating mechanism for the biological action of thyroid hormones. TSH is a very sensitive and specific parameter for assessing thyroid function and is particularly suitable for early detection or exclusion of disorders in the central regulating circuit between the hypothalamus, pituitary and thyroid. LIMITATION Presence of autoantibodies may cause unexpected high value of TSH

----- End Of Report -----

TEST REPORT

Reg. No : 2107102624 Name : Ratan Singh

: 24-Jul-2021 Collected On : 24-Jul-2021 10:27

Age/Sex

: 55 Years / Male

Approved On : 24-Jul-2021 11:46

Ref. By

Parameter

: MEDIWHEEL WELLNESS

Printed On

Reg. Date

: 24-Jul-2021 11:46

Client

Result

<u>Unit</u>

Reference Interval

PROSTATE SPECIFIC ANTIGEN

PSA 1.78 ng/mL

0 - 4

Chemiluminescence

----- End Of Report -----

This is an electronically authenticated report.

Page 10 of 12

TEST REPORT

Reg. No : 2107102624 : Ratan Singh Name Age/Sex : 55 Years / Male

Collected On : 24-Jul-2021 10:27

Reg. Date

Ref. By

Client

Approved On : 24-Jul-2021 11:46 **Printed On** : 24-Jul-2021 11:46

: 24-Jul-2021

Parameter

: MEDIWHEEL WELLNESS

Result <u>Unit</u> Reference Interval

URINE ROUTINE EXAMINATION

PHYSICAL EXAMINATION

Quantity 20 cc

Pale Yellow Colour

Clear **Appearance**

CHEMICAL EXAMINATION (BY REFLECTANCE PHOTOMETRIC METHOD)

5.0 - 8.0рΗ 6.0 1.015 1.002 - 1.03 Sp. Gravity

Nil Protein Glucose Nil **Ketone Bodies** Nil Urine Bile salt and Bile Pigment Nil Urine Bilirubin Nil Nitrite Nil Leucocytes Nil Blood Nil

MICROSCOPIC EXAMINATION (MANUAL BY MCIROSCOPY)

Nil

Leucocytes (Pus Cells) Nil Erythrocytes (Red Cells) Nil **Epithelial Cells** 1-2/hpf **Amorphous Material** Nil Casts Nil Nil Crystals **Bacteria** Nil

----- End Of Report -----

Page 11 of 12

Monilia

DR PS RAO Approved by:

TEST REPORT

2107102624 Reg. No Name Ratan Singh Age/Sex

Reg. Date : 24-Jul-2021 **Collected On** : 24-Jul-2021 10:27

: 55 Years / Male

Approved On : 24-Jul-2021 11:46

Ref. By

<u>Parameter</u>

MEDIWHEEL WELLNESS

Printed On : 24-Jul-2021 11:46

Client

<u>Unit</u> Reference Interval

STOOL EXAMINATION

Consistency Semi Solid

Result

CHEMICAL EXAMINATION

Occult Blood Negative

Peroxidase Reaction with o-

Dianisidine

Acidic Reaction

pH Strip Method

Reducing Substance Absent

Benedict's Method

MICROSCOPIC EXAMINATION

Mucus Nil

1 - 2/hpf Pus Cells

Red Cells Nil **Epithelial Cells** Nil Vegetable Cells Nil **Trophozoites** Nil Cysts Nil Ova Nil **Neutral Fat** Nil

Note: Stool occult blood test is highly sensitive to peroxidase like activity of free hemoglobin.

Nil

False negative: False negative occult blood test may be observed in case of excess (>250mg/day) Vitamin C intake and in case of occassinal unruptured RBCs.

False positive: False positive occult blood test may be observed in stool samples containing vegetable peroxidase (turnips, horseradish, cauliflower, brocoli, cantaloupe, parsnips) and myoglobin from food (meat diet) intake.

----- End Of Report -----

Page 12 of 12

Monilia