



॥ ॐ गणेशाय नमः ॥

GANESH DIAGNOSTIC

DR. LOKESH GOYAL

MBBS (KGMC), MD (RADIOLOGY)

CONSULTANT INTERVENTIONAL RADIOLOGIST
FORMER SR. REGISTRAR - APOLLO HOSPITAL, NEW DELHI
LIFE MEMBER OF IRIA

Timings : 9:00 am to 9:00 pm, Sunday 9.00 am to 3.00 pm

☎ 8392957683, 6395228718

MRS. SONI GANGWAR
DR. NITIN AGARWAL, DM

11-03-2023

REPORT

EXAMINATION PERFORMED: X-RAY CHEST

B/L lung fields are clear

Both of the CP angles are clear.

Both hila show a normal pattern .

Cardiac and mediastinal borders appear normal.

Visualized bony thorax and soft tissue of the chest wall appear normal.

IMPRESSION ---NO SIGNIFICANT ABNORMALITY IS SEEN

Not for medico-legal purpose

DR LOKESH GOYAL
MD
RADIOLOGIST

डि
सी

डिजिटल एक्स-रे, मल्टी स्लाइस
सी. टी. स्कैन सुविधा उपलब्ध है।



NOT VALID FOR
MEDICO LEGAL PURPOSE

Dr. Nitin Agarwal

MD, DM (Cardiology)

Consultant Interventional Cardiologist

Cell : +91-94578 33777

Formerly at :

Escorts Heart Institute & Research Centre, Delhi

Dr. Ram Manohar Lohia Hospital, Delhi



APPLE
CARDIAC CARE

DR. NITIN AGARWAL'S HEART CLINIC

Some features

11/2/23

*

T - cycle (200)

0 - - -

T - cycle (100)

0 - - -

limi Area for

in Area

Eye and

0

0

in Area

A-3, EKTA NAGAR, (OPP. CARE HOSPITAL) STADIUM ROAD, NEAR DELAPEER CHAURAHA, BARREILLY - 243 122 (U.P.)

OPD Timings : 12.00 Noon to 04.00 pm, Sunday : 12.00 Noon to 3.00 pm

स्वास्थ्य के लिए फोन करें : 09458888448, 07599031977

VALID FOR 5 DAYS.

पचास दिनों के लिए मान्य

Timings : 12 noon

A Venture of Apple Cardiac Care

A-3, Ekta Nagar, Stadium Road,
(Opp. Care Hospital),
Bareilly - 243 122 (U.P.) India
Tel. : 07599031977, 09458888448



Reg.NO. : 116
NAME : **Mrs. SONY GANGWAR**
REFERRED BY : Dr.Nitin Agarwal (D M)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 27 Yrs.
SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
HAEMATOLOGY			
COMPLETE BLOOD COUNT (CBC)			
HAEMOGLOBIN	13.1	gm/dl	12.0-15.0
TOTAL LEUCOCYTE COUNT	6,700	/cumm	4,000-11,000
DIFFERENTIAL LEUCOCYTE COUNT(DLC)			
Neutrophils	67	%	40-75
Lymphocytes	30	%	20-45
Eosinophils	03	%	01-08
Monocytes	00	%	01-06
Basophils	00	%	00-02
TOTAL R.B.C. COUNT	4.43	million/cumm	3.5-6.5
P.C.V./ Haematocrit value	41.7	%	35-54
M C V	94.1	fL	76-96
M C H	29.6	pg	27.00-32.00
M C H C	31.4	g/dl	30.50-34.50
PLATELET COUNT	2.20	lacs/mm ³	1.50 - 4.50
E.S.R (WINTROBE METHOD)			
-in First hour	14	mm	00- 20
BLOOD GROUP	B+		
Blood Group	POSITIVE		
BIOCHEMISTRY			
BLOOD SUGAR F.	86	mg/dl	60-100

HAEMATOLOGY

Report is not valid for medicolegal purpose

Adventure of Apple Cardiac Care

73, Ekta Nagar, Stadium Road,
(Opp. Care Hospital),
Bareilly - 243 122 (U.P.) India
Tel. : 07599031977, 09458888448



APPLE
PATHOLOGY
TRUSTED RESULT

Reg.NO. : 116
NAME : Mrs. SONY GANGWAR
REFERRED BY : Dr. Nitin Agarwal (D.M.)
SAMPLE : BLOOD
DATE : 11/03/2023
AGE : 27 Yrs.
SEX : FEMALE

TEST NAME	RESULTS	UNITS	BIOLOGICAL REF. RANGE
GLYCOSYLATED HAEMOGLOBIN	5.7		

EXPECTED RESULTS :

 Non diabetic patients : 4.0% to 6.0%
 Good Control : 6.0% to 7.0%
 Fair Control : 7.0% to -8%
 Poor Control : Above 8%

*ADA: American Diabetes Association

The glycosylated hemoglobin assay has been validated as a reliable indicator of mean blood glucose levels for a period of 8-12 week period prior to HBA1C determination. ADA recommends the testing twice a year in patients with stable blood glucose, and quarterly, if treatment changes, or if blood glucose levels are unstable.

METHOD : ADVANCED IMMUNO ASSAY.

BIOCHEMISTRY

Gamma Glutamyl Transferase (GGT)	37	U/L	11-50
----------------------------------	----	-----	-------

BLOOD UREA	24	mg/dL.	10-40
------------	----	--------	-------

* Low serum urea is usually associated with status of overhydration severe hepatic failure.

* A urea level of 10-45 mg/dl indicates normal glomerular function and a level of 100-250 mg/dl indicates a serious impairement of renal function. In chronic renal failure , urea correlates better with the symptoms of uremia than does serum creatinine.

* Urine/Serum urea is more than 9 in prerenal and less than 3 in renal uremia.

SERUM CREATININE	0.6	mg/dL.	0.5-1.4
------------------	-----	--------	---------

Report is not valid for medicolegal purpose

Reg.NO. : 116 DATE : **11/03/2023**
 NAME : **Mrs. SONY GANGWAR** AGE : 27 Yrs.
 REFERRED BY : Dr.Nitin Agarwai (D M) SEX : FEMALE
 SAMPLE : BLOOD

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
URIC ACID	5.7	mg/dl	3.0-6.0

CLINICAL SIGNIFICANCE:

Analysis of synovial fluid plays a major role in the diagnosis of joint disease.

SERUM SODIUM (Na)	137	m Eq/litre.	135 - 155
SERUM POTASSIUM (K)	3.9	m Eq/litre.	3.5 - 5.5
SERUM CALCIUM	9.0	mg/dl	8.5 - 10.5

LIVER PROFILE

SERUM BILIRUBIN			
TOTAL	0.5	mg/dL	0.3-1.2
DIRECT	0.3	mg/dL	0.2-0.6
INDIRECT	0.2	mg/dL	0.1-0.4
SERUM PROTEINS			
Total Proteins	6.6	Gm/dL	6.4 8.3
Albumin	4.1	Gm/dL	3.5 - 5.5
Globulin	2.5	Gm/dL	2.3 - 3.5
A : G Ratio	1.64		0.0-2.0
SGOT	56	IU/L	0-40
SGPT	52	IU/L	0-40
SERUM ALK.PHOSPHATASE	71	IU/L	00-115

NORMAL RANGE : BILIRUBIN TOTAL

Premature infants. 0 to 1 day: <8 mg/dL Premature infants. 1 to 2 days: <12 mg/dL Adults: 0.3-1 mg/dL.
 Premature infants. 3 to 5 days: <16 mg/dL Neonates, 0 to 1 day: 1.4-8.7 mg/dL
 Neonates, 1 to 2 days: 3.4-11.5 mg/dL Neonates, 3 to 5 days: 1.5-12 mg/dL Children 6 days to 18 years: 0.3-1.2 mg/dL

COMMENTS-

Total and direct bilirubin determination in serum is used for the diagnosis, differentiation and follow -up of jaundice. Elevation of SGPT is found in liver and kidney diseases such as infectious or toxic hepatitis, IM and cirrhosis. Organs rich in SGOT are heart, liver and skeletal muscles. When any of these organs are damaged, the serum SGOT level rises in proportion to the severity of damage. Elevation of Alkaline Phosphatase in serum or plasma is found in hepatitis, biliary obstructions, hyperparathyroidism, steatorrhea and bone diseases.

Report is not valid for medicolegal purpose

Reg.NO. : 116
NAME : Mrs. SONY GANGWAR
REFERRED BY : Dr. Nitin Agarwal (D.M.)
SAMPLE : BLOOD

DATE : 11/03/2023
AGE : 27 Yrs.
SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
LIPID PROFILE			
SERUM CHOLESTEROL	177	mg/dL.	130 - 200
SERUM TRIGLYCERIDE	97	mg/dl.	30 - 160
HDL CHOLESTEROL	48	mg/dL.	30-70
VLDL CHOLESTEROL	19.4	mg/dL.	15 - 40
LDL CHOLESTEROL	109.60	mg/dL.	00-130
CHOL/HDL CHOLESTEROL RATIO	3.69	mg/dl	
LDL/HDL CHOLESTEROL RATIO	2.28	mg/dl	

INTERPRETATION

TRIGLYCERIDE level > 250mg/dL is associated with an approximately 2-fold greater risk of coronary vascular disease. Elevation of triglycerides can be seen with obesity, medication, fast less than 12 hrs., alcohol intake, diabetes melitus, and pancreatitis.

CHOLESTEROL, its fractions and triglycerides are the important plasma lipids in defining cardiovascular risk factors and in the management of cardiovascular disease. Highest acceptable and optimum values of cholesterol values of cholesterol vary with age. Values above 220 mgm/dl are associated with increased risk of CHD regardless of HDL & LDL values.

HDL-CHOLESTEROL level <35 mg/dL is associated with an increased risk of coronary vascular disease even in the face of desirable levels of cholesterol and LDL - cholesterol.

LDL - CHOLESTEROL & TOTAL CHOLESTEROL levels can be strikingly altered by thyroid, renal and liver disease as well as hereditary factors. Based on total cholesterol, LDL- cholesterol, and total cholesterol/HDL - cholesterol ratio, patients may be divided into the three risk categories.

URINE EXAMINATION

Report is not valid for medicolegal purpose

Reg.NO. : 116
 NAME : **Mrs. SONY GANGWAR**
 REFERRED BY : Dr.Nitin Agarwal (D.M)
 SAMPLE : BLOOD
 DATE : **11/03/2023**
 AGE : 27 Yrs.
 SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
URINE EXAMINATION REPORT			
PHYSICAL EXAMINATION			
pH	6.0		
TRANSPARENCY			
Volume	25	ml	
Colour	Light Yellow		
Appearance	Clear		Nil
Sediments	Nil		
Specific Gravity	1.020		1.015-1.025
Reaction	Acidic		
BIOCHEMICAL EXAMINATION			
UROBILINOGEN	Nil		NIL
BILIRUBIN	Nil		NEGATIVE
URINE KETONE	Nil		NEGATIVE
Sugar	Nil		Nil
Albumin	Nil		Nil
Phosphates	Absent		Nil
MICROSCOPIC EXAMINATION			
Red Blood Cells	Nil	/H.P.F.	
Pus Cells	100-120	/H.P.F.	
Epithelial Cells	10-15	/H.P.F.	
Crystals	NIL		NIL
Casts	NIL	/H.P.F.	
Bacteria	NIL		
Other	NIL		

BIOCHEMISTRY

Report is not valid for medicolegal purpose

enture of Apple Cardiac Care
F-3, Ekta Nagar, Stadium Road,
(Opp. Care Hospital),
Bareilly - 243 122 (U.P.) India
Tel. : 07599031977, 09458888448



APPLE
PATHOLOGY
TRUSTED RESULT

Reg.NO. : 116
NAME : **Mrs. SONY GANGWAR**
REFERRED BY : Dr.Nitin Agarwal (D.M)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 27 Yrs.
SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
BLOOD SUGAR P.P.	119	mg/dl	80-140

--{End of Report}--

Dr. Shweta Agarwal
MD(Pathology), Apple Pathology
Bareilly (UP)



NAME	Mrs. SONY GANGWAR	AGE/SEX	27 Y/F
Ref. By	Dr. NITIN AGARWAL (DM)	DATE	11/03/2023

ECHOCARDIOGRAPHY AND COLOUR DOPPLER STUDY

<u>MEASUREMENTS</u>	<u>VALUE</u>	<u>NORMAL DIMENSIONS</u>
LVID (d)	4.5 cm	(3.7 –5.6 cm)
LVID (s)	2.4 cm	(2.2 –3.9 cm)
RVID (d)	2.4 cm	(0.7 –2.5 cm)
IVS (ed)	1.0 cm	(0.6 –1.1 cm)
LVPW (ed)	1.0 cm	(0.6 –1.1 cm)
AO	2.2 cm	(2.2 –3.7 cm)
LA	3.0 cm	(1.9 –4.0 cm)
<u>LV FUNCTION</u>		
EF	60 %	(54 –76 %)
FS	30 %	(25 –44 %)

- LEFT VENTRICLE** : No regional wall motion abnormality
No concentric left Ventricle Hypertrophy
- MITRAL VALVE** : Thin, PML moves posteriorly during Diastole
No SAM, No Subvalvular pathology seen.
No mitral valve prolapse calcification .
- TRICUSPID VALVE** : Thin, opening wells. No calcification, No doming .
No Prolapse.
Tricuspid inflow velocity= 0.7 m/sec
- AORTIC VALVE** : Thin, tricuspid, opening well, central closer,
no flutter.
No calcification
Aortic velocity = 1.3 m/sec
- PULMONARY VALVE** : Thin, opening well, Pulmonary artery is normal
EF slope is normal.
Pulmonary Velocity = 0.9 m /sec



ON DOPPLER INTERROGATION THERE WAS :

- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

MITRAL FLOW

E= 0.8 m/sec

A= 0.6 m/sec

ON COLOUR FLOW:


- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

COMMENTS:

- No LA /LV clot
- No pericardial effusion
- No intracardiac mass
- IAS/IVS Intact
- Inferior vena cava – normal in size with normal respiratory variation

FINAL IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY
- NORMAL LV DIASTOLIC FUNCTION
- NORMAL LV SYSTOLIC FUNCTION (LVEF~60%)
- NORMAL CARDIAC CHAMBER DIMENSIONS
- NORMAL VALVULAR COLOUR FLOW PATTERN


DR.NITIN AGARWAL
DM (Cardiology)
Consultant Cardiologist

This opinion is to be correlated with the clinically findings and if required, please re-evaluate / reconfirm with further investigation.



PARAS MRI & ULTRASOUND CENTRE

MOST ADVANCED 32 CHANNEL 3T 3D WHOLE BODY MRI

261, ASHAPURAM, OPP. DR. BASU EYE HOSPITAL, STADIUM ROAD, BAREILLY

• Helpline : 7300761761 • E-mail : parasmribly@gmail.com

REPORT

4D / 5D ULTRASOUND

COLOR DOPPLER

TVS/ TRUS

MUSCULOSKELETAL USG

Date : 11.03.2023
Name : SONY GANGWAR 27Y/F
Ref.By : APPLE CARDIAC CARE

ULTRASOUND WHOLE ABDOMEN

LIVER - Liver is normal in size and outline. It shows increased echogenicity. No obvious focal pathology is seen. The intra hepatic biliary radicals are not dilated. PV –normal.

GALL BLADDER -Gall Bladder is normal in size, has normal wall thickness with no evidence of calculi. Fat planes between GB and liver are well maintained.The CBD appears normal.

PANCREAS - Pancreas is normal in size and echogenicity. Its outlines are distinct. No obvious focal lesion, calcification or ductile dilatation is seen.

SPLEEN - Spleen is normal in size and echogenicity. There is no evidence of collaterals

KIDNEYS - Both kidneys are normal in position, outline and echogenicity. No evidence of calculi is seen. CMD is maintained. No evidence of hydronephrosis is seen on either side.

URINARY BLADDER -Urinary Bladder is normal in size and outline. There is no evidence of any obvious intraluminal or paramedical pathology. Wall is not thickened.

Uterus - Uterus is retroflexed and normal in size. The myometrial and endometrial echoes are normal. The endometrial thickness measures 6mm. No focal lesion is seen.

Both ovaries are normal in size and shows uniform parenchymal echogenicity and smooth outlines. No evidence of cyst or mass is seen.

No evidence of ascites /pleural effusion/adenopathy is seen. Bowel loops are not dilated. Bilateral iliac fossa appears normal.

IMPRESSION:



❖ Grade I fatty liver.


Adv clinical correlation.


Dr. Puja Tripathi

M.B.B.S., M.D.

MBBS, MD (Radiodiagnosis, SGPGI)





PAT. 10. SON, GORDON 11/8/23

~~REGULAR SINUS BRADYCARDIA~~
~~REGULAR SINUS BRADYCARDIA~~
~~REGULAR SINUS BRADYCARDIA~~


DPL CARDIAC RIBRI
10mm/10V 25mm/50Hz 200µV

DPL CARDIAC RIBRI
10mm/10V 25mm/50Hz 200µV

DPL CARDIAC RIBRI
10mm/10V 25mm/50Hz 200µV

DPL CARDIAC RIBRI
10mm/10V 25mm/50Hz 200µV





R



Visit ID : MBAR39351	Registration : 11/Mar/2023 05:21PM
UHID/MR No : ABAR.0000039339	Collected : 11/Mar/2023 05:24PM
Patient Name : Ms.SONY	Received : 11/Mar/2023 05:27PM
Age/Gender : 27 Y 0 M 0 D /F	Reported : 11/Mar/2023 07:11PM
Ref Doctor : Dr.NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240,Sanjay Nagar Bareilly (UP)	Barcode No : A3619624

DEPARTMENT OF HORMONE ASSAYS

Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

THYROID PROFILE (T3,T4,ULTRASENSITIVE TSH)

Sample Type : SERUM

T3	1.15	ng/ml	0.61-1.81	CLIA
T4	8.4	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	1.671	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

- Serum T3, T4 and TSH are the measurements form three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
- Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
- Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
- Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propanolol and propylthiouracil.
- Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
- Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
- Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
- TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

REFERENCE RANGE:

PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 – 2.500
2nd Trimester	0.200 – 3.000
3rd Trimester	0.300 – 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

- During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
- TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***



 Dr. Miti Gupta
 DNB ; MD [Pathology]
