

CERTIFICATE OF MEDICAL FITNESS

NAME: <u>mita madheismeta</u>
AGE/GENDER: 38 you / remale
HEIGHT: 16000 WEIGHT: 58.7kg
IDENTIFICATION MARK: Black mole In left Neck
BLOOD PRESSURE: 110 / formon leg
PULSE: 76/min
CVG.
RS:P & NO Gronal
ANY OTHER DISEASE DIAGNOSED IN THE PAST: 1/1/
ALLERGIES, IF ANY: Nil
LIST OF PRESCRIBED MEDICINES: Nil/
ANY OTHER REMARKS: NO
of Ms Ponder to Krimined Mr/Mrs. <u>Meta madhalineta</u> son/daughter of Ms Ponder to Krimined Mr/Mrs. In the presence. He/ she has no physical
disease and is fit for employment. Dr. BINDURAJ. R
MBBS, MD
Signature of candidate Signature of Medical Officer
Place: Spackgrown Diagnostics Phealth case,
Date: 16/10/23
Disclaiman, The matient to the state of the

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined







Dr. Ashok S Bsc., MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827

DATE: 16-10-2-3

EYE EXAMINATION

NAME: Mss. onfa ma	a dhu AGE: 355)	GENDER : F / M
	RIGHT EYE	LEFT EYE
Vision	619:06	6/3:00
Vision With glass	Ellin-	- Elino
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nill	Nill
Diagnosis/ impression	Normal	Normal
	Ey Cons	ORSARODARS C., M.B.B.S., D.O.M.S. Sultant & Surgeon MC 31827
		pthalmologist)







NAME	AGE	GENDER
Mr. Mordinguita	38711	Sende.

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY

O: OTHERS

grone.
Singated 3/8; Extrem recommended.

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS: ç.

Tr.

SIGNATURE OF THE DENTAL SURGEON

SEAL

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA)

DATE

Reg. No: 2247/A





0.15~35Hz AC50 25mm/s] aVL	avr			ID: 1610230058 16-10-2023 MRS MITA MADHUSMITA MISHRA HR Female 38Years PR ORS ORS OT/QTc P/QRS/I RV5/SV1
mm/s 10mm/mV 2*5.0s \(\pi\)64 \(\nabla\)2.2						10:43:10 64 bpm 99 ms 150 ms 87 ms 87 ms 383/396 ms 58/52/49 ° 0.783/0.612 mV
SEMIP V181 SPECTRUM DIAGNOSTICS & HEALTH CARE	V6	Vs	V4	V3		Diagnosis Information: Sinus Rhythm ****Normal ECG**** Report Confirmed by:
CS & HEALTH CARE						

SPECTRUM DIAGNOSTICS & HEALTH CARE

#9/1 TEJAS ARCADE, DR. RAJKUMAR ROAD, RAJAJINAGAR-560010 AUDIOGRAN




Patient ID: 0915

Name: MITA MADHUSMITHA MISHRA

CR Number: 20231016113138 Registration Date: 16-Oct-2023 Age: 38

Gender: Female

Operator: spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 Hz	1500 Hz	2000 Hz	3000 Hz	4000 Hz	6000 Hz	8000 Hz
X - Air Left	25	35	20	25	25	20	20	20	15	15	10
O - Air Right	30	35	25	25	30	25	20	25	20	20	15
> - Bone Left											
< - Bone Right											

Clinical Notes:

Not Found	
	ETICC
	SHOTH SHO
	13/4 7
	I DENGALIE
	18
nttps://www.rmsindia.com © RMS Audiometer(HERMES v3.0.0.7) Print Date:16-Oct-2023	4



NAME : MRS.MITA MADHUSMITA MISHRA	DATE :16/10/2023 REG NO:1610230058
AGE/SEX : 38 YEARS/FEMALE	REG NO:1010230030
REF BY : APOLO CLINIC	

CHEST PA VIEW

Lung fields are clear.

Cardiovascular shadows are within normal limits.

Both CP angles are free.

Domes of diaphragm and bony thoracic cage are normal.

IMPRESSION: NORMAL CHEST RADIOGRAPH.

DR.RAM PRAKASH G MDRD CONSULTANT RADIOLOGIST

RH1-14

Your suggestion / feedback is a valuable input for improving our services





PATIENT NAME	MRS. MITA MADHUSMITA MISHRA	ID NO	1610230058
AGE	38YEARS	SEX	FEMALE
REF BY	DR. APOLO CLINIC	DATE	16.10.2023

2D ECHO CARDIOGRAHIC STUDY

M-MODE

141.	TVIODE	
AORTA	22mm	
LEFT ATRIUM	27mm	
RIGHT VENTRICLE	20mm	
LEFT VENTRICLE (DIASTOLE)	38mm	
LEFT VENTRICLE(SYSTOLE)	26mm	
VENTRICULAR SEPTUM (DIASTOLE)	07mm	
VENTRICULAR SEPTUM (SYSTOLE)	07mm	
POSTERIOR WALL (DIASTOLE)	09mm	-
POSTERIOR WALL (SYSTOLE)	09mm	
FRACTIONAL SHORTENING	30%	-
EJECTION FRACTION	60%	

DOPPLER /COLOUR FLOW

MITRAL VALVE	E-1.06m/sec	A-0.62m/sec	TRIVIAL MR
AORTIC VALVE	1.18 m/sec		NO AR
PULMONARY VALVE	0.89 m/sec		NO PR
TRISCUSPID VALVE			MILD TR





PATIENT NAME	MRS. MITA MADHUSMITA	ID NO	1610230058
	MISHRA		
AGE	38YEARS	SEX	FEMALE
REF BY	DR. APOLO CLINIC	DATE	16.10.2023

2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE : NORMAL	
LEFT ATRIUM : NORMAL	
RIGHT ATRIUM: NORMAL	
MITRAL VALVE : NORMAL	
AORTIC VALVE : NORMAL	
PULMONARY VALVE: NORMAL	
TRICUSPID VALVE: NORMAL	
INTER ATRIAL SEPTUM :INTACT	
INTER VENTRICULAR SEPTUM: INTACT	
PERICARDIUM : NORMAL	
OTHERS : - NIL	

IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- > NORMAL VALVES AND DIMENSIONS
- ➢ NORMAL LV FUNCTION, LVEF- 60%
- > TRIVIAL MR / MILD TR / NO PAH
- NO CLOT / VEGETATION / EFFUSION
- NO ASD / VSD / PDA / Coa SEEN

The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.







NAME AND LAB NO	MRS.MITA MADHUSMITA MISHRA	REG-0058
AGE & SEX	38 YRS	FEMALE
DATE AND AREA OF INTEREST	16.10.2023	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	THE STREET OF LEVIS

USG ABDOMEN AND PELVIS

LIVER:

Measures 12.7cm. Normal in size and echotexture.

No e/o IHBR dilatation. No evidence of SOL.

Portal vein appears normal.

CBD appears normal. . No e/o calculus / SOL

GALL BLADDER:

Collapsed.

SPLEEN:

Measures 9.0 cm. Normal in size and echotexture. No e/o SOL/ calcification.

PANCREAS:

Head and body appears normal in size and echotexture.

Pancreatic duct appears normal. No e/o calculus / calcifications.

RETROPERITONEUM:

Poor window.

RIGHT KIDNEY:

Measures 10.9 x 4.0 cm. Right kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

LEFT KIDNEY:

Measures 10.7 x 5.4 cm .Left kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

URETERS:

Bilateral ureters are not dilated.

URINARY BLADDER:

Well distended. No wall thickening/ calculi.

UTERUS:

Normal in size and echotexture

Endometrium is normal.ET -8.0 mm. Small seedlings fibroid measuring 10 x 7

mm in the posterior wall.

OVARIES:

B/L ovaries normal in size and echotexture.

No evidence of ascites/pleural effusion.

IMPRESSION:

No significant sonological abnormality detected in the abdomen and pelvis.

DR.AKSHATHA R BHAT MDRD DNB FRCR



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org





: 38 years / Female : Dr. APOLO CLINIC

Ref. By Dr. Reg. No. : 1610230058

Age / Gender

C/o : Apollo Clinic **Bill Date** : 16-Oct-2023 09:13 AM Sample Col. Date: 16-Oct-2023 09:13 AM

Result Date : 16-Oct-2023 04:35 PM **Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole I	Blood EDTA			
Haemoglobin (HB)	12.20	g/dL	Male: 14.0-17.0 Female:12.0-15.0 Newborn:16.50 - 19.50	Spectrophotmeter
Red Blood Cell (RBC)	4.42	million/cur	nm3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	38.30	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	86.70	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)		pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	31.80	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	45.30	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	15.70	%	Male: 11.80-14.50 Female: 12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	8.80	fL	8.0-15.0	Volumetric Impedance
Platelet	3.36	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	11.20	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)		cells/cumm	Male: 4000.0-11000.0 Female 4000.0-11000.0 Children: 6000.0-17500.0 Infants: 9000.0-30000.0	Volumetric Impedance
Neutrophils	62.70	%	40.0-75.0	Light
Lymphocytes	32.80	%	20.0-40.0	scattering/Manual Light
Eosinophils	1.40	%	0.0-8.0	scattering/Manual Light scattering/Manual

UHID

: 1610230058

1610230058









Age / Gender : 38 years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 1610230058

C/o : Apollo Clinic

Bill Date : 16-Oct-2023 09:13 AM : 1610230058

Sample Col. Date: 16-Oct-2023 09:13 AM

Result Date : 16-Oct-2023 04:35 PM

Report Status :	Final	
-----------------	-------	--

Test Name	Result	Unit	Reference Value	Method
Monocytes	3.10	%	0.0-10.0	Light
Basophils	0.00	%	0.0-1.0	scattering/Manual Light
Absolute Neutrophil Count	3.53	10^3/uL	2.0- 7.0	scattering/Manual Calculated
Absolute Lymphocyte Count	1.85	10^3/uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.17	10^3/uL	0.20-1.00	Calculated
Absolute Eosinophil Count	80.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.00	10^3/uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	12	mm/hr	Female: 0.0-20.0 Male: 0.0-10.0	Westergren

1610230058

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.

WBC'S : Are normal in total number, morphology and distribution.

: Adequate in number and normal in morphology. **Platelets**

No abnormal cells or hemoparasites are present.

Impression: Normocytic Normochromic Blood picture.



Printed By

+91 77604 97644 | 080 2337 1555

: spectrum

Printed On

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist







Age / Gender : 38 years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1610230058

C/o : Apollo Clinic **Bill Date**

: 16-Oct-2023 09:13 AM

Result Date

Sample Col. Date: 16-Oct-2023 09:13 AM : 16-Oct-2023 04:35 PM

Report Status

: Final

Test Name	Result	Unit	Reference Value	Method
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)
Fasting Blood Sugar (FBS)- Plasma	92	mg/dL	60.0-110.0	Hexo Kinase

1610230058

: 1610230058

UHID

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By

: spectrum

Printed On : 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C, MD, Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalurមិខ្មែរទី៦២០មិ www.spectrumdiagnostics.org





: 38 years / Female

: Dr. APOLO CLINIC

Reg. No. : 1610230058

Age / Gender

Ref. By Dr.

C/o : Apollo Clinic Bill Date : 16-Oct-2023 09:13 AM

Sample Col. Date: 16-Oct-2023 09:13 AM Result Date : 16-Oct-2023 04:35 PM

Report Status : Final

Test Name Result Unit Reference Value Method

1610230058

Postprandial Urine glucose-Urine

Negative

98

Negative

: 1610230058

Dipstick/Benedicts

(Manual)

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

UHID

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

Post prandial Blood Glucose (PPBS)-Plasma

mg/dL

70-140

Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total



Printed By

: spectrum

Printed On

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalur^{Pag}්ත්රේව @ www.spectrumdiagnostics.org







: 38 years / Female

: Dr. APOLO CLINIC

Reg. No. : 1610230058

Age / Gender

Ref. By Dr.

C/o : Apollo Clinic Bill Date

: 16-Oct-2023 09:13 AM

Sample Col. Date: 16-Oct-2023 09:13 AM **Result Date**

: 16-Oct-2023 04:35 PM

Report Status	: Final
---------------	---------

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA				
Glycosylated Haemoglobin (HbA1c)	5.30	%	Non diabetic adults:<5.7 At risk (Prediabetes): 5.7 - 6.4 Diagnosing Diabetes:>= 6.5 Diabetes Excellent Control: 6-7 Fair to good Control: 7-8 Unsatisfactory Control: 8-10	HPLC
Estimated Average Glucose(eAG)	105.41	mg/dL	Poor Control :>10	Calculated

1610230058

: 1610230058

UHID

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By

: spectrum

Printed On

+91 77604 97644 | 080 2337 1555

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalur වීමේන් රාජ් info@spectrumdiagnostics.org @ www.spectrumdiagnostics.org





Age / Gender : 38 years / Female

: Dr. APOLO CLINIC

Reg. No. : 1610230058

C/o : Apollo Clinic

Ref. By Dr.

Bill Date : 16-Oct-2023 09:13 AM

Sample Col. Date: 16-Oct-2023 09:13 AM **Result Date**

Report Status : Final

: 16-Oct-2023 04:35 PM

Test Name Result Unit Reference Value Method Blood Group & Rh Typing-Whole Blood EDTA **Blood Group** Slide/Tube agglutination Rh Type Positive Slide/Tube agglutination

1610230058

: 1610230058

UHID

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By

+91 77604 97644 | 080 2337 1555

: spectrum

Printed On : 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalurមិទ្ធ១៦០៦៤ lnfo@spectrumdiagnostics.org @ www.spectrumdiagnostics.org





Age / Gender : 38 years / Female

: Dr. APOLO CLINIC

Reg. No. : 1610230058

Ref. By Dr.

C/o : Apollo Clinic **Bill Date** : 16-Oct-2023 09:13 AM

Sample Col. Date: 16-Oct-2023 09:13 AM Result Date

Report Status : Final

: 16-Oct-2023 04:35 PM

Test Name	Result	Unit	Reference Value	Method
Calcium, Total- Serum	9.70	mg/dL	8.50-10.10	Spectrophotometry (O- Cresolphthalein
Gamma-Glutamyl Transferase (GGT)-Serum	11.00	U/L	Male: 15.0-85.0 Female: 5.0-55.0	complexone) Other g-Glut-3- carboxy-4 nitro

1610230058

: 1610230058

UHID

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By

: spectrum

Printed On

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



SCAN FOR LOCATION





Age / Gender : 38 years / Female

Ref. By Dr. : Dr. APOLO CLINIC

: 1610230058

C/o : Apollo Clinic

Reg. No.

Bill Date : 16-Oct-2023 09:13 AM

Sample Col. Date: 16-Oct-2023 09:13 AM **Result Date**

: 16-Oct-2023 04:35 PM Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination	on-Urine			
Physical Examination				(*9)
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.00		1.000-1.030	Dipstick
Biochemical Examination	1		11000 11000	Dipstick
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	
Ketone Bodies	Negative		Negative	Dipstick/Fouchets Dipstick/Rotheras
Urobilinogen	Normal		Normal	AND DESCRIPTION OF THE PROPERTY OF THE PROPERT
Nitrite	Negative		Negative	Dipstick/Ehrlichs Dipstick
Microscopic Examination	_			Dipstick
Pus Cells	1-2	hpf	0.0-5.0	Microscopy
Epithelial Cells	3-4	hpf	0.0-10.0	
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent	-	Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy Microscopy

UHID

: 1610230058

1610230058

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By

: spectrum

Printed On

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalur විදේශීර්ණි @+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org





Age / Gender : 38 years / Female

Ref. By Dr. : Dr. APOLO CLINIC **Reg. No.** : 1610230058

C/o : Apollo Clinic

Bill Date : 16-Oct-2023 09:13 AM : 1610230058 Sample Col. Date : 16-Oct-2023 09:13 AM

Sample Col. Date: 16-Oct-2023 09:13 AM **Result Date** : 16-Oct-2023 04:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test) Blood Urea Nitrogen (BUN)- Serum	: 11.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.75	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	5.37	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	140.0	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.23	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	102.60	mmol/L	94.0-110.0	Ion-Selective Electrodes (ISE)

1610230058



Printed By

: spectrum

Printed On : 16 Oct, 2023 04:57 pm

Millim.

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalur (1925) 177604 97644 | 080 2337 1555 info@spectrumdiagnostics.org





Name : MRS. MITA MADHUSMITA MISHRA Bill Date : 16-Oct-2023 09:13 AM

 Age / Gender
 : 38 years / Female
 UHID
 : 1610230058
 Sample Col. Date : 16-Oct-2023 09:13 AM

 Ref. By Dr.
 : Dr. APOLO CLINIC
 Result Date
 : 16-Oct-2023 04:35 PM

Reg. No. : 1610230058 1610230058 Report Status : Final

C/o : Apollo Clinic

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)- Serum				
Tri-Iodo Thyronine (T3)-Seru	ım 1.05	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	9.70	μg/dL	Female: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormon (TSH)-Serum	e 2.92	μIU/mL	Female: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15

Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years: 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days :>6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments:TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.

els: Graves disease, Autonomous thyroid hormone secretion, TSH defic

Printed By

: spectrum

Printed On

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



SCAN FOR LOCATION





Bill Date

: 16-Oct-2023 09:13 AM

Age / Gender

: 38 years / Female

: 1610230058

Sample Col. Date: 16-Oct-2023 09:13 AM

Ref. By Dr.

: Dr. APOLO CLINIC

Result Date

: 16-Oct-2023 04:35 PM

Reg. No.

C/o

: 1610230058 : Apollo Clinic 1610230058

UHID

Report Status

: Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	183.00	mg/dL	Female: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	62.00	mg/dL	Female: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	56.00	mg/dL	Female: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	127	mg/dL	Female: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	99.00	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	12	mg/dL	Female: 0.0 - 40	Calculated
	3.27	Ratio	Female: 0.0 - 5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: spectrum

Printed On

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist







: 38 years / Female

: Dr. APOLO CLINIC

Reg. No. : 1610230058

Age / Gender

Ref. By Dr.

C/o

: Apollo Clinic

Bill Date : 16-Oct-2023 09:13 AM

Sample Col. Date: 16-Oct-2023 09:13 AM

Result Date : 16-Oct-2023 04:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serun	1			
Bilirubin Total-Serum	0.50	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.13	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.37	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	17.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	16.00	U/L	Male:16.0-63.0 Female:14.0-59.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)- Serum	62.00	U/L	Adult: 45.0-117.0 Children: 48.0-445.0 Infants: 81.90-350.30	PNPP,AMP- Buffer
Protein, Total-Serum	7.74	g/dL	6.40-8.20	Biuret/Endpoint- With Blank
Albumin-Serum	4.43	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	3.31	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.34	Ratio	0.80-1.20	Calculated ·

UHID

: 1610230058

1610230058



Printed By

: spectrum

Printed On

: 16 Oct, 2023 04:57 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



SCAN FOR LOCATION

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru 550000 +91 77604 97644 | 080 2337 1555 @info@spectrumdiagnostics.org