Name	: Mr. MALAGRI PUNDLIK			
PID No.	: MED111087640	Register On	: 14/05/2022 9:45 AM	M
SID No.	: 712214641	Collection On	: 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On	: 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On	: 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel			

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	17.1	g/dL	13.5 - 18.0
INTERPRETATION: Haemoglobin values vary in Men blood loss, renal failure etc. Higher values are often due t			
PCV (Packed Cell Volume) / Haematocrit (EDTA Blood/Derived)	48.4	%	42 - 52
RBC Count (EDTA Blood/Automated Blood cell Counter)	5.45	mill/cu.mm	4.7 - 6.0
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	89.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	31.3	pg	27 - 32
MCHC (Mean Corpuscular Haemoglobin concentration) (EDTA Blood/Derived)	35.2	g/dL	32 - 36
RDW-CV (Derived)	11.56	%	11.5 - 16.0
RDW-SD (Derived)	36.01	fL	39 - 46
Total WBC Count (TC) (EDTA Blood/Derived from Impedance)	7720	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	55	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	33	%	20 - 45



Name	: Mr. MALAGRI PUNDLIK		
PID No.	: MED111087640	Register On : 14/05/2022 9:45 AM	\mathbf{M}
SID No.	: 712214641	Collection On : 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On : 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On : 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	05	%	01 - 06
Monocytes (Blood/Impedance Variation & Flow Cytometry)	07	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	00	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.25	10^3 / µl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.55	10^3 / µl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.39	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.54	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.00	10^3 / µl	< 0.2
Platelet Count (EDTA Blood/Derived from Impedance)	254	10^3 / µl	150 - 450
MPV (Blood/Derived)	08.83	fL	7.9 - 13.7
PCT	0.22	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Automated ESR analyser)	06	mm/hr	< 15



Name	: Mr. MALAGRI PUNDLIK		
PID No.	: MED111087640	Register On : 14/05/2022 9:45 AM	\mathbf{C}
SID No.	: 712214641	Collection On : 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On : 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On : 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.7	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.50	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.0	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.4	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.60	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.69		1.1 - 2.2
INTERPRETATION: Remark : Electrophoresis is the providence of the second	referred method		
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC / Kinetic)	42	U/L	5 - 40
Remark: Kindly correlate clinically.			
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	66	U/L	5 - 41
Remark: Kindly correlate clinically.			
Alkaline Phosphatase (SAP) (Serum/PNPP / Kinetic)	80	U/L	53 - 128
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	36	U/L	< 55



Name	: Mr. MALAGRI PUNDLIK		
PID No.	: MED111087640	Register On : 14/05/2022 9:45 AM	\mathbf{O}
SID No.	: 712214641	Collection On : 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On : 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On : 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Lipid Profile			
Cholesterol Total (Serum/Oxidase / Peroxidase method)	141	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	81	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	34	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
Remark: Kindly correlate clinically.			
LDL Cholesterol (Serum/Calculated)	90.8	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	16.2	mg/dL	< 30
	K	7	



Name	: Mr. MALAGRI PUNDLIK			
PID No.	: MED111087640	Register On	: 14/05/2022 9:45 AM	M
SID No.	: 712214641	Collection On	: 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On	: 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On	: 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel			

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Non HDL Cholesterol (Serum/ <i>Calculated</i>)	107.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >=220

INTERPRETATION: 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	4.1	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/ <i>Calculated</i>)	2.4	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.7	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0



Name	: Mr. MALAGRI PUNDLIK			
PID No.	: MED111087640	Register On	: 14/05/2022 9:45 AM	\mathbf{C}
SID No.	: 712214641	Collection On	: 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On	: 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On	: 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel			

Investigation Glycosylated Haemoglobin (HbA1c)	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval		
HbA1C (Whole Blood/HPLC)	5.2	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5		
INTERPRETATION: If Diabetes - Good control : 6.1 - 7.0 %, Fair control : 7.1 - 8.0 %, Poor control >= 8.1 %					

Estimated Average Glucose	102.54	mg/dL
Estimated Trefage Glacose	10110	8

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



APPROVED BY

Name	: Mr. MALAGRI PUNDLIK		
PID No.	: MED111087640	Register On : 14/05/2022 9:45 AM	\mathbf{C}
SID No.	: 712214641	Collection On : 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On : 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On : 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
IMMUNOASSAY			
<u>THYROID PROFILE / TFT</u>			
T3 (Triiodothyronine) - Total (Serum/Chemiluminescent Immunometric Assay (CLIA)) INTERPRETATION: Comment : Total T3 variation can be seen in other condition like pres	1.06 gnancy, drugs, neph	ng/ml rosis etc. In such cases,	0.7 - 2.04 Free T3 is recommended as it is
Metabolically active. T4 (Thyroxine) - Total (Serum/ <i>Chemiluminescent Immunometric Assay</i> (<i>CLIA</i>))	10.20	Microg/dl	4.2 - 12.0
INTERPRETATION: Comment : Total T4 variation can be seen in other condition like pre- Metabolically active.	gnancy, drugs, neph	rosis etc. In such cases,	Free T4 is recommended as it is
TSH (Thyroid Stimulating Hormone) (Serum/Chemiluminescent Immunometric Assay (CLIA))	2.089	µIU/mL	0.35 - 5.50
INTERPRETATION: Reference range for cord blood - upto 20 1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0 (Indian Thyroid Society Guidelines) Comment : 1.TSH reference range during pregnancy depends on Iod 2.TSH Levels are subject to circadian variation, reaching of the order of 50%,hence time of the day has influence of 3.Values&lt0.03 μIU/mL need to be clinically correl	peak levels betwee on the measured serv	n 2-4am and at a minim am TSH concentrations	um between 6-10PM. The variation can be



Name	: Mr. MALAGRI PUNDLIK		
PID No.	: MED111087640	Register On : 14/05/2022 9:45 AM	\mathbf{C}
SID No.	: 712214641	Collection On : 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On : 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On : 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
CLINICAL PATHOLOGY			
PHYSICAL EXAMINATION			
Colour (Urine/Physical examination)	Pale yellow		Yellow to Amber
Volume (Urine/Physical examination)	30		ml
Appearance (Urine)	Clear		
CHEMICAL EXAMINATION			
pH (Urine)	6.0		4.5 - 8.0
Specific Gravity (Urine/Dip Stick o''Reagent strip method)	1.015		1.002 - 1.035
Protein (Urine/Dip Stick ó"Reagent strip method)	Negative		Negative
Glucose (Urine)	Nil		Nil
Ketone (Urine/Dip Stick ó"Reagent strip method)	Nil		Nil
Leukocytes (Urine)	Negative	leuco/uL	Negative
Nitrite (Urine/Dip Stick ó"Reagent strip method)	Nil		Nil
Bilirubin (Urine)	Negative	mg/dL	Negative



Name	: Mr. MALAGRI PUNDLIK		
PID No.	: MED111087640	Register On : 14/05/2022 9:45 AM	\mathbf{C}
SID No.	: 712214641	Collection On : 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On : 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On : 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Blood	Nil		Nil
(Urine)			
Urobilinogen	Normal		Within normal limits
(Urine/Dip Stick ó"Reagent strip method)			
<u>Urine Microscopy Pictures</u>			
RBCs	Nil	/hpf	NIL
(Urine/Microscopy)			
Pus Cells	3-4	/hpf	< 5
(Urine/Microscopy)			
Epithelial Cells	3-4	/hpf	No ranges
(Urine/Microscopy)			
Others	Nil		Nil
(Urine)			

Dr Shouree K.R MBBS MD DNB Consultant Pathologist Reg No : KMC 103138 APPROVED BY

Name	:	Mr. MALAGRI PUNDLIK					
PID No.	:	MED111087640	Register On	:	14/05/2022 9:45 AM	\mathbf{C}	
SID No.	:	712214641	Collection On	:	14/05/2022 11:22 AM		
Age / Sex	:	31 Year(s) / Male	Report On	:	14/05/2022 4:16 PM	MEDALL	
Туре	:	OP	Printed On	:	04/06/2022 12:32 PM		
Ref. Dr	:	MediWheel					

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
<u>Stool Analysis - ROUTINE</u>			
Colour (Stool)	Brownish		Brown
Blood (Stool)	Not present		Not present
Mucus (Stool)	Not present		Not present
Reaction (Stool)	Alkaline		Alkaline
Consistency (Stool)	Semi solid		Semi solid
Ova (Stool)	Nil		Nil
Others (Stool)	Nil		Nil
Cysts (Stool)	Nil		Nil
Trophozoites (Stool)	Nil		Nil
RBCs (Stool)	Nil	/hpf	Nil
Pus Cells (Stool)	2-3	/hpf	Nil
Macrophages (Stool)	Nil		Nil
Epithelial Cells (Stool)	Nil	/hpf	Nil



Name	: Mr. MALAGRI PUNDLIK			
PID No.	: MED111087640	Register On	: 14/05/2022 9:45 AM	\mathbf{C}
SID No.	: 712214641	Collection On	: 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On	: 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On	: 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel			

Investigation

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination) Remark: Test to be confirmed by Gel method. 'O' 'Positive'

Observed

<u>Value</u>



<u>Unit</u>

Biological Reference Interval

APPROVED BY

Name	: Mr. MALAGRI PUNDLIK			
PID No.	: MED111087640	Register On	: 14/05/2022 9:45 AM	\mathbf{C}
SID No.	: 712214641	Collection On	: 14/05/2022 11:22 AM	
Age / Sex	: 31 Year(s) / Male	Report On	: 14/05/2022 4:16 PM	MEDALL
Туре	: OP	Printed On	: 04/06/2022 12:32 PM	
Ref. Dr	: MediWheel			

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
BIOCHEMISTRY			
BUN / Creatinine Ratio	11.7		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	84	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Urine sugar, Fasting	Nil	Nil	
(Urine - F)			
Glucose Postprandial (PPBS)	124	mg/dL	70 - 140
(Plasma - PP/GOD - POD)			

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Sugar (PP-2 hours) (Urine - PP)	Negative	Negative		
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	11.7	mg/dL	7.0 - 21	
Creatinine	1.0	mg/dL	0.9 - 1.3	

(Serum/Jaffe Kinetic)

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid			5.1	mg/dL	3.5 - 7.2	
(0			• • •			

(Serum/Uricase/Peroxidase)



-- End of Report --