



BHARM010672181

CLIENT CODE: C000138394

CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT ID:

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP

ACCESSION NO: **0181VK000873** AGE: 50 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 19/11/2022 09:54 REPORTED: 19/11/2022 21:48

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Final Results Biological Reference Interval Units

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE 40 MALE

BLOOD COUNTS,EDTA WHOLE BLOOD						
HEMOGLOBIN (HB)	10.9	Low	13.0 - 17.0	g/dL		
METHOD : SLS- HEMOGLOBIN DETECTION METHOD						
RED BLOOD CELL (RBC) COUNT	3.12	Low	4.5 - 5.5	mil/μL		
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION						
WHITE BLOOD CELL (WBC) COUNT	3.59	Low	4.0 - 10.0	thou/µL		
METHOD: FLUORESCENCE FLOW CYTOMETRY						
PLATELET COUNT	240		150 - 410	thou/µL		
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION						
RBC AND PLATELET INDICES						
HEMATOCRIT (PCV)	33.1	Low	40.0 - 50.0	%		
METHOD : CUMULATIVE PULSE HEIGHT DETECTION METHOD						
MEAN CORPUSCULAR VOLUME (MCV)	106.1	High	83.0 - 101.0	fL		
METHOD: CALCULATED FROM RBC & HCT						
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	34.9	High	27.0 - 32.0	pg		
METHOD: CALCULATED FROM THE RBC & HGB						
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	32.9		31.5 - 34.5	g/dL		
METHOD : CALCULATED FROM THE HGB & HCT						
RED CELL DISTRIBUTION WIDTH (RDW)	15.8	High	11.6 - 14.0	%		
METHOD: CALCULATED FROM RBC SIZE DISTRIBUTION CURVE						
MENTZER INDEX	34.0					
MEAN PLATELET VOLUME (MPV)	11.1	High	6.8 - 10.9	fL		
METHOD : CALCULATED FROM PLATELET COUNT & PLATELET HEMATOCRIT						
WBC DIFFERENTIAL COUNT						
NEUTROPHILS	65		40 - 80	%		
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING						
LYMPHOCYTES	27		20 - 40	%		
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING						
MONOCYTES	5		2 - 10	%		
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING						
EOSINOPHILS	3		1 - 6	%		
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING						
ABSOLUTE NEUTROPHIL COUNT	2.32		2.0 - 7.0	thou/µL		
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING						
		-				

0.98

Low 1.0 - 3.0



ABSOLUTE LYMPHOCYTE COUNT



thou/µL





CLIENT CODE: C000138394

CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110020

NEW DELHI 110030 DELHI INDIA

8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP PATIENT ID: BHARM010672181

ACCESSION NO: **0181VK000873** AGE: 50 Years SEX: Male ABHA NO:

RECEIVED: 19/11/2022 09:54 19/11/2022 21:48 DRAWN: REPORTED:

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results		Biological Reference Interva	al Units
METHOD : FLOW CYTOMETR	Y WITH LIGHT SCATTERING				
ABSOLUTE MONOCYTE		0.17	Low	0.2 - 1.0	thou/µL
METHOD : FLOW CYTOMETRY	Y WITH LIGHT SCATTERING				,,
ABSOLUTE EOSINOPHI	L COUNT	0.09		0.02 - 0.50	thou/µL
METHOD : FLOW CYTOMETRY	Y WITH LIGHT SCATTERING				
NEUTROPHIL LYMPHOO	CYTE RATIO (NLR)	2.4			
MORPHOLOGY					
RBC		MACROCYTOSIS			
WBC		LEUCOPENIA			
METHOD : MICROSCOPIC EX	(AMINATION				
PLATELETS		ADEQUATE			
ERYTHROCYTE SEDI	MENTATION RATE (ESR),WHOLE			
E.S.R		11		< 15	mm at 1 hr
GLYCOSYLATED HEM BLOOD	IOGLOBIN(HBA1C), EDT	A WHOLE			
HBA1C METHOD: HPLC		4.6		Non-diabetic Adult < 5.7 Pre-diabetes 5.7 - 6.4 Diabetes diagnosis: > or = 6.5 Therapeutic goals: < 7.0 Action suggested : > 8.0 (ADA Guideline 2021)	%
ESTIMATED AVERAGE	GLUCOSE(FAG)	85.3		< 116.0	mg/dL
METHOD : CALCULATED PAR	` ,	00.0			9, ==
GLUCOSE FASTING,F	LUORIDE PLASMA				
FBS (FASTING BLOOD	SUGAR)	96		Normal 75 - 99 Pre-diabetics: 100 - 125 Diabetic: > or = 126	mg/dL
METHOD : ENZYMATIC REFE	RENCE METHOD WITH HEXOKINAS	E			
GLUCOSE, POST-PRA	NDIAL, PLASMA				
PPBS(POST PRANDIAL	BLOOD SUGAR)	93		70 - 139	mg/dL
	RENCE METHOD WITH HEXOKINAS	E			
LIPID PROFILE, SER	UM				
CHOLESTEROL, TOTAL		148		Desirable cholesterol level < 200 Borderline high cholesterol 200 - 239 High cholesterol > / = 240	mg/dL

METHOD: ENZYMATIC COLORIMETRIC ASSAY









CLIENT CODE: C000138394

CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

ABHA NO:

REPORTED:

PATIENT NAME: BHARAT P JAGTAP

PATIENT ID:

BHARM010672181

DRAWN:

ACCESSION NO: **0181VK000873**

AGE: 50 Years SEX: Male

RECEIVED: 19/11/2022 09:54

19/11/2022 21:48

REFERRING DOCTOR: SELF

CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
TRIGLYCERIDES	80	Normal: < 150 mg/dL Borderline high: 150 - 199 High: 200 - 499 Very High: >/= 500
METHOD: ENZYMATIC COLORIMETRIC ASSAY	27	Law Low UDI Chalastaval (40 mag/di
HDL CHOLESTEROL	37	Low Low HDL Cholesterol <40 mg/dL
METHOD , ENZYMATIC COLODIMETRIC		High HDL Cholesterol >/= 60
METHOD: ENZYMATIC, COLORIMETRIC CHOLESTEROL LDL	95	Adult levels: mg/dL Optimal < 100 Near optimal/above optimal: 100- 129 Borderline high : 130-159 High : 160-189 Very high : = 190
METHOD: ENZYMATIC COLORIMETRIC ASSAY		very mgm i 150
NON HDL CHOLESTEROL	111	Desirable : < 130 mg/dL Above Desirable : 130 -159 Borderline High : 160 - 189 High : 190 - 219 Very high : > / = 220
CHOL/HDL RATIO	4.0	Low Risk : 3.3 - 4.4 Average Risk : 4.5 - 7.0 Moderate Risk : 7.1 - 11.0 High Risk : > 11.0
LDL/HDL RATIO	2.6	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk
VERY LOW DENSITY LIPOPROTEIN	16.0	< OR = 30.0 mg/dL
LIVER FUNCTION PROFILE, SERUM		
BILIRUBIN, TOTAL METHOD: COLORIMETRIC DIAZO	1.08	Upto 1.2 mg/dL
BILIRUBIN, DIRECT	0.30	< 0.30 mg/dL
BILIRUBIN, INDIRECT	0.78	0.1 - 1.0 mg/dL
TOTAL PROTEIN METHOD: COLORIMETRIC	7.8	6.0 - 8.0 g/dL
ALBUMIN METHOD: COLORIMETRIC	4.9	3.97 - 4.94 g/dL
GLOBULIN	2.9	2.0 - 3.5 g/dL
ALBUMIN/GLOBULIN RATIO	1.7	1.0 - 2.1 RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT) METHOD: UV ABSORBANCE	19	< OR = 50 U/L

METHOD: UV ABSORBANCE



Page 3 Of 10

Scan to View Report





CLIENT CODE: C000138394

REFERRING DOCTOR: SELF

CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030

DELHI INDIA 8800465156 SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

REPORTED:

PATIENT NAME: BHARAT P JAGTAP

PATIENT ID:

BHARM010672181

ACCESSION NO: **0181VK000873**

AGE: 50 Years

SEX: Male ABHA NO:

19/11/2022 21:48

DRAWN:

RECEIVED: 19/11/2022 09:54

CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units
ALANINE AMINOTRANCEDACE (ALT/CORT)	10	< OR = 50	11/1
ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: UV ABSORBANCE	19	< UK = 50	U/L
ALKALINE PHOSPHATASE	48	40 - 129	U/L
METHOD: COLORIMETRIC			
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: ENZYMATIC, COLORIMETRIC	8	0 - 60	U/L
LACTATE DEHYDROGENASE METHOD: UV ABSORBANCE	205	125 - 220	U/L
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	10	6 - 20	mg/dL
METHOD: ENZYMATIC ASSAY			
CREATININE, SERUM			
CREATININE	0.80	0.7 - 1.2	mg/dL
METHOD : COLORIMETRIC			
BUN/CREAT RATIO			
BUN/CREAT RATIO	12.50	8.0 - 15.0	
URIC ACID, SERUM			
URIC ACID	5.7	3.4 - 7.0	mg/dL
METHOD: ENZYMATIC COLORIMETRIC ASSAY			
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	7.8	6.0 - 8.0	g/dL
METHOD: COLORIMETRIC			
ALBUMIN, SERUM			
ALBUMIN	4.9	3.97 - 4.94	g/dL
METHOD: COLORIMETRIC			
GLOBULIN			
GLOBULIN	2.9	2.0 - 3.5	g/dL
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM, SERUM	138	136 - 145	mmol/L
POTASSIUM, SERUM	4.90	3.5 - 5.1	mmol/L
CHLORIDE, SERUM	103	98 - 107	mmol/L
PHYSICAL EXAMINATION, URINE			
COLOR	PALE YELLOW		
APPEARANCE	CLEAR		
CHEMICAL EXAMINATION, URINE			
PH	6.0	5.00 - 7.50	











CLIENT CODE: C000138394

CLIENT'S NAME AND ADDRESS:
ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110020

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP

PATIENT ID: BHARM010672181

ACCESSION NO: **0181VK000873** AGE: 50 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 19/11/2022 09:54 REPORTED: 19/11/2022 21:48

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interva	l Units	
SPECIFIC GRAVITY	1.015	1.010 - 1.030		
PROTEIN	NOT DETECTED	NOT DETECTED		
GLUCOSE	NOT DETECTED	NOT DETECTED		
KETONES	NOT DETECTED	NOT DETECTED		
BLOOD	NOT DETECTED	NOT DETECTED		
UROBILINOGEN	NORMAL	NORMAL		
NITRITE	NOT DETECTED	NOT DETECTED		
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED		
MICROSCOPIC EXAMINATION, URINE				
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF	
PUS CELL (WBC'S)	1-2	0-5	/HPF	
EPITHELIAL CELLS	0-1	0-5	/HPF	
CASTS	NOT DETECTED			
CRYSTALS	NOT DETECTED			
BACTERIA	NOT DETECTED	NOT DETECTED		
YEAST	NOT DETECTED	NOT DETECTED		
THYROID PANEL, SERUM				
ТЗ	92.8	80 - 200	ng/dL	
METHOD: ELECTROCHEMILUMINESCENCE				
T4	6.35	5.1 - 14.1	µg/dL	
METHOD: ELECTROCHEMILUMINESCENCE				
TSH (ULTRASENSITIVE)	1.800	0.27 - 4.2	μIU/mL	
METHOD: ELECTROCHEMILUMINESCENCE				



Scan to View Report





CLIENT CODE: C000138394 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA**

8800465156

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN-U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP PATIENT ID: BHARM010672181

ACCESSION NO: 0181VK000873 AGE: 50 Years SEX: Male ABHA NO:

RECEIVED: 19/11/2022 09:54 DRAWN: REPORTED: 19/11/2022 21:48

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units <u>Final</u>

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyporthyroidism, TSH levels are low. owidctlparowidctlparBelow mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

STOOL: OVA & PARASITE

COLOUR SAMPLE NOT RECEIVED

METHOD: VISUAL

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE AB

METHOD: GEL COLUMN AGGLUTINATION METHOD.

RH TYPE **NEGATIVE**









CLIENT CODE: C000138394

CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030**

DELHI INDIA 8800465156 SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN-U74899PB1995PLC045956

PATIENT ID:

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP

ABHA NO:

BHARM010672181

ACCESSION NO: 0181VK000873 AGE: 50 Years SEX: Male

DRAWN: RECEIVED: 19/11/2022 09:54 REPORTED: 19/11/2022 21:48

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

METHOD: GEL COLUMN AGGLUTINATION METHOD.

XRAY-CHEST	RESULT PENDING
TMT OR ECHO	RESULT PENDING
ECG	RESULT PENDING
MEDICAL HISTORY	RESULT PENDING
ANTHROPOMETRIC DATA & BMI	RESULT PENDING
GENERAL EXAMINATION	RESULT PENDING
CARDIOVASCULAR SYSTEM	RESULT PENDING
RESPIRATORY SYSTEM	RESULT PENDING
PER ABDOMEN	RESULT PENDING
CENTRAL NERVOUS SYSTEM	RESULT PENDING
MUSCULOSKELETAL SYSTEM	RESULT PENDING
BASIC EYE EXAMINATION	RESULT PENDING
BASIC ENT EXAMINATION	RESULT PENDING
BASIC DENTAL EXAMINATION	RESULT PENDING
SUMMARY	RESULT PENDING
FITNESS STATUS	RESULT PENDING

BLOOD COUNTS, EDTA WHOLE BLOODThe cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-

Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-

The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504

This ratio element is a calculated parameter and out of NABL scope. ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION:

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy,

Estrogen medication, Aging.
Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS



Page 7 Of 10 Scan to View Report





CLIENT CODE: C000138394

CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA**

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP

PATIENT ID:

BHARM010672181

8800465156

ACCESSION NO: 0181VK000873

AGE: 50 Years SEX: Male ABHA NO:

REPORTED:

19/11/2022 21:48

DRAWN:

RECEIVED: 19/11/2022 09:54

CLIENT PATIENT ID:

Test Report Status

REFERRING DOCTOR: SFLF

<u>Final</u>

Results

Biological Reference Interval Units

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine, salicylates)

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- Diagnosing diabetes.
 Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range. 1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.

- 2. eAG gives an evaluation of blood glucose levels for the last couple of months.
 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to :I.Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic

anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

III. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results. IV.Interference of hemoglobinopathies in HbA1c estimation is seen in

a.Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b.Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy GLUCOSE FASTING,FLUORIDE PLASMA-**TEST DESCRIPTION**

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine.

Increased in

Diabetes mellitus, Cushing's syndrome (10 - 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in

Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency, hypopituitarism,diffuse liver disease, malignancy (adrenocortical, stomach,fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia),Drugs- insulin, ethanol, propranolol sulfonylureas,tolbutamide, and other oral hypoglycemic agents.

NOTE:

Hypoglycemia is defined as a glucoseof < 50 mg/dL in men and < 40 mg/dL in women.

While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin (HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic

index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.
GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin

treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic

hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparatyrioldism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also











CLIENT CODE: C000138394

CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP

PATIENT ID:

BHARM010672181

ACCESSION NO: 0181VK000873

AGE: 50 Years

SEX: Male ABHA NO:

REPORTED:

19/11/2022 21:48

DRAWN:

RECEIVED: 19/11/2022 09:54

CLIENT PATIENT ID:

Test Report Status

REFERRING DOCTOR: SFLF

<u>Final</u>

Results

Biological Reference Interval Units

known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular

permeability or decreased lymphatic clearance,malnutrition and wasting etc BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH. CREATININE, SERUM-Higher than normal level may be due to:

- Blockage in the urinary tract
- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
 Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis
- Muscular dystrophy URIC ACID, SERUM-

Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic syndrome Causes of decreased levels-Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome,Protein-losing enteropathy etc. ALBUMIN, SERUM-

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc.

ABO GROUP & RH TYPE, EDTA WHOLE BLOODBlood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.



Page 9 Of 10 Scan to View Report







CLIENT CODE: C000138394 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN-U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: BHARAT P JAGTAP

PATIENT ID:

BHARM010672181

ACCESSION NO: 0181VK000873

AGE: 50 Years SEX: Male ABHA NO: REPORTED:

19/11/2022 21:48

DRAWN:

RECEIVED: 19/11/2022 09:54

CLIENT PATIENT ID:

Test Report Status

<u>Final</u>

Results

Units

MEDI WHEEL FULL BODY HEALTH CHECK UP ABOVE ASOLMIANTENDING

ULTRASOUND ABDOMEN

REFERRING DOCTOR: SELF

RESULT PENDING

End Of Report Please visit www.srlworld.com for related Test Information for this accession

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- 8. Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062





Scan to View Report