

Visit ID : MBAR39301	Registration : 11/Mar/2023 01:31PM
UHID/MR No : ABAR.0000039289	Collected : 11/Mar/2023 08:53PM
Patient Name : Mrs.AAKRATI	Received : 11/Mar/2023 08:53PM
Age/Gender : 31 Y 0 M 0 D /F	Reported : 11/Mar/2023 08:54PM
Ref Doctor : Dr.NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240,Sanjay Nagar Bareilly (UP)	Barcode No : A3619646

DEPARTMENT OF HORMONE ASSAYS

Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

THYROID PROFILE (T3,T4,ULTRASENSITIVE TSH)

Sample Type : SERUM

T3	1.00	ng/ml	0.61-1.81	CLIA
T4	7.3	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	0.976	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

- Serum T3, T4 and TSH are the measurements form three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
- Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
- Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
- Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propanolol and propylthiouracil.
- Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
- Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
- Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
- TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

9. REFERENCE RANGE:

PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 – 2.500
2nd Trimester	0.200 – 3.000
3rd Trimester	0.300 – 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

- During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
- TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***



 Dr. Miti Gupta
 DNB ; MD [Pathology]


Visit ID : MBAR39301	Registration : 11/Mar/2023 01:31PM
UHID/MR No : ABAR.0000039289	Collected : 11/Mar/2023 08:53PM
Patient Name : Mrs.AAKRATI	Received : 11/Mar/2023 08:53PM
Age/Gender : 31 Y 0 M 0 D /F	Reported : 11/Mar/2023 08:54PM
Ref Doctor : Dr.NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240,Sanjay Nagar Bareilly (UP)	Barcode No : A3619646

DEPARTMENT OF HORMONE ASSAYS

Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

THYROID PROFILE (T3,T4,ULTRASENSITIVE TSH)

Sample Type : SERUM

T3	1.00	ng/ml	0.61-1.81	CLIA
T4	7.3	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	0.976	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

- Serum T3, T4 and TSH are the measurements form three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
- Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
- Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
- Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propanolol and propylthiouracil.
- Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
- Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
- Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
- TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

9. REFERENCE RANGE:

PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 – 2.500
2nd Trimester	0.200 – 3.000
3rd Trimester	0.300 – 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

- During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
- TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***



 Dr. Miti Gupta
 DNB ; MD [Pathology]





Pat. BAKSHI

11/3/23

डॉ. निरमल अग्रवाल
हृदय रोग विशेषज्ञ

Pat. ID.....



CARDIART 6108T

10mm/mV 25mm/sec 25Hz

BPL CARDIART 6108T

10mm/mV 25mm/sec 25Hz


BPL CARDIART

V3

V4

V5

V6



Pat. ID.

Pat. ID.

CARDIART

Dr. Nitin Agarwal

MD, DM (Cardiology)

Consultant Interventional Cardiologist

Cell +91-98578 33777

Formerly at

Escorts Heart Institute & Research Centre, Delhi

Dr. Ram Manohar Lohia Hospital, Delhi



APPLE
CARDIAC CARE

DR. NITIN AGARWAL'S HEART CLINIC

A Agarwal

Amman

11/3/20

12/10/20

26/10

20

Amman 10/10/20

0

0

A-3, EKTA NAGAR, (OPP CARE HOSPITAL) STADIUM ROAD, NEAR DELAPEER CHAURAHA, BAREILLY - 243 122 (U.P)

OPD Timings : 12.00 Noon to 04.00 pm, Sunday : 12.00 Noon to 3.00 pm

संख्या लगाने के लिए फोन करें : 09458888448, 07599031977

VALID FOR 5 DAYS.

पचास पाँच दिन के लिये मान्य





PARAS MRI & ULTRASOUND CENTRE

MOST ADVANCED 32 CHANNEL 3T 3D WHOLE BODY MRI

261, ASHAPURAM, OPP. DR. BASU EYE HOSPITAL, STADIUM ROAD, BAREILLY

• Helpline : 7300761761 • E-mail : parasmribly@gmail.com

REPORT

4D / 5D ULTRASOUND

COLOR DOPPLER

TVS/ TRUS

MUSCULOSKELETAL USG

Date : 11.03.2023
Name : AAKRATI AGARWAL 31Y/F
Ref.By : DR APPLE CARDIAC CARE

ULTRASOUND WHOLE ABDOMEN

LIVER - Liver is normal in size and outline. It shows increased echogenicity. No obvious focal pathology is seen. The intra hepatic biliary radicals are not dilated. PV -5.0 mm

GALL BLADDER -Gall Bladder is normal in size, has normal wall thickness with no evidence of calculi. Fat planes between GB and liver are well maintained.The CBD appears normal.

PANCREAS - Pancreas is normal in size and echogenicity. Its outlines are distinct. No obvious focal lesion, calcification or ductile dilatation is seen.

SPLEEN - Spleen is normal in size and echogenicity. There is no evidence of collaterals

KIDNEYS - Both kidneys are normal in position, outline and echogenicity. No evidence of calculi is seen. CMD is maintained. No evidence of hydronephrosis is seen on either side.

URINARY BLADDER -Urinary Bladder is normal in size and outline. There is no evidence of any obvious intraluminal or paramedical pathology. Wall is not thickened.

Uterus - Uterus is anteverted and normal in size. The myometrial and endometrial echoes are normal. The endometrial thickness measures 7 mm. No focal lesion is seen.

Both ovaries are normal in size and shows uniform parenchymal echogenicity and smooth outlines. No evidence of cyst or mass is seen.

No evidence of ascites /pleural effusion/adenopathy is seen. Bowel loops are not dilated. Bilateral iliac fossa appears normal.

IMPRESSION:


❖ Grade I fatty liver.






Adv clinical correlation.


Dr. Puja Tripathi

M.B.B.S., M.D.

MBBS, MD (Radiodiagnosis, SGPGI)







NAME	Ms. AAKARTI AGARWAL	AGE/SEX	31 Y/F
Reff. By	Dr. NITIN AGARWAL (DM)	DATE	11/03/2023

ECHOCARDIOGRAPHY AND COLOUR DOPPLER STUDY

<u>MEASUREMENTS</u>	<u>VALUE</u>	<u>NORMAL DIMENSIONS</u>
LVID (d)	4.5	cm (3.7 –5.6 cm)
LVID (s)	2.4	cm (2.2 –3.9 cm)
RVID (d)	2.4	cm (0.7 –2.5 cm)
IVS (ed)	1.0	cm (0.6 –1.1 cm)
LVPW (ed)	1.0	cm (0.6 –1.1 cm)
AO	2.2	cm (2.2 –3.7 cm)
LA	3.0	cm (1.9 –4.0 cm)
<u>LV FUNCTION</u>		
EF	60	% (54 –76 %)
FS	30	% (25 –44 %)

LEFT VENTRICLE : No regional wall motion abnormality
No concentric left Ventricle Hypertrophy

MITRAL VALVE : Thin, PML moves posteriorly during Diastole
No SAM, No Subvalvular pathology seen.
No mitral valve prolapse calcification .

TRICUSPID VALVE : Thin, opening wells. No calcification, No doming .
No Prolapse.
Tricuspid inflow velocity= 0.7 m/sec

AORTIC VALVE : Thin, tricuspid, opening well, central closer,
no flutter.
No calcification
Aortic velocity = 1.3 m/sec

PULMONARY VALVE : Thin, opening well, Pulmonary artery is normal
EF slope is normal.
Pulmonary Velocity = 0.9 m /sec



ON DOPPLER INTERROGATION THERE WAS :

- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

MITRAL FLOW E= 0.8 m/sec A= 0.6 m/sec

ON COLOUR FLOW:

- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

COMMENTS:

- No LA /LV clot
- No pericardial effusion
- No intracardiac mass
- IAS/IVS Intact
- Inferior vena cava – normal in size with normal respiratory variation

FINAL IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY
- NORMAL LV DIASTOLIC FUNCTION
- NORMAL LV SYSTOLIC FUNCTION (LVEF~60%)
- NORMAL CARDIAC CHAMBER DIMENSIONS
- NORMAL VALVULAR COLOUR FLOW PATTERN



DR. NITIN AGARWAL
DM (Cardiology)
Consultant Cardiologist

This opinion is to be correlated with the clinically findings and if required, please re-evaluate / re-evaluate with further investigation.

ON DOPPLER INTERROGATION THERE WAS :

- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

MITRAL FLOW

E= 0.8 m/sec

A= 0.6 m/sec

ON COLOUR FLOW:

- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

COMMENTS:

- No LA /LV clot
- No pericardial effusion
- No intracardiac mass
- IAS/IVS Intact
- Inferior vena cava – normal in size with normal respiratory variation

FINAL IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY
- NORMAL LV DIASTOLIC FUNCTION
- NORMAL LV SYSTOLIC FUNCTION (LVEF~60%)
- NORMAL CARDIAC CHAMBER DIMENSIONS
- NORMAL VALVULAR COLOUR FLOW PATTERN



DR. NITIN AGARWAL
DM (Cardiology)
Consultant Cardiologist

This opinion is to be correlated with the clinically findings and if required, please re-evaluate / reconfirm with further investigation.



॥ ॐ गणेशाय नमः ॥

GANESH DIAGNOSTIC

DR. LOKESH GOYAL
MBBS (KGMCI), MD (RADIOLOGY)
CONSULTANT INTERVENTIONAL RADIOLOGIST
FORMER SR REGISTRAR - APOLLO HOSPITAL, NEW DELHI
LIFE MEMBER OF RIA

Timings : 9:00 am to 9:00 pm, Sunday 9.00 am to 3.00 pm

☎ 8392957683, 6395228718

MR. AAKRATI AGARWAL
DR. NITIN AGARWAL, DM

11-03-2023

REPORT

EXAMINATION PERFORMED: X-RAY CHEST

R/L Lung fields are clear

Both of the CP angles are clear.

Both hila show a normal pattern

Cardiac and mediastinal borders appear normal.

Visualized bony thorax and soft tissue of the chest wall appear normal.

IMPRESSION ---NO SIGNIFICANT ABNORMALITY IS SEEN

Not for medico-legal purpose

DR LOKESH GOYAL
MD
RADIOLOGIST



॥ ॐ गणेशाय नमः ॥
DR. NITIN AGARWAL, DM

**NOT VALID FOR
MEDICO LEGAL PURPOSE**

A Venture of Apple Cardiac Care

A-3, Ekta Nagar, Stadium Road,
 (Opp. Care Hospital),
 Bareilly - 243 122 (U.P.) India
 Tel. : 07599031977, 09458888448



APPLE
PATHOLOGY
 TRUSTED RESULT

Reg.NO. : 111
 NAME : **Mrs. AAKRATI AGARWAL**
 REFERRED BY : Dr.Nitin Agarwal (D M)
 SAMPLE : BLOOD

DATE : **11/03/2023**
 AGE : 31 Yrs.
 SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
HAEMATOLOGY			
COMPLETE BLOOD COUNT (CBC)	12.6	gm/dl	12.0-15.0
HAEMOGLOBIN	8,600	/cumm	4,000-11,000
TOTAL LEUCOCYTE COUNT	63	%	40-75
DIFFERENTIAL LEUCOCYTE COUNT(DLC)	34	%	20-45
Neutrophils	03	%	01-08
Lymphocytes	00	%	01-06
Eosinophils	00	%	00-02
Monocytes	4.09	million/cumm	3.5-6.5
Basophils	38.0	%	35-54
TOTAL R.B.C. COUNT	92.9	fL	76-96
P.C.V./ Haematocrit value	30.8	pg	27.00-32.00
M C V	33.2	g/dl	30.50-34.50
M C H	2.93	lacs/mm ³	1.50 - 4.50
M C H C	15	mm	00- 20
PLATELET COUNT			
E.S.R (WINTROBE METHOD)			
-in First hour			
BIOCHEMISTRY			
BLOOD SUGAR F.	76	mg/dl	60-100
HAEMATOLOGY			

Report is not valid for medicolegal purpose

Reg.NO. : 111	DATE : 11/03/2023
NAME : Mrs. AAKRATI AGARWAL	AGE : 31 Yrs.
REFERRED BY : Dr. Nitin Agarwal (D M)	SEX : FEMALE
SAMPLE : BLOOD	

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
GLYCOSYLATED HAEMOGLOBIN	5.7		

EXPECTED RESULTS :

Non diabetic patients	: 4.0% to 6.0%
Good Control	: 6.0% to 7.0%
Fair Control	: 7.0% to -8%
Poor Control	: Above 8%

***ADA: American Diabetes Association**

The glycosylated hemoglobin assay has been validated as a reliable indicator of mean blood glucose levels for a period of 8-12 week period prior to HBA1C determination. ADA recommends the testing twice a year in patients with stable blood glucose, and quarterly, if treatment changes, or if blood glucose levels are unstable.

METHOD : ADVANCED IMMUNO ASSAY.

BIOCHEMISTRY

Gamma Glutamyl Transferase (GGT)	29	U/L	11-50
----------------------------------	----	-----	-------

HAEMATOLOGY

Blood Group	O+
Rh	POSITIVE

BIOCHEMISTRY

BLOOD UREA	19	mg/dL.	10-40
------------	----	--------	-------

* Low serum urea is usually associated with status of overhydration severe hepatic failure.

* A urea level of 10-45 mg/dl indicates normal glomerular function and a level of 100-250 mg/dl indicates a serious impairment of renal function. In chronic renal failure, urea correlates better with the symptoms of uremia than does serum creatinine.

* Urine/Serum urea is more than 9 in prerenal and less than 3 in renal uremia.

Report is not valid for medicolegal purpose

Reg.NO. : 111
NAME : **Mrs. AAKRATI AGARWAL**
REFERRED BY : Dr.Nitin Agarwal (D M)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 31 Yrs.
SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
SERUM CREATININE	0.5	mg/dL.	0.5-1.4
URIC ACID	6.8	mg/dl	3.0-6.0

CLINICAL SIGNIFICANCE:

Analysis of synovial fluid plays a major role in the diagnosis of joint disease.

SERUM SODIUM (Na)	138	m Eq/litre.	135 - 155
SERUM POTASSIUM (K)	4.1	m Eq/litre.	3.5 - 5.5
SERUM CALCIUM	9.0	mg/dl	8.5 - 10.5

Report is not valid for medicolegal purpose

Reg.NO. : 111
 NAME : Mrs. AAKRATI AGARWAL
 REFERRED BY : Dr.Nitin Agarwal (D M)
 SAMPLE : BLOOD

DATE : 11/03/2023
 AGE : 31 Yrs.
 SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
LIVER PROFILE			
SERUM BILIRUBIN			
TOTAL	0.6	mg/dL	0.3-1.2
DIRECT	0.4	mg/dL	0.2-0.6
INDIRECT	0.2	mg/dL	0.1-0.4
SERUM PROTEINS			
Total Proteins	6.6	Gm/dL	6.4 - 8.3
Albumin	4.1	Gm/dL	3.5 - 5.5
Globulin	2.5	Gm/dL	2.3 - 3.5
A : G Ratio	1.64		0.0-2.0
SGOT	21	IU/L	0-40
SGPT	19	IU/L	0-40
SERUM ALK.PHOSPHATASE	67	IU/L	00-115

NORMAL RANGE : BILIRUBIN TOTAL

Premature infants. 0 to 1 day: <8 mg/dL Premature infants. 1 to 2 days: <12 mg/dL Adults: 0.3-1 mg/dL.
 Premature infants. 3 to 5 days: <16 mg/dL Neonates, 0 to 1 day: 1.4-8.7 mg/dL
 Neonates, 1 to 2 days: 3.4-11.5 mg/dL Neonates, 3 to 5 days: 1.5-12 mg/dL Children 6 days to 18 years: 0.3-1.2 mg/dL.

COMMENTS-

Total and direct bilirubin determination in serum is used for the diagnosis, differentiation and follow -up of jaundice. Elevation of SGPT is found in liver and kidney diseases such as infectious or toxic hepatitis, IM and cirrhosis. Organs rich in SGOT are heart, liver and skeletal muscles. When any of these organs are damaged, the serum SGOT level rises in proportion to the severity of damage. Elevation of Alkaline Phosphatase in serum or plasma is found in hepatitis, biliary obstructions, hyperparathyroidism, steatorrhea and bone diseases.

Report is not valid for medicolegal purpose

Reg.NO. : 111
NAME : **Mrs. AAKRATI AGARWAL**
REFERRED BY : Dr.Nitin Agarwal (D.M.)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 31 Yrs.
SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
LIPID PROFILE			
SERUM CHOLESTEROL	170	mg/dL.	130 - 200
SERUM TRIGLYCERIDE	97	mg/dl.	30 - 160
HDL CHOLESTEROL	49	mg/dL.	30-70
VLDL CHOLESTEROL	19.4	mg/dL.	15 - 40
LDL CHOLESTEROL	101.60	mg/dL.	00-130
CHOL/HDL CHOLESTEROL RATIO	3.47	mg/dl	
LDL/HDL CHOLESTEROL RATIO	2.07	mg/dl	

INTERPRETATION

TRIGLYCERIDE level > 250mg/dL is associated with an approximately 2-fold greater risk of coronary vascular disease. Elevation of triglycerides can be seen with obesity, medication, fast less than 12 hrs., alcohol intake, diabetes melitus, and pancreatitis.

CHOLESTEROL, its fractions and triglycerides are the important plasma lipids in defining cardiovascular risk factors and in the management of cardiovascular disease. Highest acceptable and optimum values of cholesterol values of cholesterol vary with age. Values above 220 mgm/dl are associated with increased risk of CHD regardless of HDL & LDL values.

HDL-CHOLESTEROL level <35 mg/dL is associated with an increased risk of coronary vascular disease even in the face of desirable levels of cholesterol and LDL - cholesterol.

LDL - CHOLESTEROL & TOTAL CHOLESTEROL levels can be strikingly altered by thyroid, renal and liver disease as well as hereditary factors. Based on total cholesterol, LDL- cholesterol, and total cholesterol/HDL - cholesterol ratio, patients may be divided into the three risk categories.

URINE EXAMINATION

Report is not valid for medicolegal purpose

Reg.NO. : 111
 NAME : Mrs. AAKRATI AGARWAL
 REFERRED BY : Dr.Nitin Agarwal (D M)
 SAMPLE : BLOOD

DATE : 11/03/2023
 AGE : 31 Yrs.
 SEX : FEMALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
URINE EXAMINATION REPORT			
PHYSICAL EXAMINATION			
pH	5.0		
TRANSPARENCY			
Volume	20	ml	
Colour	Light Yellow		
Appearance	Clear		Nil
Sediments	Nil		
Specific Gravity	1.020		1.015-1.025
Reaction	Acidic		
BIOCHEMICAL EXAMINATION			
UROBILINOGEN	Nil		NIL
BILIRUBIN	Nil		NEGATIVE
URINE KETONE	Nil		NEGATIVE
Sugar	Nil		Nil
Albumin	Nil		Nil
Phosphates	Absent		Nil
MICROSCOPIC EXAMINATION			
Red Blood Cells	Nil	/H.P.F.	
Pus Cells	1-2	/H.P.F.	
Epithelial Cells	2-3	/H.P.F.	
Crystals	NIL		NIL
Casts	NIL	/H.P.F.	
Bacteria	NIL		
Other	NIL		

Report is not valid for medicolegal purpose

Apple Cardiac Care
Ekta Nagar, Stadium Road,
Apple Care Hospital,
Bareilly - 243 122 (U.P.) India
Tel. : 07599031977, 09458888448



Reg.NO. : 111
NAME : **Mrs. AAKRATI AGARWAL**
REFERRED BY : Dr.Nitin Agarwal (D.M.)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 31 Yrs.
SEX : FEMALE

TEST NAME

RESULTS

UNITS

BIOLOGICAL REF. RANGE

--{End of Report}--

Dr. Shweta Agarwal
MD(Pathology), Apple Pathology
Bareilly (UP)



॥ ॐ गणेशाय नमः ॥

GANESH DIAGNOSTIC

DR. LOKESH GOYAL

MBBS (KMC), MD (RADIOLOGY)

CONSULTANT INTERVENTIONAL RADIOLOGIST
FORMER SR. REGISTRAR - APOLLO HOSPITAL, NEW DELHI
LIFE MEMBER OF IRIA

Timings : 9:00 am to 9:00 pm, Sunday 9.00 am to 3.00 pm ☎ 8392957683, 6395228718

MR. AAKRATI AGARWAL
DR. NITIN AGARWAL, DM

11-03-2023

REPORT

EXAMINATION PERFORMED: X-RAY CHEST

B/L lung fields are clear

Both of the CP angles are clear.

Both hila show a normal pattern .

Cardiac and mediastinal borders appear normal.

Visualized bony thorax and soft tissue of the chest wall appear normal.

IMPRESSION ---NO SIGNIFICANT ABNORMALITY IS SEEN

Not for medico-legal purpose

DR LOKESH GOYAL
MD
RADIODIAGNOSIS

डिजिटल एक्स-रे, मल्टी स्लाइस
सी. टी. स्कैन सुविधा उपलब्ध है।



NOT VALID FOR
MEDICO LEGAL PURPOSE